

catena-Poly[[pyridine- κ N)copper(II)]- μ_3 -pyridine-2,6-dicarboxylato- κ^3 O²:O^{2'},N,O⁶:O^{6'}]

Manoj Trivedi,^{a*}‡ Daya Shankar Pandey^a and Nigam P. Rath^b

^aDepartment of Chemistry, Faculty of Science, Banaras Hindu University, Varanasi 221005, India, and ^bDepartment of Chemistry and Biochemistry, and Centre for Nanoscience, University of Missouri-St Louis, One University Boulevard, St Louis, MO 63121-4499, USA

Correspondence e-mail: manojtri@gmail.com


Received 20 January 2009; accepted 13 February 2009

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.029; wR factor = 0.074; data-to-parameter ratio = 11.0.

In the title compound, $[\text{Cu}(\text{C}_7\text{H}_3\text{NO}_4)(\text{C}_5\text{H}_5\text{N})]_n$, the Cu^{II} atom is in a slightly distorted octahedral coordination environment. Each Cu^{II} atom is bound to two N atoms and one O atom of the pyridinedicarboxylate (PDA) ligand in a tridentate manner, one N atom of the pyridine molecule and two bridging carboxylate O atoms of adjacent PDA ligands, leading to a linear one-dimensional chain running along the c axis. These chains are further assembled *via* weak $\text{C}-\text{H}\cdots\text{O}$ and $\pi-\pi$ interactions into a three-dimensional supramolecular network structure. The centroid-centroid distance between the $\pi-\pi$ interacting pyridine rings is 3.9104 (13) Å. The two N atoms are *trans* to each other with respect to Cu.

Related literature

For background information on coordination polymers, see: Kitagawa *et al.* (2004); Kirillov *et al.* (2008); Hoskins & Robson (1990); Eddaoudi *et al.* (2001). For related polymeric structures of PDA complexes, see, for example: Zhao *et al.* (2003); Choi *et al.* (2003); Ghosh *et al.* (2004); Xie *et al.* (2004). For related structures of Cu complexes, see: Uçar *et al.* (2007); Manna *et al.* (2007); Gao *et al.* (2006).



Experimental

Crystal data

$[\text{Cu}(\text{C}_7\text{H}_3\text{NO}_4)(\text{C}_5\text{H}_5\text{N})]$	$V = 1069.2$ (2) Å ³
$M_r = 307.74$	$Z = 4$
Monoclinic, $C2/c$	Mo $K\alpha$ radiation
$a = 7.8042$ (9) Å	$\mu = 2.06$ mm ⁻¹
$b = 13.6152$ (17) Å	$T = 100$ K
$c = 10.0667$ (12) Å	$0.21 \times 0.13 \times 0.08$ mm
$\beta = 91.687$ (4)°	

Data collection

Bruker APEXII CCD area-detector diffractometer	3530 measured reflections
Absorption correction: multi-scan (SADABS; Bruker, 2007)	981 independent reflections
$T_{\min} = 0.671$, $T_{\max} = 0.848$	859 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.036$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.029$	89 parameters
$wR(F^2) = 0.074$	H-atom parameters not refined
$S = 1.10$	$\Delta\rho_{\text{max}} = 0.41$ e Å ⁻³
981 reflections	$\Delta\rho_{\text{min}} = -0.60$ e Å ⁻³

Table 1

Selected bond lengths (Å).

Cu1—N2	1.896 (3)	Cu1—O1	2.0110 (18)
Cu1—N1	1.944 (3)	Cu1—O1 ⁱ	2.0110 (18)

Symmetry code: (i) $-x + 1, y, -z + \frac{1}{2}$.

Table 2

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C3}-\text{H3}\cdots\text{O2}^{\text{ii}}$	0.95	2.44	3.187 (3)	135
$\text{C5}-\text{H5}\cdots\text{O1}^{\text{i}}$	0.95	2.48	3.070 (3)	120
$\text{C6}-\text{H6}\cdots\text{O1}^{\text{iii}}$	0.95	2.48	3.394 (3)	162

Symmetry codes: (i) $-x + 1, y, -z + \frac{1}{2}$; (ii) $-x + \frac{1}{2}, -y - \frac{1}{2}, -z$; (iii) $x + \frac{1}{2}, -y + \frac{1}{2}, z + \frac{1}{2}$.

‡ Present Address: Department of Chemistry, University of Delhi, Delhi 110007, India.

The authors gratefully acknowledge financial support by the Council of Scientific and Industrial Research, New Delhi, and

the University Grants Commission, New Delhi [grant No. F4-2/2006(BSR)/13-76/2008(BSR)]. The authors also thank Dr Rajamani Nagarajan and the Head, Department of Chemistry, University of Delhi, and the Head, Department of Chemistry, Faculty of Science, Banaras Hindu University, Varanasi, for their support. Funding from the National Science Foundation (grant No. CHE0420497) for the purchase of the APEXII diffractometer is acknowledged.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: IS2384).

References

- Bruker (2006). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2007). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Choi, K.-Y., Ryu, H., Lim, Y.-M., Sung, N.-D., Shin, U.-S. & Suh, M. (2003). *Inorg. Chem. Commun.* **6**, 412–415.
- Eddaoudi, M., Moler, D. B., Li, H., Chen, B., Reinecke, T. M., O'Keeffe, M. & Yaghi, O. M. (2001). *Acc. Chem. Res.* **34**, 319–330.
- Gao, H.-L., Yi, L., Zhao, B., Zhao, X.-Q., Cheng, P., Liao, D.-Z. & Yan, S.-P. (2006). *Inorg. Chem.* **45**, 5980–5988.
- Ghosh, S. K., Ribas, J. & Bharadwaj, P. K. (2004). *CrystEngComm*, **6**, 250–256.
- Hoskins, B. F. & Robson, R. (1990). *J. Am. Chem. Soc.* **112**, 1546–1554.
- Kirillov, A. M., Karabach, Y. Y., Haukka, M., Guedes da Silva, M. F. C., Sanchiz, J., Kopylovich, M. N. & Pombeiro, A. J. L. (2008). *Inorg. Chem.* **47**, 162–175.
- Kitagawa, S., Kitaura, R. & Noro, S. (2004). *Angew. Chem. Int. Ed.* **43**, 2334–2375.
- Manna, S. C., Ribas, J., Zangrando, E. & Chaudhuri, N. R. (2007). *Inorg. Chim. Acta*, **360**, 2589–2597.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Uçar, İ., Karabulut, B., Bulut, A. & Büyükgüngör, O. (2007). *J. Mol. Struct.* **834–836**, 336–344.
- Xie, C., Zhang, Z., Wang, X., Liu, X., Shen, G., Wang, R. & Shen, D. (2004). *J. Coord. Chem.* **57**, 1173–1178.

supplementary materials

Acta Cryst. (2009). E65, m303-m304 [doi:10.1107/S1600536809005212]

***catena*-Poly[[*(pyridine-κN)*copper(II)]-*μ*₃-*pyridine-2,6-dicarboxylato-κ³O²:O^{2'},N,O⁶:O^{6'}*]**

M. Trivedi, D. S. Pandey and N. P. Rath

Comment

The rapidly expanding field of the crystal engineering (the design of crystalline materials) of polymeric coordination networks stems has recently attracted great interest because of their potential applications as zeolite-like materials for molecular selection, ion exchange, and catalysis, as well as in the variety of architectures and topologies (Kitagawa *et al.*, 2004; Kirillov *et al.*, 2008). The main strategy popularly used in this area is a building-block approach (Hoskins & Robson, 1990; Eddaoudi *et al.*, 2001). 2,6-Pyridinedicarboxylic acid (H₂PDA) is an efficient ligand. Polymeric structure of PDA complexes with transition and lanthanide metals have been reported, in which PDA not only chelates but also bridges to form diversified structures with three coordination sites (Zhao *et al.*, 2003; Choi *et al.*, 2003; Ghosh *et al.*, 2004; Xie *et al.*, 2004). We report the synthesis, and crystal structures of one compound, [Cu(*μ*-2,6-PDA)(py)]_n (1).

Molecular structure of (1) shows a slightly distorted octahedral coordination geometry. The equatorial sites are occupied by an NO₂ donor from the carboxylate groups at the pyridine-2,6-position of PDA (N2, O1, O1ⁱ) and one N atom from pyridine (N1). Two O atoms from two other neighboring PDA ligands occupy the axial sites (O2, O2ⁱ) at a distance of 2.761 Å (Fig. 1). The equatorial Cu—O and Cu—N bond lengths of are normal [Cu1—O1 = 2.0110 (18) Å, Cu1—O1ⁱ = 2.0110 (18) Å, Cu1—N2 = 1.896 (3) Å, Cu1—N1 = 1.944 (3) Å], which are within ranges reported in other copper complexes (Uçar *et al.*, 2007; Manna *et al.*, 2007; Gao *et al.*, 2006). The pyridine is essentially planar with no deviation from planarity for pyridyl N1-atom. The C—C—C angles about the pyridyl ring are 118.2 (3) to 128.5 (2)°, indicating *sp*² hybridization. Two carboxylate O atoms (O2 and O2ⁱ) which are coordinated to the adjacent copper atom, have C—O distances [O2—C1 = 1.228 (3) Å, O1—C1 = 1.290 (3) Å] which are generally shorter than C—O distances, indicating the conjugation of the double bond after deprotonation. In this way, the PDA ligands bridge adjacent Cu atoms to form a [Cu(*μ*-2,6-PDC)(py)]_n linear chains extending in the [001] direction (Fig. 2). The separations between the two Cu atom in the linear chains are 5.332 Å. The PDA ligand and pyridine are *trans* to each other (N2—Cu1—N1 = 180°). However, one-dimensional polymeric chains are connected in the solid state through weak C—H⋯O and π-π interactions. Weak C—H⋯O interactions that connects polymeric chains into two-dimensional network (Fig. 3). Contact distances for C—H⋯O interactions are 2.43–2.76 Å (Table 1). The weak π-π interactions are present in (1). Further, the importance of π-π stacking interactions between aromatic rings has widely been recognized in the intercalation of drugs with DNA especially in biological systems, which lie in the range 3.4–3.5 Å. The complex (1) exhibits intermolecular face-to-face π-π interactions [π-pyridyl/π-pyridyl ct/ct distance 3.9104 (13) Å; Fig. 4].

Experimental

A mixture of [Cu(NO₃)₂·6H₂O] (0.466 g, 2 mmol), H₂PDA (0.167 g, 1 mmol), 2-Pyridine thiol(0.111 g, 1 mmol) was dissolved in a mixture of MeOH (5 ml) and water (5 mL) and add pyridine (in excess). The solution was stirred for 24 h at room temperature. Slowly, color of the solution changes from blue to dark green. The resulting solution was filtered and left

supplementary materials

at room temperature for two days, which resulted in blue needle crystals which are suitable for X-ray diffraction analysis (yield 0.184 g, 60%). Anal. Calc. for $C_{12}H_8N_2O_4Cu$: C 48.83, H 2.62, N 9.10%; found: C 47.65, H 2.56, N 9.30%.

Refinement

All H atoms were added in their calculated positions ($C-H = 0.95 \text{ \AA}$) and were treated using appropriate riding models, with $U_{iso}(H) = 1.2U_{eq}(C)$.

Figures




Fig. 1. A view of the structure of (1), showing the atom-numbering scheme and the Cu coordination octahedra; displacement ellipsoids are drawn at the 50% probability level.



Fig. 2. Packing view of (1), showing linear chains along the [001] direction.



Fig. 3. Packing view of (1), showing connectivity with other polymeric chains through weak $C-H \cdots O$ hydrogen bond interactions.



Fig. 4. Packing view of (1), showing face-to-face $\pi-\pi$ interactions.

catena-Poly[[*(pyridine- κ N)*copper(II)]- μ_3 -*pyridine-2,6-dicarboxylato- $\kappa^3 O^2:O^2',N,O^6:O^6'$*]

Crystal data

$[Cu(C_7H_3NO_4)(C_5H_5N)]$

$M_r = 307.74$

Monoclinic, $C2/c$

Hall symbol: $-C 2yc$

$a = 7.8042 (9) \text{ \AA}$

$b = 13.6152 (17) \text{ \AA}$

$c = 10.0667 (12) \text{ \AA}$

$\beta = 91.687 (4)^\circ$

$V = 1069.2 (2) \text{ \AA}^3$

$Z = 4$

$F_{000} = 620$

$D_x = 1.912 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation

$\lambda = 0.71073 \text{ \AA}$

Cell parameters from 1607 reflections

$\theta = 3.0-25.3^\circ$

$\mu = 2.06 \text{ mm}^{-1}$

$T = 100 \text{ K}$

Needle, blue

$0.21 \times 0.13 \times 0.08 \text{ mm}$

Data collection

Bruker APEXII CCD area-detector diffractometer	981 independent reflections
Radiation source: fine-focus sealed tube	859 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.036$
$T = 100$ K	$\theta_{\text{max}} = 25.3^\circ$
φ and ω scans	$\theta_{\text{min}} = 3.0^\circ$
Absorption correction: multi-scan (SADABS; Bruker, 2007)	$h = -9 \rightarrow 8$
$T_{\text{min}} = 0.671$, $T_{\text{max}} = 0.848$	$k = -16 \rightarrow 16$
3530 measured reflections	$l = -12 \rightarrow 11$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.029$	H-atom parameters not refined
$wR(F^2) = 0.074$	$w = 1/[\sigma^2(F_o^2) + (0.0347P)^2 + 1.5139P]$
$S = 1.10$	where $P = (F_o^2 + 2F_c^2)/3$
981 reflections	$(\Delta/\sigma)_{\text{max}} < 0.001$
89 parameters	$\Delta\rho_{\text{max}} = 0.41 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	$\Delta\rho_{\text{min}} = -0.60 \text{ e } \text{\AA}^{-3}$
	Extinction correction: none

Special details

Experimental. All H atoms were added in their calculated positions and were treated using appropriate riding models.

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Cu1	0.5000	0.06463 (3)	0.2500	0.01055 (18)
O1	0.3461 (2)	0.04164 (13)	0.08928 (18)	0.0119 (4)
O2	0.2652 (2)	-0.07906 (13)	-0.05100 (19)	0.0137 (4)
N1	0.5000	0.2074 (2)	0.2500	0.0099 (7)

supplementary materials

N2	0.5000	-0.0746 (2)	0.2500	0.0096 (7)
C1	0.3334 (3)	-0.04888 (19)	0.0526 (3)	0.0111 (6)
C2	0.4161 (3)	-0.1210 (2)	0.1514 (3)	0.0100 (6)
C3	0.4135 (3)	-0.2221 (2)	0.1484 (3)	0.0120 (6)
H3	0.3542	-0.2564	0.0789	0.014*
C4	0.5000	-0.2728 (3)	0.2500	0.0130 (8)
H4	0.5000	-0.3426	0.2500	0.016*
C5	0.5748 (3)	0.2583 (2)	0.3507 (3)	0.0124 (6)
H5	0.6286	0.2230	0.4218	0.015*
C6	0.5765 (4)	0.3594 (2)	0.3549 (3)	0.0154 (6)
H6	0.6290	0.3931	0.4281	0.018*
C7	0.5000	0.4112 (3)	0.2500	0.0159 (9)
H7	0.5000	0.4810	0.2500	0.019*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cu1	0.0144 (3)	0.0059 (3)	0.0109 (3)	0.000	-0.00649 (18)	0.000
O1	0.0151 (10)	0.0074 (10)	0.0129 (10)	-0.0014 (8)	-0.0066 (8)	0.0000 (8)
O2	0.0173 (10)	0.0117 (11)	0.0118 (10)	-0.0026 (8)	-0.0068 (8)	-0.0017 (8)
N1	0.0095 (16)	0.0095 (17)	0.0108 (17)	0.000	-0.0009 (13)	0.000
N2	0.0085 (16)	0.0102 (17)	0.0100 (16)	0.000	-0.0009 (13)	0.000
C1	0.0114 (14)	0.0106 (14)	0.0114 (15)	-0.0017 (11)	-0.0007 (11)	0.0004 (11)
C2	0.0080 (13)	0.0125 (15)	0.0095 (14)	-0.0020 (11)	-0.0011 (11)	-0.0023 (11)
C3	0.0122 (14)	0.0131 (15)	0.0108 (14)	-0.0006 (11)	-0.0019 (11)	-0.0018 (11)
C4	0.014 (2)	0.009 (2)	0.016 (2)	0.000	0.0001 (16)	0.000
C5	0.0124 (14)	0.0138 (15)	0.0110 (15)	0.0001 (11)	-0.0012 (11)	0.0007 (11)
C6	0.0163 (15)	0.0136 (15)	0.0163 (16)	-0.0035 (12)	0.0025 (12)	-0.0042 (12)
C7	0.016 (2)	0.009 (2)	0.023 (2)	0.000	0.0055 (17)	0.000

Geometric parameters (\AA , $^\circ$)

Cu1—N2	1.896 (3)	C2—C3	1.378 (4)
Cu1—N1	1.944 (3)	C3—C4	1.392 (3)
Cu1—O1	2.0110 (18)	C3—H3	0.9500
Cu1—O1 ⁱ	2.0110 (18)	C4—C3 ⁱ	1.392 (3)
O1—C1	1.290 (3)	C4—H4	0.9500
O2—C1	1.228 (3)	C5—C6	1.378 (4)
N1—C5 ⁱ	1.347 (3)	C5—H5	0.9500
N1—C5	1.347 (3)	C6—C7	1.390 (3)
N2—C2	1.332 (3)	C6—H6	0.9500
N2—C2 ⁱ	1.332 (3)	C7—C6 ⁱ	1.390 (3)
C1—C2	1.527 (4)	C7—H7	0.9500
N2—Cu1—N1	180.0	N2—C2—C1	111.7 (2)
N2—Cu1—O1	81.05 (5)	C3—C2—C1	128.5 (2)
N1—Cu1—O1	98.95 (5)	C2—C3—C4	118.2 (3)
N2—Cu1—O1 ⁱ	81.05 (5)	C2—C3—H3	120.9
N1—Cu1—O1 ⁱ	98.95 (5)	C4—C3—H3	120.9

O1—Cu1—O1 ⁱ	162.10 (10)	C3—C4—C3 ⁱ	120.6 (4)
C1—O1—Cu1	114.71 (16)	C3—C4—H4	119.7
C5 ⁱ —N1—C5	118.1 (3)	C3 ⁱ —C4—H4	119.7
C5 ⁱ —N1—Cu1	120.97 (16)	N1—C5—C6	122.8 (3)
C5—N1—Cu1	120.96 (16)	N1—C5—H5	118.6
C2—N2—C2 ⁱ	123.4 (3)	C6—C5—H5	118.6
C2—N2—Cu1	118.28 (16)	C5—C6—C7	118.7 (3)
C2 ⁱ —N2—Cu1	118.28 (16)	C5—C6—H6	120.7
O2—C1—O1	126.2 (2)	C7—C6—H6	120.7
O2—C1—C2	120.2 (2)	C6 ⁱ —C7—C6	119.0 (4)
O1—C1—C2	113.6 (2)	C6 ⁱ —C7—H7	120.5
N2—C2—C3	119.8 (3)	C6—C7—H7	120.5
N2—Cu1—O1—C1	-6.98 (18)	Cu1—N2—C2—C3	-179.96 (18)
N1—Cu1—O1—C1	173.02 (18)	C2 ⁱ —N2—C2—C1	-179.7 (2)
O1 ⁱ —Cu1—O1—C1	-6.98 (18)	Cu1—N2—C2—C1	0.3 (2)
O1—Cu1—N1—C5 ⁱ	-6.84 (14)	O2—C1—C2—N2	172.9 (2)
O1 ⁱ —Cu1—N1—C5 ⁱ	173.16 (14)	O1—C1—C2—N2	-6.2 (3)
O1—Cu1—N1—C5	173.16 (14)	O2—C1—C2—C3	-6.8 (4)
O1 ⁱ —Cu1—N1—C5	-6.84 (14)	O1—C1—C2—C3	174.1 (3)
O1—Cu1—N2—C2	3.29 (14)	N2—C2—C3—C4	-0.1 (4)
O1 ⁱ —Cu1—N2—C2	-176.71 (14)	C1—C2—C3—C4	179.6 (2)
O1—Cu1—N2—C2 ⁱ	-176.71 (14)	C2—C3—C4—C3 ⁱ	0.03 (18)
O1 ⁱ —Cu1—N2—C2 ⁱ	3.29 (14)	C5 ⁱ —N1—C5—C6	0.47 (19)
Cu1—O1—C1—O2	-170.2 (2)	Cu1—N1—C5—C6	-179.53 (19)
Cu1—O1—C1—C2	8.8 (3)	N1—C5—C6—C7	-0.9 (4)
C2 ⁱ —N2—C2—C3	0.04 (18)	C5—C6—C7—C6 ⁱ	0.44 (18)

Symmetry codes: (i) $-x+1, y, -z+1/2$.

Hydrogen-bond geometry ($\text{\AA}, ^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
C3—H3 \cdots O2 ⁱⁱ	0.95	2.44	3.187 (3)	135
C5—H5 \cdots O1 ⁱ	0.95	2.48	3.070 (3)	120
C6—H6 \cdots O1 ⁱⁱⁱ	0.95	2.48	3.394 (3)	162

Symmetry codes: (ii) $-x+1/2, -y-1/2, -z$; (i) $-x+1, y, -z+1/2$; (iii) $x+1/2, -y+1/2, z+1/2$.

Fig. 1




Fig. 2




Fig. 3




Fig. 4

