

4-Chloro-2-methyl-N-phenylbenzene-sulfonamide

B. Thimme Gowda,^{a*} Sabine Foro,^b P. G. Nirmala,^a K. S. Babitha^a and Hartmut Fuess^b

^aDepartment of Chemistry, Mangalore University, Mangalagangotri 574 199, Mangalore, India, and ^bInstitute of Materials Science, Darmstadt University of Technology, Petersenstrasse 23, D-64287 Darmstadt, Germany
Correspondence e-mail: gowdabt@yahoo.com


Received 17 January 2009; accepted 2 February 2009

Key indicators: single-crystal X-ray study; $T = 299$ K; mean $\sigma(\text{C}-\text{C}) = 0.007 \text{ \AA}$; R factor = 0.057; wR factor = 0.176; data-to-parameter ratio = 14.4.

There are two molecules in the asymmetric unit of the title compound, $C_{13}H_{12}ClNO_2S$, with similar conformations. The orientations of the *ortho*-methyl groups in the sulfonyl benzene rings are in the direction of the N—H bonds of the sulfonamide groups. In the crystal, the molecules are each linked into centrosymmetric dimers through N—H···O hydrogen bonds and packed into a layered structure diagonally in the *bc* plane.

Related literature

For related structures, see: Gelbrich *et al.* (2007); Gowda *et al.* (2008a,b, 2009); Perlovich *et al.* (2006)



Experimental

Crystal data

$C_{13}H_{12}ClNO_2S$	$\gamma = 105.79(1)^\circ$
$M_r = 281.75$	$V = 1341.6(3) \text{ \AA}^3$
Triclinic, $P\bar{1}$	$Z = 4$
$a = 8.609(1) \text{ \AA}$	Cu $K\alpha$ radiation
$b = 11.143(1) \text{ \AA}$	$\mu = 3.93 \text{ mm}^{-1}$
$c = 14.726(2) \text{ \AA}$	$T = 299(2) \text{ K}$
$\alpha = 98.618(7)^\circ$	$0.33 \times 0.23 \times 0.08 \text{ mm}$
$\beta = 90.951(8)^\circ$	

Data collection

Enraf–Nonius CAD-4 diffractometer
Absorption correction: ψ scan (North *et al.*, 1968)
 $T_{\min} = 0.350$, $T_{\max} = 0.729$
7620 measured reflections

4784 independent reflections
2980 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.045$
3 standard reflections
frequency: 120 min
intensity decay: 1.0%

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.057$
 $wR(F^2) = 0.176$
 $S = 1.03$
4784 reflections
333 parameters
12 restraints

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.41 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.42 \text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N1—H1N···O2 ⁱ	0.91 (5)	2.02 (5)	2.922 (4)	175 (4)
N2—H2N···O3 ⁱⁱ	0.88 (4)	2.03 (5)	2.906 (4)	173 (4)

Symmetry codes: (i) $-x + 1, -y + 2, -z$; (ii) $-x, -y + 1, -z + 1$.

Data collection: *CAD-4-PC* (Enraf–Nonius, 1996); cell refinement: *CAD-4-PC*; data reduction: *REDU4* (Stoe & Cie, 1987); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2003); software used to prepare material for publication: *SHELXL97*.

BTG thanks the Alexander von Humboldt Foundation, Bonn, Germany, for extensions of his research fellowship.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: FJ2192).

References

- Enraf–Nonius (1996). *CAD-4-PC*. Enraf–Nonius, Delft, The Netherlands.
- Gelbrich, T., Hursthouse, M. B. & Threlfall, T. L. (2007). *Acta Cryst. B* **63**, 621–632.
- Gowda, B. T., Foro, S., Babitha, K. S. & Fuess, H. (2008a). *Acta Cryst. E* **64**, o1692.
- Gowda, B. T., Foro, S., Babitha, K. S. & Fuess, H. (2008b). *Acta Cryst. E* **64**, o1825.
- Gowda, B. T., Foro, S., Nirmala, P. G., Babitha, K. S. & Fuess, H. (2009). *Acta Cryst. E* **65**. Submitted.
- North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst. A* **24**, 351–359.
- Perlovich, G. L., Tkachev, V. V., Schaper, K.-J. & Raevsky, O. A. (2006). *Acta Cryst. E* **62**, o780–o782.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2003). *J. Appl. Cryst. 36*, 7–13.
- Stoe & Cie (1987). *REDU4*. Stoe & Cie GmbH, Darmstadt, Germany.

supplementary materials

Acta Cryst. (2009). E65, o476 [doi:10.1107/S1600536809003845]

4-Chloro-2-methyl-N-phenylbenzenesulfonamide

B. T. Gowda, S. Foro, P. G. Nirmala, K. S. Babitha and H. Fuess

Comment

In the present work, as part of a study of substituent effects on the structures of *N*-(aryl)-arylsulfonamides, the structure of *N*-(phenyl)-2-methyl-4-chlorobenzenesulfonamide (NP2M4CBSA) has been determined (Gowda *et al.* 2008*a,b*, 2009). The asymmetric unit of NP2M4CBSA contains 2 molecules. The orientations of the *ortho*- methyl groups in the sulfonyl benzene rings are in the direction of the N—H bonds of the sulfonamido groups (Fig. 1). The opposite signs of the C—S—N—C torsion angles in the two independent molecules, -61.9 (4) $^{\circ}$ (molecule 1) and 69.7 (4) $^{\circ}$ (molecule 2), indicates that they have opposite chirality, although the choice of the chirality of the second molecule relative to the first may be arbitrary. The two benzene rings in NP2M4CBSA are tilted relative to each other by 86.6 (2) $^{\circ}$ in the molecule 1 and 83.0 (2) $^{\circ}$ in molecule 2, compared with the values of 67.5 (1) $^{\circ}$ (molecule 1) and 72.9 (1) $^{\circ}$ (molecule 2) for *N*-(phenyl)-2,4-dimethylbenzenesulfonamide (NP24DMBSA) (Gowda *et al.*, 2009). The other bond parameters in NP2M4CBSA are similar to those observed for *N*-(2-methylphenyl)-benzenesulfonamide (Gowda *et al.*, 2008*a*), NP24DMBSA and other aryl sulfonamides (Perlovich *et al.*, 2006; Gelbrich *et al.*, 2007; Gowda *et al.*, 2008*b*). The crystal packing of molecules in NP2M4CBSA *via* intermolecular N—H···O hydrogen bonds (Table 1) is shown in Fig.2.

Experimental

The solution of *m*-chlorotoluene (10 cc) in chloroform (40 cc) was treated dropwise with chlorosulfonic acid (25 cc) at 0 °C. After the initial evolution of hydrogen chloride subsided, the reaction mixture was brought to room temperature and poured into crushed ice in a beaker. The chloroform layer was separated, washed with cold water and allowed to evaporate slowly. The residual 2-methyl-4-chlorobenzenesulfonylchloride was treated with aniline in the stoichiometric ratio and boiled for ten minutes. The reaction mixture was then cooled to room temperature and added to ice cold water (100 cc). The resultant solid *N*-(phenyl)-2-methyl-4-chlorobenzenesulfonamide was filtered under suction and washed thoroughly with cold water. It was then recrystallized to constant melting point from dilute ethanol. The purity of the compound was checked and characterized by recording its infrared and NMR spectra. The single crystals used in X-ray diffraction studies were grown in ethanolic solution by slow evaporation at room temperature.

Refinement

The H atoms of the NH groups were located in a difference map and their positions refined, with N—H = 0.88 (4)–0.91 (5) Å. The carbon-bound H atoms were positioned with idealized geometry and refined using a riding model, with C—H distances 0.93–0.96 Å. All H atoms were refined with isotropic displacement parameters (set to 1.2 times of the U_{eq} of the parent atom). For methyl group $U_{\text{iso}}(\text{H}) = 1.5 U_{\text{eq}}$.

supplementary materials

Figures



Fig. 1. Molecular structure of the title compound, showing the atom labeling scheme. The displacement ellipsoids are drawn at the 50% probability level. The H atoms are represented as small spheres of arbitrary radii.




Fig. 2. Molecular packing of the title compound with hydrogen bonding shown as dashed lines.

4-Chloro-2-methyl-N-phenylbenzenesulfonamide

Crystal data

$C_{13}H_{12}ClNO_2S$	$Z = 4$
$M_r = 281.75$	$F_{000} = 584$
Triclinic, $P\bar{1}$	$D_x = 1.395 \text{ Mg m}^{-3}$
Hall symbol: -P 1	$\text{Cu } K\alpha \text{ radiation}$
$a = 8.609 (1) \text{ \AA}$	$\lambda = 1.54180 \text{ \AA}$
$b = 11.143 (1) \text{ \AA}$	Cell parameters from 25 reflections
$c = 14.726 (2) \text{ \AA}$	$\theta = 5.6\text{--}21.8^\circ$
$\alpha = 98.618 (7)^\circ$	$\mu = 3.93 \text{ mm}^{-1}$
$\beta = 90.951 (8)^\circ$	$T = 299 \text{ K}$
$\gamma = 105.79 (1)^\circ$	Prism, colourless
$V = 1341.6 (3) \text{ \AA}^3$	$0.33 \times 0.23 \times 0.08 \text{ mm}$

Data collection

Enraf–Nonius CAD-4 diffractometer	$R_{\text{int}} = 0.045$
Radiation source: fine-focus sealed tube	$\theta_{\text{max}} = 67.0^\circ$
Monochromator: graphite	$\theta_{\text{min}} = 3.0^\circ$
$T = 299 \text{ K}$	$h = -10 \rightarrow 10$
$\omega/2\theta$ scans	$k = -13 \rightarrow 13$
Absorption correction: ψ scan (North <i>et al.</i> , 1968)	$l = -17 \rightarrow 7$
$T_{\text{min}} = 0.350$, $T_{\text{max}} = 0.729$	3 standard reflections
7620 measured reflections	every 120 min
4784 independent reflections	intensity decay: 1.0%
2980 reflections with $I > 2\sigma(I)$	

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.057$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.176$	$w = 1/[\sigma^2(F_o^2) + (0.0962P)^2 + 0.1233P]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.03$	$(\Delta/\sigma)_{\max} = 0.005$
4784 reflections	$\Delta\rho_{\max} = 0.41 \text{ e \AA}^{-3}$
333 parameters	$\Delta\rho_{\min} = -0.42 \text{ e \AA}^{-3}$
12 restraints	Extinction correction: none
Primary atom site location: structure-invariant direct methods	

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C11	0.4618 (4)	0.43860 (18)	0.24063 (15)	0.1670 (10)
S1	0.46343 (11)	0.96896 (11)	0.15504 (7)	0.0586 (3)
O1	0.4383 (4)	1.0364 (3)	0.2403 (2)	0.0802 (9)
O2	0.3532 (3)	0.9562 (3)	0.07717 (19)	0.0722 (8)
N1	0.6377 (4)	1.0345 (3)	0.1199 (2)	0.0611 (9)
H1N	0.647 (5)	1.039 (4)	0.059 (3)	0.073*
C1	0.4664 (4)	0.8182 (4)	0.1749 (3)	0.0565 (10)
C2	0.4758 (8)	0.7240 (5)	0.1044 (3)	0.0907 (16)
C3	0.4778 (10)	0.6088 (5)	0.1273 (4)	0.120 (2)
H3	0.4876	0.5444	0.0816	0.144*
C4	0.4653 (7)	0.5877 (5)	0.2179 (4)	0.0910 (16)
C5	0.4560 (6)	0.6791 (5)	0.2862 (3)	0.0768 (13)
H5	0.4494	0.6642	0.3467	0.092*
C6	0.4563 (5)	0.7946 (5)	0.2653 (3)	0.0663 (11)
H6	0.4495	0.8586	0.3122	0.080*
C7	0.7894 (4)	1.0635 (3)	0.1698 (3)	0.0522 (9)

supplementary materials

C8	0.9247 (5)	1.0883 (4)	0.1201 (3)	0.0701 (12)
H8	0.9151	1.0848	0.0567	0.084*
C9	1.0763 (6)	1.1187 (5)	0.1656 (4)	0.0865 (15)
H9	1.1685	1.1358	0.1323	0.104*
C10	1.0914 (6)	1.1239 (5)	0.2591 (4)	0.0834 (15)
H10	1.1933	1.1440	0.2891	0.100*
C11	0.9570 (6)	1.0995 (4)	0.3074 (3)	0.0718 (12)
H11	0.9671	1.1026	0.3707	0.086*
C12	0.8054 (5)	1.0701 (4)	0.2639 (3)	0.0653 (11)
H12	0.7141	1.0547	0.2980	0.078*
C13	0.4894 (9)	0.7419 (4)	0.0007 (3)	0.102 (2)
H13A	0.5834	0.8097	-0.0050	0.123*
H13B	0.3947	0.7616	-0.0205	0.123*
H13C	0.4985	0.6653	-0.0357	0.123*
Cl2	0.1160 (4)	0.3570 (3)	0.99174 (14)	0.1919 (12)
S2	-0.06931 (11)	0.32374 (9)	0.57752 (8)	0.0560 (3)
O3	-0.1468 (3)	0.4185 (3)	0.5635 (2)	0.0703 (8)
O4	-0.1553 (3)	0.1953 (3)	0.5486 (2)	0.0727 (8)
N2	0.0946 (4)	0.3586 (3)	0.5248 (2)	0.0581 (9)
H2N	0.114 (5)	0.431 (4)	0.503 (3)	0.070*
C14	-0.0126 (5)	0.3404 (4)	0.6948 (3)	0.0569 (10)
C15	0.0592 (7)	0.4573 (4)	0.7478 (4)	0.0803 (13)
C16	0.0983 (8)	0.4591 (6)	0.8395 (4)	0.115 (2)
H16	0.1457	0.5359	0.8770	0.138*
C17	0.0679 (8)	0.3489 (7)	0.8762 (4)	0.110 (2)
C18	-0.0041 (7)	0.2348 (6)	0.8243 (4)	0.0906 (16)
H18	-0.0267	0.1612	0.8502	0.109*
C19	-0.0424 (5)	0.2309 (4)	0.7334 (3)	0.0679 (11)
H19	-0.0894	0.1533	0.6967	0.081*
C20	0.2114 (4)	0.2902 (3)	0.5100 (2)	0.0477 (8)
C21	0.1911 (5)	0.1719 (4)	0.5338 (3)	0.0610 (10)
H21	0.0998	0.1346	0.5632	0.073*
C22	0.3085 (6)	0.1093 (4)	0.5135 (3)	0.0692 (12)
H22	0.2954	0.0292	0.5286	0.083*
C23	0.4443 (6)	0.1657 (5)	0.4708 (3)	0.0742 (13)
H23	0.5231	0.1238	0.4574	0.089*
C24	0.4630 (5)	0.2827 (5)	0.4484 (3)	0.0723 (12)
H24	0.5551	0.3209	0.4201	0.087*
C25	0.3464 (4)	0.3451 (4)	0.4674 (3)	0.0577 (10)
H25	0.3596	0.4247	0.4513	0.069*
C26	0.0961 (9)	0.5821 (4)	0.7100 (4)	0.111 (2)
H26A	0.1530	0.5749	0.6548	0.133*
H26B	0.1619	0.6486	0.7550	0.133*
H26C	-0.0033	0.6010	0.6965	0.133*

Atomic displacement parameters (\AA^2)

U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
----------	----------	----------	----------	----------	----------

Cl1	0.280 (3)	0.1079 (13)	0.1210 (15)	0.0456 (15)	0.0296 (16)	0.0575 (11)
S1	0.0529 (5)	0.0868 (7)	0.0417 (5)	0.0282 (5)	0.0004 (4)	0.0114 (5)
O1	0.092 (2)	0.114 (3)	0.0494 (17)	0.0576 (19)	0.0072 (15)	0.0056 (16)
O2	0.0544 (16)	0.121 (3)	0.0516 (17)	0.0361 (16)	-0.0077 (13)	0.0244 (16)
N1	0.0573 (19)	0.085 (2)	0.0421 (18)	0.0178 (17)	-0.0045 (15)	0.0176 (17)
C1	0.052 (2)	0.076 (3)	0.039 (2)	0.0125 (18)	0.0021 (16)	0.0133 (18)
C2	0.146 (5)	0.079 (3)	0.047 (3)	0.030 (3)	0.012 (3)	0.017 (2)
C3	0.230 (8)	0.074 (3)	0.057 (3)	0.036 (4)	0.021 (4)	0.018 (3)
C4	0.123 (4)	0.081 (3)	0.066 (3)	0.017 (3)	0.006 (3)	0.025 (3)
C5	0.082 (3)	0.098 (4)	0.053 (3)	0.018 (3)	0.012 (2)	0.030 (3)
C6	0.059 (2)	0.091 (3)	0.048 (2)	0.015 (2)	0.0018 (19)	0.018 (2)
C7	0.057 (2)	0.051 (2)	0.049 (2)	0.0156 (17)	-0.0071 (17)	0.0100 (16)
C8	0.062 (3)	0.088 (3)	0.062 (3)	0.026 (2)	0.003 (2)	0.009 (2)
C9	0.058 (3)	0.112 (4)	0.089 (4)	0.024 (3)	0.002 (3)	0.012 (3)
C10	0.070 (3)	0.080 (3)	0.096 (4)	0.025 (2)	-0.029 (3)	-0.001 (3)
C11	0.081 (3)	0.063 (3)	0.065 (3)	0.012 (2)	-0.025 (2)	0.010 (2)
C12	0.065 (3)	0.074 (3)	0.050 (2)	0.007 (2)	-0.0075 (19)	0.015 (2)
C13	0.226 (7)	0.068 (3)	0.026 (2)	0.061 (4)	0.018 (3)	0.0116 (19)
Cl2	0.244 (3)	0.239 (3)	0.0830 (12)	0.035 (2)	-0.0157 (15)	0.0602 (15)
S2	0.0490 (5)	0.0529 (5)	0.0731 (7)	0.0173 (4)	0.0041 (4)	0.0259 (5)
O3	0.0608 (16)	0.0743 (19)	0.095 (2)	0.0364 (14)	0.0125 (15)	0.0398 (16)
O4	0.0641 (17)	0.0545 (17)	0.097 (2)	0.0071 (13)	-0.0088 (16)	0.0213 (15)
N2	0.0595 (19)	0.0517 (19)	0.078 (2)	0.0282 (16)	0.0181 (17)	0.0319 (17)
C14	0.057 (2)	0.055 (2)	0.066 (3)	0.0187 (18)	0.0156 (19)	0.0258 (19)
C15	0.103 (4)	0.063 (3)	0.073 (3)	0.019 (3)	0.007 (3)	0.015 (2)
C16	0.152 (6)	0.096 (4)	0.081 (4)	0.010 (4)	0.001 (4)	0.009 (3)
C17	0.131 (5)	0.131 (6)	0.071 (4)	0.027 (4)	0.003 (4)	0.041 (4)
C18	0.093 (4)	0.103 (4)	0.089 (4)	0.029 (3)	0.011 (3)	0.053 (3)
C19	0.070 (3)	0.063 (3)	0.079 (3)	0.021 (2)	0.007 (2)	0.032 (2)
C20	0.053 (2)	0.0466 (19)	0.046 (2)	0.0183 (16)	-0.0057 (16)	0.0085 (15)
C21	0.062 (2)	0.057 (2)	0.070 (3)	0.0214 (19)	0.001 (2)	0.019 (2)
C22	0.081 (3)	0.057 (2)	0.076 (3)	0.032 (2)	-0.010 (2)	0.008 (2)
C23	0.076 (3)	0.089 (3)	0.070 (3)	0.051 (3)	-0.004 (2)	0.001 (2)
C24	0.063 (3)	0.087 (3)	0.078 (3)	0.032 (2)	0.015 (2)	0.024 (3)
C25	0.056 (2)	0.061 (2)	0.058 (2)	0.0166 (18)	-0.0001 (18)	0.0164 (19)
C26	0.171 (6)	0.045 (3)	0.104 (4)	0.007 (3)	-0.003 (4)	0.010 (3)

Geometric parameters (\AA , $^\circ$)

Cl1—C4	1.735 (6)	Cl2—C17	1.727 (6)
S1—O1	1.415 (3)	S2—O4	1.417 (3)
S1—O2	1.441 (3)	S2—O3	1.431 (3)
S1—N1	1.611 (4)	S2—N2	1.604 (3)
S1—C1	1.754 (4)	S2—C14	1.756 (4)
N1—C7	1.419 (5)	N2—C20	1.419 (4)
N1—H1N	0.91 (5)	N2—H2N	0.88 (4)
C1—C2	1.382 (6)	C14—C19	1.385 (5)
C1—C6	1.395 (5)	C14—C15	1.389 (6)
C2—C3	1.380 (7)	C15—C16	1.383 (7)

supplementary materials

C2—C13	1.571 (6)	C15—C26	1.529 (7)
C3—C4	1.390 (7)	C16—C17	1.378 (8)
C3—H3	0.9300	C16—H16	0.9300
C4—C5	1.340 (7)	C17—C18	1.360 (8)
C5—C6	1.367 (6)	C18—C19	1.366 (7)
C5—H5	0.9300	C18—H18	0.9300
C6—H6	0.9300	C19—H19	0.9300
C7—C8	1.372 (5)	C20—C25	1.367 (5)
C7—C12	1.380 (5)	C20—C21	1.381 (5)
C8—C9	1.388 (6)	C21—C22	1.389 (5)
C8—H8	0.9300	C21—H21	0.9300
C9—C10	1.372 (7)	C22—C23	1.379 (6)
C9—H9	0.9300	C22—H22	0.9300
C10—C11	1.353 (7)	C23—C24	1.361 (6)
C10—H10	0.9300	C23—H23	0.9300
C11—C12	1.377 (6)	C24—C25	1.378 (5)
C11—H11	0.9300	C24—H24	0.9300
C12—H12	0.9300	C25—H25	0.9300
C13—H13A	0.9600	C26—H26A	0.9600
C13—H13B	0.9600	C26—H26B	0.9600
C13—H13C	0.9600	C26—H26C	0.9600
O1—S1—O2	118.88 (18)	O4—S2—O3	118.30 (18)
O1—S1—N1	110.6 (2)	O4—S2—N2	110.14 (19)
O2—S1—N1	103.80 (17)	O3—S2—N2	104.50 (17)
O1—S1—C1	106.88 (19)	O4—S2—C14	107.02 (18)
O2—S1—C1	109.17 (19)	O3—S2—C14	109.55 (19)
N1—S1—C1	107.02 (18)	N2—S2—C14	106.81 (18)
C7—N1—S1	126.6 (3)	C20—N2—S2	128.7 (3)
C7—N1—H1N	113 (3)	C20—N2—H2N	116 (3)
S1—N1—H1N	119 (3)	S2—N2—H2N	115 (3)
C2—C1—C6	120.1 (4)	C19—C14—C15	120.7 (4)
C2—C1—S1	122.2 (3)	C19—C14—S2	117.1 (3)
C6—C1—S1	117.7 (3)	C15—C14—S2	122.3 (3)
C3—C2—C1	117.7 (4)	C16—C15—C14	117.3 (5)
C3—C2—C13	118.5 (4)	C16—C15—C26	119.1 (5)
C1—C2—C13	123.8 (4)	C14—C15—C26	123.6 (5)
C2—C3—C4	120.8 (5)	C17—C16—C15	121.0 (6)
C2—C3—H3	119.6	C17—C16—H16	119.5
C4—C3—H3	119.6	C15—C16—H16	119.5
C5—C4—C3	121.4 (5)	C18—C17—C16	121.4 (6)
C5—C4—C11	120.5 (4)	C18—C17—Cl2	119.5 (5)
C3—C4—C11	118.0 (4)	C16—C17—Cl2	119.1 (6)
C4—C5—C6	118.7 (4)	C17—C18—C19	118.4 (5)
C4—C5—H5	120.6	C17—C18—H18	120.8
C6—C5—H5	120.6	C19—C18—H18	120.8
C5—C6—C1	121.2 (4)	C18—C19—C14	121.1 (5)
C5—C6—H6	119.4	C18—C19—H19	119.4
C1—C6—H6	119.4	C14—C19—H19	119.4
C8—C7—C12	119.8 (4)	C25—C20—C21	120.2 (3)

C8—C7—N1	116.9 (4)	C25—C20—N2	116.9 (3)
C12—C7—N1	123.3 (4)	C21—C20—N2	123.0 (3)
C7—C8—C9	119.3 (4)	C20—C21—C22	119.3 (4)
C7—C8—H8	120.4	C20—C21—H21	120.4
C9—C8—H8	120.4	C22—C21—H21	120.4
C10—C9—C8	120.7 (5)	C23—C22—C21	120.1 (4)
C10—C9—H9	119.7	C23—C22—H22	120.0
C8—C9—H9	119.7	C21—C22—H22	120.0
C11—C10—C9	119.5 (4)	C24—C23—C22	119.9 (4)
C11—C10—H10	120.2	C24—C23—H23	120.1
C9—C10—H10	120.2	C22—C23—H23	120.1
C10—C11—C12	120.9 (4)	C23—C24—C25	120.5 (4)
C10—C11—H11	119.6	C23—C24—H24	119.7
C12—C11—H11	119.6	C25—C24—H24	119.7
C11—C12—C7	119.9 (4)	C20—C25—C24	120.1 (4)
C11—C12—H12	120.1	C20—C25—H25	119.9
C7—C12—H12	120.1	C24—C25—H25	119.9
C2—C13—H13A	109.5	C15—C26—H26A	109.5
C2—C13—H13B	109.5	C15—C26—H26B	109.5
H13A—C13—H13B	109.5	H26A—C26—H26B	109.5
C2—C13—H13C	109.5	C15—C26—H26C	109.5
H13A—C13—H13C	109.5	H26A—C26—H26C	109.5
H13B—C13—H13C	109.5	H26B—C26—H26C	109.5
O1—S1—N1—C7	54.0 (4)	O4—S2—N2—C20	−46.3 (4)
O2—S1—N1—C7	−177.4 (3)	O3—S2—N2—C20	−174.3 (3)
C1—S1—N1—C7	−62.0 (4)	C14—S2—N2—C20	69.6 (4)
O1—S1—C1—C2	174.7 (4)	O4—S2—C14—C19	6.9 (4)
O2—S1—C1—C2	44.9 (5)	O3—S2—C14—C19	136.3 (3)
N1—S1—C1—C2	−66.8 (4)	N2—S2—C14—C19	−111.1 (3)
O1—S1—C1—C6	−4.2 (4)	O4—S2—C14—C15	−173.1 (4)
O2—S1—C1—C6	−134.0 (3)	O3—S2—C14—C15	−43.7 (4)
N1—S1—C1—C6	114.2 (3)	N2—S2—C14—C15	68.9 (4)
C6—C1—C2—C3	−1.2 (8)	C19—C14—C15—C16	−0.3 (7)
S1—C1—C2—C3	179.9 (5)	S2—C14—C15—C16	179.7 (4)
C6—C1—C2—C13	−179.3 (5)	C19—C14—C15—C26	179.9 (5)
S1—C1—C2—C13	1.7 (8)	S2—C14—C15—C26	−0.1 (7)
C1—C2—C3—C4	2.1 (10)	C14—C15—C16—C17	0.5 (9)
C13—C2—C3—C4	−179.7 (6)	C26—C15—C16—C17	−179.6 (6)
C2—C3—C4—C5	−2.0 (11)	C15—C16—C17—C18	−1.3 (11)
C2—C3—C4—Cl1	177.8 (6)	C15—C16—C17—Cl2	−178.9 (5)
C3—C4—C5—C6	1.0 (9)	C16—C17—C18—C19	1.8 (10)
Cl1—C4—C5—C6	−178.9 (4)	Cl2—C17—C18—C19	179.4 (4)
C4—C5—C6—C1	−0.1 (7)	C17—C18—C19—C14	−1.5 (8)
C2—C1—C6—C5	0.3 (7)	C15—C14—C19—C18	0.8 (7)
S1—C1—C6—C5	179.2 (3)	S2—C14—C19—C18	−179.2 (4)
S1—N1—C7—C8	163.5 (3)	S2—N2—C20—C25	−176.1 (3)
S1—N1—C7—C12	−17.8 (6)	S2—N2—C20—C21	5.9 (6)
C12—C7—C8—C9	0.5 (6)	C25—C20—C21—C22	−0.5 (6)
N1—C7—C8—C9	179.2 (4)	N2—C20—C21—C22	177.4 (4)

supplementary materials


C7—C8—C9—C10	0.1 (7)	C20—C21—C22—C23	0.7 (6)
C8—C9—C10—C11	-0.2 (8)	C21—C22—C23—C24	-0.2 (7)
C9—C10—C11—C12	-0.3 (7)	C22—C23—C24—C25	-0.5 (7)
C10—C11—C12—C7	0.9 (7)	C21—C20—C25—C24	-0.2 (6)
C8—C7—C12—C11	-1.0 (6)	N2—C20—C25—C24	-178.2 (4)
N1—C7—C12—C11	-179.7 (4)	C23—C24—C25—C20	0.7 (7)

Hydrogen-bond geometry (\AA , $^\circ$)

$D\text{—H}\cdots A$	$D\text{—H}$	$H\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
N1—H1N \cdots O2 ⁱ	0.91 (5)	2.02 (5)	2.922 (4)	175 (4)
N2—H2N \cdots O3 ⁱⁱ	0.88 (4)	2.03 (5)	2.906 (4)	173 (4)

Symmetry codes: (i) $-x+1, -y+2, -z$; (ii) $-x, -y+1, -z+1$.

Fig. 1



supplementary materials

Fig. 2

