

Dimethyl (1-hydroxy-1,2-diphenylethyl)-phosphonate

Nurcan Acar,^a M. Nawaz Tahir,^{b*} Hamza Yilmaz,^a
Muhammad Saeed Ahmad Chishti^b and Muhammad Ali
Malik^b

^aDepartment of Chemistry, Faculty of Science, University of Ankara, Ankara, Turkey,

and ^bDepartment of Physics, University of Sargodha, Sargodha, Pakistan

Correspondence e-mail: dmntahir_uos@yahoo.com


Received 30 January 2009; accepted 3 February 2009

Key indicators: single-crystal X-ray study; $T = 296\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.005\text{ \AA}$;
 R factor = 0.064; wR factor = 0.159; data-to-parameter ratio = 14.6.

In the molecule of the title compound, $\text{C}_{16}\text{H}_{19}\text{O}_4\text{P}$, the coordination around the P atom is distorted tetrahedral. The aromatic rings are oriented at a dihedral angle of $72.28(11)^\circ$. Intramolecular C—H···O hydrogen bonding result in the formation of five- and six-membered rings. In the crystal structure, intermolecular C—H···O hydrogen bonds link the molecules. There is also a weak C—H···π interaction.

Related literature

For related structures, see: Hudson *et al.* (1993); Tahir *et al.* (2007); Wroblewski *et al.* (2000).



Experimental

Crystal data

$\text{C}_{16}\text{H}_{19}\text{O}_4\text{P}$

$M_r = 306.28$

Monoclinic, $P2_1/c$

$a = 6.0671(12)\text{ \AA}$

$b = 17.0962(11)\text{ \AA}$

$c = 15.0502(12)\text{ \AA}$

$\beta = 95.0021(11)^\circ$

$V = 1555.1(3)\text{ \AA}^3$

$Z = 4$

Mo $K\alpha$ radiation

$\mu = 0.19\text{ mm}^{-1}$

$T = 296\text{ K}$

$0.28 \times 0.12 \times 0.10\text{ mm}$

Data collection

Enraf–Nonius CAD-4 diffractometer

Absorption correction: ψ scan (North *et al.*, 1968)
 $T_{\min} = 0.968$, $T_{\max} = 0.982$
2800 measured reflections

2794 independent reflections
1655 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.059$
3 standard reflections
frequency: 120 min
intensity decay: -1.3%

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.064$
 $wR(F^2) = 0.159$
 $S = 1.01$
2794 reflections

191 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.40\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.35\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H} \cdots A$	$D-\text{H}$	$\text{H} \cdots A$	$D \cdots A$	$D-\text{H} \cdots A$
O1—H1···O2 ⁱ	0.8200	1.9000	2.710 (3)	172.00
C2—H2···O4	0.9300	2.5200	2.956 (4)	109.00
C6—H6···O1	0.9300	2.3000	2.671 (4)	103.00
C14—H14···O1 ⁱⁱ	0.9300	2.5700	3.432 (5)	154.00
C16—H16A···O3 ⁱⁱ	0.9600	2.4700	3.336 (5)	149.00
C15—H15B···CgA	0.9600	2.7200	3.608 (5)	154.00

Symmetry codes: (i) $-x + 2, -y, -z$; (ii) $x - 1, y, z$. CgA is centroid of the C1—C6 ring.

Data collection: *CAD-4 EXPRESS* (Enraf–Nonius, 1994); cell refinement: *CAD-4 EXPRESS*; data reduction: *XCAD4* (Harms & Wocadlo, 1995); program(s) used to solve structure: *SHELXS86* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997) and *PLATON* (Spek, 2003); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HK2619).

References

- Enraf–Nonius (1994). *CAD-4 EXPRESS*. Enraf–Nonius, Delft, The Netherlands.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Harms, K. & Wocadlo, S. (1995). *XCAD4*. University of Marburg, Germany.
- Hudson, H. R., McPartlin, M., Matthews, R. W., Powell, H. R., Yusuf, R. O., Jaszy, Z. M., Keglevich, G., Petnehazy, I. & Toke, L. (1993). *Phosphorus Sulfur Silicon Relat. Elem.* **79**, 239–243.
- North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst.* **A24**, 351–359.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.
- Tahir, M. N., Acar, N., Yilmaz, H., Danish, M. & Ülkü, D. (2007). *Acta Cryst.* **E63**, o3817–o3818.
- Wroblewski, A. E., Maniukiewicz, W. & Karolczak, W. (2000). *J. Chem. Soc. Perkin Trans. 1*, pp. 1433–1435.

supplementary materials

Acta Cryst. (2009). E65, o481 [doi:10.1107/S1600536809004036]

Dimethyl (1-hydroxy-1,2-diphenylethyl)phosphonate

N. Acar, M. N. Tahir, H. Yilmaz, M. S. A. Chishti and M. A. Malik

Comment

The crystal structure of (*R*)-dimethyl [(2-chlorophenyl)hydroxymethyl]- phosphonate (Tahir *et al.*, 2007), which is a member of α -hydroxy phosphonates, has been reported, previously. In continuation to the study of such organic compounds, we report herein the synthesis and crystal structure of the title compound, (I).

In the molecule of (I), (Fig. 1) the coordination around the P atom is a distorted tetrahedral. The crystal structure of dimethyl α -chloromethyl- α -hydroxybenzylphosphonate, (II) (Hudson *et al.*, 1993) and C₂₄H₂₆NO₄P, (III) (Wroblewski *et al.*, 2000) have also been reported, in which both of them have similar coordinations around the C-atom having α -hydroxy group. In (I), the benzene rings A (C1-C6) and B (C9-C14) are oriented at a dihedral angle of 72.28 (11) $^{\circ}$. The intramolecular C-H \cdots O hydrogen bonds (Table 1) result in the formations of five- and six-membered rings C (O1/C1/C6/C7/H6) and D (P1/O4/C1/C2/C7/H2), respectively, having planar and boat conformations.

In the crystal structure, intermolecular O-H \cdots O and C-H \cdots O hydrogen bonds (Table 1) link the molecules (Fig. 2), in which they may be effective in the stabilization of the structure. There is also a weak C—H \cdots π interaction (Table 1).

Experimental

For the preparation of the title compound, 2-phenylacetophenone (1.96 g, 10 mmol) was dissolved in dimethylphosphonate (1.10 g, 10 mmol) at room temperature. Then, KF (2.5 g) and ψ -Al₂O₃ (2.5 g) were added partwise and the mixture was kept at room temperature for 72 h. The product was extracted twice with 50 ml portions of a dichloromethane/methanol mixture (1:1). After the evaporation of the solvent on a rotary evaporator, the residue was crystallized from a mixture of diethyl ether/acetone (3:1) (m.p. 414 K).

Refinement

H atoms were positioned geometrically, with O-H = 0.82 Å (for OH), C-H = 0.93, 0.96 and 0.97 Å for aromatic, methyl and methylene H, and constrained to ride on their parent atoms with U_{iso}(H) = xU_{eq}(C), where x = 1.5 for methyl H, and x = 1.2 for all other H atoms.

Figures




Fig. 1. The molecular structure of the title molecule, with the atom-numbering scheme. Displacement ellipsoids are drawn at the 50% probability level.

supplementary materials




Fig. 2. A partial packing diagram of the title compound. Hydrogen bonds are shown as dashed lines.

Dimethyl (1-hydroxy-1,2-diphenylethyl)phosphonate

Crystal data

C ₁₆ H ₁₉ O ₄ P	$F_{000} = 648$
$M_r = 306.28$	$D_x = 1.308 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Melting point = 83–86 K
Hall symbol: -P 2ybc	Mo $K\alpha$ radiation
$a = 6.0671 (12) \text{ \AA}$	$\lambda = 0.71073 \text{ \AA}$
$b = 17.0962 (11) \text{ \AA}$	Cell parameters from 15 reflections
$c = 15.0502 (12) \text{ \AA}$	$\theta = 10.0\text{--}11.3^\circ$
$\beta = 95.0021 (11)^\circ$	$\mu = 0.19 \text{ mm}^{-1}$
$V = 1555.1 (3) \text{ \AA}^3$	$T = 296 \text{ K}$
$Z = 4$	Needle, colorless
	$0.28 \times 0.12 \times 0.10 \text{ mm}$

Data collection

Enraf–Nonius CAD-4 diffractometer	$\theta_{\max} = 25.2^\circ$
$\omega/2\theta$ scans	$\theta_{\min} = 2.4^\circ$
Absorption correction: ψ scan (North <i>et al.</i> , 1968)	$h = 0\rightarrow 7$
$T_{\min} = 0.968$, $T_{\max} = 0.982$	$k = 0\rightarrow 20$
2800 measured reflections	$l = -18\rightarrow 17$
2794 independent reflections	3 standard reflections
1655 reflections with $I > 2\sigma(I)$	every 120 min
$R_{\text{int}} = 0.059$	intensity decay: –1.3%

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	H-atom parameters constrained
$R[F^2 > 2\sigma(F^2)] = 0.064$	$w = 1/[\sigma^2(F_o^2) + (0.088P)^2]$
$wR(F^2) = 0.159$	where $P = (F_o^2 + 2F_c^2)/3$
	$(\Delta/\sigma)_{\max} < 0.001$

$S = 1.01$ $\Delta\rho_{\max} = 0.40 \text{ e \AA}^{-3}$
 2794 reflections $\Delta\rho_{\min} = -0.35 \text{ e \AA}^{-3}$
 191 parameters Extinction correction: none
 Primary atom site location: structure-invariant direct
 methods

Special details

Geometry. Bond distances, angles *etc.* have been calculated using the rounded fractional coordinates. All su's are estimated from the variances of the (full) variance-covariance matrix. The cell e.s.d.'s are taken into account in the estimation of distances, angles and torsion angles

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
P1	0.79483 (14)	0.12098 (5)	-0.05193 (6)	0.0416 (3)
O1	1.0479 (3)	0.10996 (12)	0.09544 (16)	0.0460 (8)
O2	0.8463 (4)	0.04357 (13)	-0.08487 (15)	0.0576 (9)
O3	0.9546 (5)	0.18125 (15)	-0.08944 (18)	0.0659 (10)
O4	0.5517 (4)	0.14911 (15)	-0.07722 (15)	0.0582 (9)
C1	0.7793 (5)	0.21108 (17)	0.1007 (2)	0.0368 (10)
C2	0.5716 (5)	0.24481 (19)	0.0869 (2)	0.0444 (11)
C3	0.5409 (6)	0.3227 (2)	0.1076 (2)	0.0533 (12)
C4	0.7133 (7)	0.3671 (2)	0.1449 (3)	0.0567 (14)
C5	0.9175 (7)	0.3333 (2)	0.1623 (3)	0.0555 (14)
C6	0.9516 (6)	0.25648 (18)	0.1401 (2)	0.0470 (11)
C7	0.8226 (5)	0.12778 (17)	0.0709 (2)	0.0387 (10)
C8	0.6654 (5)	0.06725 (18)	0.1090 (2)	0.0435 (11)
C9	0.6703 (6)	0.06607 (17)	0.2091 (2)	0.0411 (10)
C10	0.8471 (6)	0.03455 (19)	0.2618 (2)	0.0522 (12)
C11	0.8457 (7)	0.0313 (2)	0.3538 (2)	0.0573 (14)
C12	0.6657 (8)	0.0583 (2)	0.3943 (3)	0.0628 (14)
C13	0.4908 (8)	0.0895 (2)	0.3432 (3)	0.0635 (16)
C14	0.4917 (6)	0.0931 (2)	0.2520 (3)	0.0537 (12)
C15	0.9185 (9)	0.2641 (2)	-0.0979 (3)	0.0863 (19)
C16	0.4445 (7)	0.1432 (3)	-0.1647 (3)	0.0733 (16)
H1	1.06744	0.06272	0.09092	0.0552*
H2	0.45212	0.21495	0.06361	0.0534*
H3	0.40178	0.34518	0.09602	0.0636*
H4	0.69192	0.41953	0.15824	0.0679*
H5	1.03397	0.36254	0.18933	0.0666*
H6	1.09151	0.23458	0.15143	0.0565*
H8A	0.51529	0.07819	0.08461	0.0520*
H8B	0.70402	0.01557	0.08879	0.0520*
H10	0.96854	0.01529	0.23501	0.0623*
H11	0.96685	0.01083	0.38820	0.0688*
H12	0.66330	0.05520	0.45591	0.0750*
H13	0.36975	0.10860	0.37031	0.0764*

supplementary materials

H14	0.37005	0.11401	0.21825	0.0644*
H15A	1.04258	0.28789	-0.12280	0.1294*
H15B	0.90211	0.28613	-0.04020	0.1294*
H15C	0.78662	0.27362	-0.13644	0.1294*
H16A	0.29675	0.16336	-0.16523	0.1095*
H16B	0.43893	0.08928	-0.18277	0.1095*
H16C	0.52532	0.17277	-0.20525	0.1095*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
P1	0.0398 (5)	0.0398 (5)	0.0458 (5)	0.0101 (4)	0.0066 (4)	-0.0001 (4)
O1	0.0323 (13)	0.0371 (12)	0.0669 (15)	0.0059 (10)	-0.0053 (11)	-0.0040 (11)
O2	0.0693 (17)	0.0495 (15)	0.0533 (14)	0.0195 (13)	0.0007 (12)	-0.0048 (11)
O3	0.0681 (18)	0.0571 (17)	0.0766 (17)	0.0082 (13)	0.0295 (14)	0.0097 (13)
O4	0.0503 (15)	0.0781 (17)	0.0450 (14)	0.0233 (13)	-0.0034 (11)	-0.0082 (12)
C1	0.0363 (18)	0.0357 (17)	0.0389 (17)	0.0016 (14)	0.0064 (14)	0.0009 (13)
C2	0.0339 (19)	0.047 (2)	0.053 (2)	0.0006 (15)	0.0075 (14)	-0.0066 (15)
C3	0.051 (2)	0.051 (2)	0.060 (2)	0.0171 (19)	0.0163 (18)	-0.0002 (17)
C4	0.071 (3)	0.0351 (19)	0.067 (2)	0.0015 (19)	0.024 (2)	-0.0090 (17)
C5	0.057 (3)	0.041 (2)	0.069 (2)	-0.0050 (18)	0.0085 (19)	-0.0118 (17)
C6	0.0391 (19)	0.042 (2)	0.059 (2)	-0.0002 (15)	0.0000 (16)	0.0002 (16)
C7	0.0306 (18)	0.0371 (17)	0.0476 (18)	0.0025 (14)	-0.0013 (14)	-0.0015 (14)
C8	0.0370 (19)	0.0395 (18)	0.053 (2)	-0.0038 (14)	-0.0009 (15)	0.0002 (15)
C9	0.0426 (19)	0.0294 (16)	0.0501 (19)	-0.0046 (14)	-0.0022 (16)	-0.0009 (14)
C10	0.054 (2)	0.048 (2)	0.054 (2)	0.0105 (17)	0.0010 (17)	0.0012 (16)
C11	0.063 (3)	0.053 (2)	0.054 (2)	0.0051 (19)	-0.0051 (19)	0.0080 (18)
C12	0.085 (3)	0.059 (2)	0.045 (2)	-0.010 (2)	0.009 (2)	-0.0025 (18)
C13	0.062 (3)	0.068 (2)	0.063 (3)	0.000 (2)	0.019 (2)	-0.004 (2)
C14	0.041 (2)	0.055 (2)	0.065 (2)	-0.0028 (16)	0.0034 (18)	0.0010 (17)
C15	0.131 (4)	0.050 (3)	0.082 (3)	-0.005 (3)	0.032 (3)	0.010 (2)
C16	0.061 (3)	0.098 (3)	0.059 (2)	0.012 (2)	-0.006 (2)	-0.001 (2)

Geometric parameters (\AA , $^\circ$)

P1—O2	1.457 (2)	C12—C13	1.364 (6)
P1—O3	1.554 (3)	C13—C14	1.374 (6)
P1—O4	1.567 (3)	C2—H2	0.9300
P1—C7	1.845 (3)	C3—H3	0.9300
O1—C7	1.418 (4)	C4—H4	0.9300
O3—C15	1.437 (4)	C5—H5	0.9300
O4—C16	1.420 (5)	C6—H6	0.9300
O1—H1	0.8200	C8—H8A	0.9700
C1—C6	1.392 (5)	C8—H8B	0.9700
C1—C7	1.523 (4)	C10—H10	0.9300
C1—C2	1.385 (4)	C11—H11	0.9300
C2—C3	1.384 (5)	C12—H12	0.9300
C3—C4	1.372 (5)	C13—H13	0.9300
C4—C5	1.371 (6)	C14—H14	0.9300

C5—C6	1.375 (5)	C15—H15A	0.9600
C7—C8	1.551 (4)	C15—H15B	0.9600
C8—C9	1.504 (4)	C15—H15C	0.9600
C9—C14	1.388 (5)	C16—H16A	0.9600
C9—C10	1.386 (5)	C16—H16B	0.9600
C10—C11	1.387 (4)	C16—H16C	0.9600
C11—C12	1.376 (6)		
O1···O2	3.094 (3)	C9···H16B ^{vi}	2.7600
O1···O3	3.045 (4)	C10···H16B ^{vi}	2.9200
O1···C10	3.154 (4)	C10···H1	3.0400
O1···O2 ⁱ	2.710 (3)	C11···H5 ^{vii}	3.0600
O2···O1 ⁱ	2.710 (3)	C12···H15C ⁱⁱⁱ	3.0100
O2···O1	3.094 (3)	C13···H15C ⁱⁱⁱ	2.9500
O3···C16 ⁱⁱ	3.336 (5)	C16···H15C	3.0500
O3···O1	3.045 (4)	H1···O2	2.8800
O4···C2	2.956 (4)	H1···C10	3.0400
O1···H14 ⁱⁱ	2.5700	H1···H8B	2.3500
O1···H6	2.3000	H1···H10	2.4400
O1···H10	2.7300	H1···O2 ⁱ	1.9000
O1···H8A ⁱⁱ	2.9100	H2···O4	2.5200
O2···H16B	2.8700	H2···C8	2.8900
O2···H8B ⁱ	2.9100	H2···H8A	2.3900
O2···H10 ⁱ	2.8000	H5···C11 ^{viii}	3.0600
O2···H1 ⁱ	1.9000	H6···O1	2.3000
O2···H8B	2.8600	H8A···O1 ^v	2.9100
O2···H1	2.8800	H8A···O4	2.7500
O3···H16A ⁱⁱ	2.4700	H8A···C2	2.8700
O4···H2	2.5200	H8A···H2	2.3900
O4···H8A	2.7500	H8A···H14	2.3500
O4···H15C	2.7500	H8B···O2	2.8600
C1···C15	3.303 (5)	H8B···H1	2.3500
C2···O4	2.956 (4)	H8B···O2 ⁱ	2.9100
C2···C9	3.590 (4)	H8B···H16B ^{vi}	2.4900
C3···C16 ⁱⁱⁱ	3.574 (5)	H10···O1	2.7300
C4···C16 ⁱⁱⁱ	3.423 (6)	H10···H1	2.4400
C6···C15	3.572 (5)	H10···O2 ⁱ	2.8000
C9···C2	3.590 (4)	H14···O1 ^v	2.5700
C10···O1	3.154 (4)	H14···H8A	2.3500
C11···C15 ⁱⁱⁱ	3.592 (5)	H15B···C1	2.6400
C12···C15 ⁱⁱⁱ	3.399 (5)	H15B···C2	2.9700
C15···C1	3.303 (5)	H15B···C6	2.7500
C15···C11 ^{iv}	3.592 (5)	H15C···O4	2.7500
C15···C12 ^{iv}	3.399 (5)	H15C···C16	3.0500
C15···C6	3.572 (5)	H15C···H16C	2.5000

supplementary materials

C16···O3 ^v	3.336 (5)	H15C···C12 ^{iv}	3.0100
C16···C3 ^{iv}	3.574 (5)	H15C···C13 ^{iv}	2.9500
C16···C4 ^{iv}	3.423 (6)	H16A···O3 ^v	2.4700
C1···H15B	2.6400	H16B···O2	2.8700
C2···H8A	2.8700	H16B···C8 ^{vi}	2.9900
C2···H15B	2.9700	H16B···C9 ^{vi}	2.7600
C3···H16C ⁱⁱⁱ	2.8300	H16B···C10 ^{vi}	2.9200
C4···H16C ⁱⁱⁱ	2.7000	H16B···H8B ^{vi}	2.4900
C6···H15B	2.7500	H16C···H15C	2.5000
C8···H16B ^{vi}	2.9900	H16C···C3 ^{iv}	2.8300
C8···H2	2.8900	H16C···C4 ^{iv}	2.7000
O2—P1—O3	108.65 (15)	C4—C3—H3	120.00
O2—P1—O4	114.93 (14)	C3—C4—H4	120.00
O2—P1—C7	113.24 (14)	C5—C4—H4	120.00
O3—P1—O4	108.19 (15)	C4—C5—H5	120.00
O3—P1—C7	108.47 (14)	C6—C5—H5	120.00
O4—P1—C7	103.05 (13)	C1—C6—H6	120.00
P1—O3—C15	126.2 (3)	C5—C6—H6	120.00
P1—O4—C16	123.3 (2)	C7—C8—H8A	109.00
C7—O1—H1	109.00	C7—C8—H8B	109.00
C2—C1—C6	118.1 (3)	C9—C8—H8A	109.00
C6—C1—C7	120.3 (3)	C9—C8—H8B	109.00
C2—C1—C7	121.6 (3)	H8A—C8—H8B	108.00
C1—C2—C3	120.4 (3)	C9—C10—H10	119.00
C2—C3—C4	120.8 (3)	C11—C10—H10	120.00
C3—C4—C5	119.3 (3)	C10—C11—H11	120.00
C4—C5—C6	120.6 (4)	C12—C11—H11	120.00
C1—C6—C5	120.8 (3)	C11—C12—H12	120.00
P1—C7—C1	110.6 (2)	C13—C12—H12	120.00
P1—C7—C8	108.9 (2)	C12—C13—H13	120.00
O1—C7—C1	108.2 (2)	C14—C13—H13	120.00
O1—C7—C8	111.7 (2)	C9—C14—H14	119.00
C1—C7—C8	112.8 (2)	C13—C14—H14	119.00
P1—C7—O1	104.4 (2)	O3—C15—H15A	109.00
C7—C8—C9	114.9 (3)	O3—C15—H15B	109.00
C8—C9—C10	121.7 (3)	O3—C15—H15C	109.00
C10—C9—C14	117.4 (3)	H15A—C15—H15B	109.00
C8—C9—C14	120.9 (3)	H15A—C15—H15C	109.00
C9—C10—C11	121.0 (3)	H15B—C15—H15C	109.00
C10—C11—C12	120.2 (4)	O4—C16—H16A	110.00
C11—C12—C13	119.3 (4)	O4—C16—H16B	109.00
C12—C13—C14	120.7 (4)	O4—C16—H16C	110.00
C9—C14—C13	121.4 (4)	H16A—C16—H16B	109.00
C1—C2—H2	120.00	H16A—C16—H16C	110.00
C3—C2—H2	120.00	H16B—C16—H16C	109.00
C2—C3—H3	120.00		
O2—P1—O3—C15	158.6 (3)	C2—C1—C7—C8	-55.0 (4)

O4—P1—O3—C15	33.2 (3)	C6—C1—C7—P1	-110.6 (3)
C7—P1—O3—C15	-77.9 (3)	C6—C1—C7—O1	3.2 (4)
O2—P1—O4—C16	-47.0 (3)	C6—C1—C7—C8	127.3 (3)
O3—P1—O4—C16	74.6 (3)	C1—C2—C3—C4	-2.3 (5)
C7—P1—O4—C16	-170.7 (3)	C2—C3—C4—C5	-0.5 (6)
O2—P1—C7—O1	60.7 (2)	C3—C4—C5—C6	2.1 (6)
O2—P1—C7—C1	176.8 (2)	C4—C5—C6—C1	-0.9 (6)
O2—P1—C7—C8	-58.8 (2)	P1—C7—C8—C9	180.0 (2)
O3—P1—C7—O1	-60.0 (2)	O1—C7—C8—C9	65.2 (3)
O3—P1—C7—C1	56.1 (2)	C1—C7—C8—C9	-56.9 (3)
O3—P1—C7—C8	-179.5 (2)	C7—C8—C9—C10	-73.5 (4)
O4—P1—C7—O1	-174.51 (18)	C7—C8—C9—C14	109.9 (3)
O4—P1—C7—C1	-58.4 (2)	C8—C9—C10—C11	-177.6 (3)
O4—P1—C7—C8	66.0 (2)	C14—C9—C10—C11	-0.8 (5)
C6—C1—C2—C3	3.4 (4)	C8—C9—C14—C13	177.4 (3)
C7—C1—C2—C3	-174.4 (3)	C10—C9—C14—C13	0.6 (5)
C2—C1—C6—C5	-1.8 (5)	C9—C10—C11—C12	1.1 (5)
C7—C1—C6—C5	176.0 (3)	C10—C11—C12—C13	-1.2 (5)
C2—C1—C7—P1	67.1 (3)	C11—C12—C13—C14	1.0 (6)
C2—C1—C7—O1	-179.1 (3)	C12—C13—C14—C9	-0.7 (6)
Symmetry codes: (i) $-x+2, -y, -z$; (ii) $x+1, y, z$; (iii) $x, -y+1/2, z+1/2$; (iv) $x, -y+1/2, z-1/2$; (v) $x-1, y, z$; (vi) $-x+1, -y, -z$; (vii) $-x+2, y-1/2, -z+1/2$; (viii) $-x+2, y+1/2, -z+1/2$.			

Hydrogen-bond geometry (\AA , $^\circ$)

$D\text{—H}\cdots A$	$D\text{—H}$	$H\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
O1—H1 ⁱ …O2 ⁱ	0.8200	1.9000	2.710 (3)	172.00
C2—H2…O4	0.9300	2.5200	2.956 (4)	109.00
C6—H6…O1	0.9300	2.3000	2.671 (4)	103.00
C14—H14…O1 ^v	0.9300	2.5700	3.432 (5)	154.00
C16—H16A…O3 ^v	0.9600	2.4700	3.336 (5)	149.00
C15—H15B…CgA	0.9600	2.7200	3.608 (5)	154.00

Symmetry codes: (i) $-x+2, -y, -z$; (v) $x-1, y, z$.

supplementary materials

Fig. 1




Fig. 2

