

1,2-Bis(di-2-pyridylphosphino)ethane

Susan J. Berners-Price, Maribel Navarro‡ and Brian W. Skelton*

Chemistry, School of Biomedical, Biomolecular and Chemical Sciences, The University of Western Australia, 35 Stirling Highway, Crawley, Perth 6009, Western Australia, Australia

Correspondence e-mail: brian.skelton@uwa.edu.au


Received 20 January 2009; accepted 9 February 2009

Key indicators: single-crystal X-ray study; $T = 110$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.052; wR factor = 0.112; data-to-parameter ratio = 17.9.

The crystal structure of the title compound, $\text{C}_{22}\text{H}_{20}\text{N}_4\text{O}_2\text{P}_2$, consists of two independent half-molecules, both of which lie on crystallographic inversion centres. There are no significant differences between the two molecules.

Related literature

For the antitumour properties of metal complexes of bidentate tertiary phosphine ligands with pyridyl substituents, see: McKeage *et al.* (2000); Barnard & Berners-Price (2007); Liu *et al.* (2008). The crystal structure of the parent 1,2-bis(di-2-pyridylphosphino)ethane molecule has been determined (Jones *et al.*, 1999). The structure of 1,2-bis(di-phenylphosphino)ethane dioxide (Calcagno *et al.*, 2000) is similar, with the two halves of the molecule related by a pseudo-inversion centre, but this is not isomorphous with the title compound.



Experimental

Crystal data

$\text{C}_{22}\text{H}_{20}\text{N}_4\text{O}_2\text{P}_2$	$\gamma = 112.559$ (7)°
$M_r = 434.36$	$V = 1055.67$ (16) Å ³
Triclinic, $P\bar{1}$	$Z = 2$
$a = 8.3760$ (6) Å	Mo $K\alpha$ radiation
$b = 8.8496$ (8) Å	$\mu = 0.23$ mm ⁻¹
$c = 16.2332$ (11) Å	$T = 110$ K
$\alpha = 105.627$ (7)°	$0.22 \times 0.10 \times 0.06$ mm
$\beta = 92.429$ (5)°	

Data collection

Oxford Diffraction Gemini diffractometer	10983 measured reflections
Absorption correction: Gaussian (<i>CrysAlis RED</i> ; Oxford Diffraction, 2008)	4842 independent reflections
$T_{\min} = 0.968$, $T_{\max} = 0.988$	2698 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.059$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.052$	271 parameters
$wR(F^2) = 0.112$	H-atom parameters constrained
$S = 0.86$	$\Delta\rho_{\text{max}} = 0.43$ e Å ⁻³
4842 reflections	$\Delta\rho_{\text{min}} = -0.34$ e Å ⁻³

Data collection: *CrysAlis CCD* (Oxford Diffraction, 2008); cell refinement: *CrysAlis RED* (Oxford Diffraction, 2008); data reduction: *CrysAlis RED*; program(s) used to solve structure: *SIR92* (Altomare *et al.*, 1994); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEPII* (Johnson, 1976); software used to prepare material for publication: *publCIF* (Westrip, 2009).

The authors thank the Australian Research Council for financial assistance.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: FJ2193).

References

- Altomare, A., Cascarano, G., Giacovazzo, C., Guagliardi, A., Burla, M. C., Polidori, G. & Camalli, M. (1994). *J. Appl. Cryst.* **27**, 435.
- Barnard, P. J. & Berners-Price, S. J. (2007). *Coord. Chem. Rev.* **251**, 1889–1902.
- Calcagno, P., Kariuki, B. M., Kitchin, S. J., Robinson, J. M. A., Philp, D. & Harris, K. D. M. (2000). *Chem. Eur. J.* **6**, 2338–2349.
- Johnson, C. K. (1976). *ORTEPII*. Report ORNL-5138. Oak Ridge National Laboratory, Tennessee, USA.
- Jones, N. D., MacFarlane, K. S., Smith, M. B., Schutte, R. P., Rettig, S. J. & James, B. R. (1999). *Inorg. Chem.* **38**, 3956–3966.
- Liu, J. J., Galettis, P., Farr, A., Maharaj, L., Samarasinha, H., McGechan, A. C., Baguley, B. C., Bowen, R. J., Berners-Price, S. J. & McKeage, M. J. (2008). *J. Inorg. Biochem.* **102**, 303–310.
- McKeage, M. J., Berners-Price, S. J., Galettis, P., Bowen, R. J., Brouwer, W., Ding, L., Zhuang, L. & Baguley, B. C. (2000). *Cancer Chemother. Pharmacol.* **46**, 343–350.
- Oxford Diffraction (2008). *CrysAlis CCD* and *CrysAlis RED*. Oxford Diffraction Ltd, Abingdon, Oxfordshire, England.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Westrip, S. P. (2009). *publCIF*. In preparation.

‡ Present Address: Laboratorio de Química Bioinorgánica, Centro de Química, Instituto Venezolano de Investigaciones Científicas (IVIC), Caracas 1020-A, Venezuela.

supplementary materials

Acta Cryst. (2009). E65, o542 [doi:10.1107/S1600536809004590]

1,2-Bis(di-2-pyridylphosphinoyl)ethane

S. J. Berners-Price, M. Navarro and B. W. Skelton

Comment

Bidentate tertiary phosphine ligands with pyridyl substituents, such as 1,2-bis(di-2-pyridylphosphino)ethane (d2pype) are of interest because a number of studies have shown that metal complexes with these ligands exhibit selective anti-tumour properties (McKeage *et al.*, 2000; Barnard and Berners-Price 2007; Liu *et al.*, 2008). During the course of our work in this area, we obtained crystals of the phosphine oxide d2pypeO₂ (**I**), which were suitable for X-ray diffraction studies.

Experimental

1,2-bis(di-2-pyridylphosphino)ethane (d2pype) was obtained from Strem Chemicals Inc. Single crystals of the title compound d2pypeO₂ (**I**) suitable for X-ray crystallographic analysis were obtained as a by-product of slow evaporation of a solution of d2pype and copper (I) iodide (molar ratio 2:1) in acetonitrile-tetrahydrofuran mixture.

Refinement

The assignments of the py ring N,C atoms were made on the basis of refinement and location of the H atoms. All H atoms were positioned geometrically and refined using a riding model with C—H = 0.95–0.99 Å and with $U_{\text{iso}}(\text{H}) = 1.2$ times $U_{\text{eq}}(\text{C})$.

Figures




Fig. 1. ORTEP drawing and atom labelling for molecule $n = 1$. Displacement ellipsoids of non-H atoms are drawn at the 50% probability level. The structure of the second, $n = 2$, molecule is very similar.

1,2-Bis(di-2-pyridylphosphinoyl)ethane

Crystal data

C₂₂H₂₀N₄O₂P₂

$M_r = 434.36$

Triclinic, $P\bar{1}$

Hall symbol: -p 1

$a = 8.3760$ (6) Å

$b = 8.8496$ (8) Å

$Z = 2$

$F_{000} = 452$

$D_x = 1.366$ Mg m⁻³

Mo $K\alpha$ radiation

$\lambda = 0.71073$ Å

Cell parameters from 2539 reflections

$\theta = 3.3$ – 32.6°

supplementary materials

$c = 16.2332 (11) \text{ \AA}$
 $\alpha = 105.627 (7)^\circ$
 $\beta = 92.429 (5)^\circ$
 $\gamma = 112.559 (7)^\circ$
 $V = 1055.67 (16) \text{ \AA}^3$

$\mu = 0.23 \text{ mm}^{-1}$
 $T = 110 \text{ K}$
Plate, colourless
 $0.22 \times 0.10 \times 0.06 \text{ mm}$

Data collection

Oxford Diffraction Gemini diffractometer
Radiation source: sealed tube
Monochromator: graphite
 $T = 110 \text{ K}$
 ω scans
Absorption correction: Gaussian (*CrysAlis RED*; Oxford Diffraction, 2008)
 $T_{\min} = 0.968$, $T_{\max} = 0.988$
10983 measured reflections

4842 independent reflections
2698 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.059$
 $\theta_{\max} = 27.5^\circ$
 $\theta_{\min} = 3.3^\circ$
 $h = -10 \rightarrow 10$
 $k = -11 \rightarrow 11$
 $l = -21 \rightarrow 21$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.052$
 $wR(F^2) = 0.112$
 $S = 0.86$
4842 reflections
271 parameters
Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map
Hydrogen site location: inferred from neighbouring sites
H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0495P)^2]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.002$
 $\Delta\rho_{\max} = 0.43 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.34 \text{ e \AA}^{-3}$
Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
P1	0.18196 (9)	0.21056 (9)	0.45935 (4)	0.02233 (18)
O1	0.3253 (2)	0.3074 (2)	0.53572 (11)	0.0272 (4)
C111	0.2511 (3)	0.0876 (3)	0.37091 (16)	0.0217 (6)
N112	0.1239 (3)	-0.0366 (3)	0.30732 (14)	0.0267 (5)

C113	0.1759 (4)	-0.1218 (4)	0.24034 (17)	0.0320 (7)
H113	0.0879	-0.2109	0.195	0.038*
C114	0.3487 (4)	-0.0891 (4)	0.23254 (18)	0.0320 (7)
H114	0.3782	-0.1528	0.1831	0.038*
C115	0.4767 (4)	0.0385 (4)	0.29857 (19)	0.0372 (8)
H115	0.5969	0.0641	0.2957	0.045*
C116	0.4284 (3)	0.1287 (4)	0.36902 (18)	0.0313 (7)
H116	0.5146	0.2172	0.4153	0.038*
C121	0.1197 (3)	0.3509 (3)	0.41523 (15)	0.0227 (6)
N122	-0.0518 (3)	0.2959 (3)	0.38395 (15)	0.0300 (6)
C123	-0.0959 (4)	0.4009 (4)	0.35135 (19)	0.0352 (7)
H123	-0.216	0.3661	0.3298	0.042*
C124	0.0236 (4)	0.5571 (4)	0.34727 (17)	0.0310 (7)
H124	-0.0142	0.6263	0.3228	0.037*
C125	0.1972 (4)	0.6111 (4)	0.37892 (17)	0.0316 (7)
H125	0.2816	0.7184	0.3773	0.038*
C126	0.2464 (3)	0.5055 (3)	0.41316 (16)	0.0256 (6)
H126	0.3659	0.5388	0.4351	0.031*
C10	-0.0180 (3)	0.0583 (3)	0.47773 (16)	0.0242 (6)
H10A	-0.0786	0.121	0.5142	0.029*
H10B	-0.0962	-0.0124	0.4216	0.029*
P2	0.68020 (8)	0.27338 (9)	0.04404 (4)	0.02132 (18)
O2	0.7095 (2)	0.3187 (2)	-0.03767 (11)	0.0278 (4)
C211	0.6499 (3)	0.4398 (3)	0.12649 (16)	0.0213 (6)
N212	0.6135 (3)	0.4071 (3)	0.20118 (14)	0.0279 (5)
C213	0.6005 (4)	0.5335 (4)	0.26420 (18)	0.0327 (7)
H213	0.5758	0.5138	0.3179	0.039*
C214	0.6210 (3)	0.6903 (4)	0.2556 (2)	0.0358 (7)
H214	0.6111	0.7759	0.3025	0.043*
C215	0.6559 (3)	0.7211 (4)	0.1781 (2)	0.0347 (7)
H215	0.6695	0.8277	0.1702	0.042*
C216	0.6707 (3)	0.5931 (4)	0.11223 (19)	0.0299 (7)
H216	0.6949	0.61	0.058	0.036*
C221	0.8672 (3)	0.2545 (3)	0.09411 (16)	0.0218 (6)
N222	0.8365 (3)	0.1354 (3)	0.13511 (14)	0.0285 (5)
C223	0.9776 (4)	0.1286 (4)	0.17259 (18)	0.0335 (7)
H223	0.9594	0.0457	0.2022	0.04*
C224	1.1484 (4)	0.2342 (4)	0.17114 (18)	0.0330 (7)
H224	1.2439	0.2227	0.1984	0.04*
C225	1.1771 (3)	0.3559 (4)	0.12949 (18)	0.0342 (7)
H225	1.2929	0.4316	0.1281	0.041*
C226	1.0340 (3)	0.3661 (4)	0.08959 (17)	0.0278 (6)
H226	1.0498	0.4481	0.0596	0.033*
C20	0.4938 (3)	0.0766 (3)	0.03363 (16)	0.0212 (6)
H20A	0.4874	0.0554	0.0905	0.025*
H20B	0.3853	0.0876	0.0159	0.025*

supplementary materials

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
P1	0.0206 (4)	0.0253 (4)	0.0224 (4)	0.0096 (3)	0.0051 (3)	0.0090 (3)
O1	0.0232 (9)	0.0318 (11)	0.0235 (10)	0.0099 (9)	0.0022 (8)	0.0063 (8)
C111	0.0238 (14)	0.0239 (15)	0.0205 (13)	0.0105 (12)	0.0034 (11)	0.0107 (12)
N112	0.0328 (13)	0.0261 (14)	0.0234 (12)	0.0148 (11)	0.0026 (10)	0.0073 (11)
C113	0.0433 (18)	0.0298 (17)	0.0235 (15)	0.0168 (15)	0.0002 (13)	0.0072 (13)
C114	0.0463 (18)	0.0262 (17)	0.0295 (16)	0.0186 (15)	0.0148 (14)	0.0115 (14)
C115	0.0328 (16)	0.0311 (18)	0.050 (2)	0.0137 (14)	0.0224 (15)	0.0120 (16)
C116	0.0261 (15)	0.0256 (17)	0.0358 (17)	0.0058 (13)	0.0070 (13)	0.0063 (14)
C121	0.0250 (14)	0.0237 (15)	0.0201 (14)	0.0105 (12)	0.0067 (11)	0.0066 (12)
N122	0.0216 (12)	0.0288 (14)	0.0420 (14)	0.0083 (11)	0.0007 (10)	0.0181 (12)
C123	0.0262 (15)	0.0337 (19)	0.0480 (19)	0.0103 (14)	0.0022 (13)	0.0197 (15)
C124	0.0354 (16)	0.0282 (17)	0.0341 (16)	0.0159 (14)	0.0042 (13)	0.0128 (14)
C125	0.0342 (16)	0.0201 (16)	0.0361 (17)	0.0037 (13)	0.0084 (13)	0.0126 (13)
C126	0.0239 (14)	0.0268 (16)	0.0232 (14)	0.0083 (12)	0.0031 (11)	0.0065 (12)
C10	0.0213 (14)	0.0279 (16)	0.0256 (14)	0.0110 (12)	0.0061 (11)	0.0104 (12)
P2	0.0193 (3)	0.0198 (4)	0.0227 (4)	0.0056 (3)	0.0010 (3)	0.0071 (3)
O2	0.0289 (10)	0.0251 (11)	0.0265 (10)	0.0079 (9)	0.0018 (8)	0.0084 (9)
C211	0.0125 (12)	0.0214 (15)	0.0253 (14)	0.0052 (11)	-0.0033 (11)	0.0037 (12)
N212	0.0272 (12)	0.0358 (15)	0.0220 (12)	0.0161 (11)	-0.0015 (10)	0.0070 (11)
C213	0.0332 (16)	0.044 (2)	0.0234 (15)	0.0229 (15)	0.0014 (12)	0.0042 (14)
C214	0.0262 (15)	0.0336 (19)	0.0426 (19)	0.0162 (14)	0.0004 (14)	-0.0008 (15)
C215	0.0232 (15)	0.0180 (16)	0.057 (2)	0.0052 (12)	0.0057 (14)	0.0069 (15)
C216	0.0195 (14)	0.0267 (17)	0.0400 (17)	0.0058 (12)	0.0060 (12)	0.0104 (14)
C221	0.0225 (14)	0.0214 (15)	0.0208 (14)	0.0089 (12)	0.0039 (11)	0.0053 (12)
N222	0.0250 (12)	0.0265 (14)	0.0348 (13)	0.0087 (11)	0.0012 (10)	0.0141 (11)
C223	0.0295 (16)	0.0307 (18)	0.0421 (18)	0.0093 (14)	0.0003 (13)	0.0191 (15)
C224	0.0234 (15)	0.0388 (19)	0.0399 (18)	0.0134 (14)	-0.0027 (13)	0.0169 (15)
C225	0.0196 (14)	0.0389 (19)	0.0412 (18)	0.0069 (13)	0.0058 (13)	0.0155 (15)
C226	0.0240 (14)	0.0279 (17)	0.0321 (16)	0.0069 (13)	0.0059 (12)	0.0157 (13)
C20	0.0185 (13)	0.0212 (15)	0.0226 (14)	0.0072 (11)	-0.0002 (11)	0.0067 (11)

Geometric parameters (\AA , $^\circ$)

P1—O1	1.4917 (18)	P2—O2	1.4897 (18)
P1—C10	1.799 (3)	P2—C20	1.798 (2)
P1—C121	1.809 (3)	P2—C211	1.811 (3)
P1—C111	1.815 (3)	P2—C221	1.819 (3)
C111—N112	1.344 (3)	C211—N212	1.341 (3)
C111—C116	1.391 (3)	C211—C216	1.384 (4)
N112—C113	1.339 (3)	N212—C213	1.340 (3)
C113—C114	1.381 (4)	C213—C214	1.378 (4)
C113—H113	0.95	C213—H213	0.95
C114—C115	1.377 (4)	C214—C215	1.378 (4)
C114—H114	0.95	C214—H214	0.95
C115—C116	1.380 (4)	C215—C216	1.382 (4)

C115—H115	0.95	C215—H215	0.95
C116—H116	0.95	C216—H216	0.95
C121—N122	1.352 (3)	C221—N222	1.343 (3)
C121—C126	1.383 (4)	C221—C226	1.386 (3)
N122—C123	1.339 (3)	N222—C223	1.336 (3)
C123—C124	1.382 (4)	C223—C224	1.382 (4)
C123—H123	0.95	C223—H223	0.95
C124—C125	1.371 (4)	C224—C225	1.372 (4)
C124—H124	0.95	C224—H224	0.95
C125—C126	1.381 (4)	C225—C226	1.383 (4)
C125—H125	0.95	C225—H225	0.95
C126—H126	0.95	C226—H226	0.95
C10—C10 ⁱ	1.516 (5)	C20—C20 ⁱⁱ	1.536 (5)
C10—H10A	0.99	C20—H20A	0.99
C10—H10B	0.99	C20—H20B	0.99
O1—P1—C10	115.87 (11)	O2—P2—C20	115.28 (11)
O1—P1—C121	112.56 (12)	O2—P2—C211	111.60 (12)
C10—P1—C121	105.88 (12)	C20—P2—C211	106.06 (11)
O1—P1—C111	110.98 (11)	O2—P2—C221	112.89 (11)
C10—P1—C111	105.53 (12)	C20—P2—C221	106.09 (12)
C121—P1—C111	105.23 (11)	C211—P2—C221	104.05 (11)
N112—C111—C116	123.1 (2)	N212—C211—C216	123.4 (2)
N112—C111—P1	116.64 (18)	N212—C211—P2	116.2 (2)
C116—C111—P1	120.2 (2)	C216—C211—P2	120.4 (2)
C113—N112—C111	116.4 (2)	C213—N212—C211	116.5 (2)
N112—C113—C114	124.6 (3)	N212—C213—C214	123.8 (3)
N112—C113—H113	117.7	N212—C213—H213	118.1
C114—C113—H113	117.7	C214—C213—H213	118.1
C115—C114—C113	118.0 (3)	C215—C214—C213	119.1 (3)
C115—C114—H114	121	C215—C214—H214	120.5
C113—C114—H114	121	C213—C214—H214	120.5
C114—C115—C116	119.3 (3)	C214—C215—C216	118.2 (3)
C114—C115—H115	120.4	C214—C215—H215	120.9
C116—C115—H115	120.4	C216—C215—H215	120.9
C115—C116—C111	118.7 (3)	C215—C216—C211	119.0 (3)
C115—C116—H116	120.6	C215—C216—H216	120.5
C111—C116—H116	120.6	C211—C216—H216	120.5
N122—C121—C126	123.0 (2)	N222—C221—C226	123.4 (2)
N122—C121—P1	117.2 (2)	N222—C221—P2	118.23 (18)
C126—C121—P1	119.86 (19)	C226—C221—P2	118.36 (19)
C123—N122—C121	116.5 (2)	C223—N222—C221	116.2 (2)
N122—C123—C124	123.6 (3)	N222—C223—C224	124.3 (3)
N122—C123—H123	118.2	N222—C223—H223	117.8
C124—C123—H123	118.2	C224—C223—H223	117.8
C125—C124—C123	119.3 (3)	C225—C224—C223	118.6 (2)
C125—C124—H124	120.4	C225—C224—H224	120.7
C123—C124—H124	120.4	C223—C224—H224	120.7
C124—C125—C126	118.4 (3)	C224—C225—C226	118.7 (3)

supplementary materials

C124—C125—H125	120.8	C224—C225—H225	120.6
C126—C125—H125	120.8	C226—C225—H225	120.6
C125—C126—C121	119.3 (2)	C225—C226—C221	118.8 (2)
C125—C126—H126	120.4	C225—C226—H226	120.6
C121—C126—H126	120.4	C221—C226—H226	120.6
C10 ⁱ —C10—P1	111.2 (2)	C20 ⁱⁱ —C20—P2	111.1 (2)
C10 ⁱ —C10—H10A	109.4	C20 ⁱⁱ —C20—H20A	109.4
P1—C10—H10A	109.4	P2—C20—H20A	109.4
C10 ⁱ —C10—H10B	109.4	C20 ⁱⁱ —C20—H20B	109.4
P1—C10—H10B	109.4	P2—C20—H20B	109.4
H10A—C10—H10B	108	H20A—C20—H20B	108

Symmetry codes: (i) $-x, -y, -z+1$; (ii) $-x+1, -y, -z$.

Fig. 1

