

n-Undecanyl 2-(4-chloroanilino)-4,4-dimethyl-6-oxocyclohex-1-enecarbo-dithioate

El Sayed H. El Ashry,^a Mohammed R. Amer,^a M. Raza Shah^a and Seik Weng Ng^{b*}

^aH.E.J. Research Institute of Chemistry, International Center for Chemical and Biological Sciences, University of Karachi, Karachi 75270, Pakistan, and

^bDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia
Correspondence e-mail: seikweng@um.edu.my


Received 19 February 2009; accepted 19 February 2009

Key indicators: single-crystal X-ray study; $T = 100\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$; R factor = 0.052; wR factor = 0.157; data-to-parameter ratio = 21.3.

The six-membered cyclohexene ring in the title compound, $\text{C}_{26}\text{H}_{38}\text{ClNO}_2$, adopts an envelope conformation, with the C atom bearing the two methyl groups representing the flap. This atom deviates by $0.642(4)\text{ \AA}$ from the plane passing through the other five atoms of the ring (r.m.s. deviation = 0.053 \AA). The molecular conformation is stabilized by an intramolecular N—H···S hydrogen bond.

Related literature

For background, see: El Ashry *et al.* (2009).



Experimental

Crystal data

$\text{C}_{26}\text{H}_{38}\text{ClNO}_2$
 $M_r = 480.14$
Triclinic, $P\bar{1}$
 $a = 7.9865(2)\text{ \AA}$
 $b = 11.8015(4)\text{ \AA}$
 $c = 14.9676(4)\text{ \AA}$
 $\alpha = 97.522(2)^\circ$
 $\beta = 94.004(3)^\circ$
 $\gamma = 106.134(2)^\circ$
 $V = 1335.13(7)\text{ \AA}^3$
 $Z = 2$
Mo $K\alpha$ radiation
 $\mu = 0.32\text{ mm}^{-1}$
 $T = 100\text{ K}$
 $0.20 \times 0.15 \times 0.10\text{ mm}$

Data collection

Bruker SMART APEX
diffractometer
Absorption correction: multi-scan
(SADABS; Sheldrick, 1996)
 $T_{\min} = 0.892$, $T_{\max} = 0.969$
12760 measured reflections
6108 independent reflections
3899 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.051$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.052$
 $wR(F^2) = 0.157$
 $S = 1.01$
6108 reflections
287 parameters
1 restraint
H atoms treated by a mixture of
independent and constrained
refinement
 $\Delta\rho_{\max} = 0.33\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.36\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N1—H1···S2	0.89 (1)	2.10 (2)	2.873 (2)	146 (3)

Data collection: *APEX2* (Bruker, 2008); cell refinement: *SAINT* (Bruker, 2008); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2009).

We thank the Higher Education Commission of Pakistan and the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT2878).

References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
- Bruker (2008). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- El Ashry, E. S. H., Amer, M. R., Raza Shah, M. & Ng, S. W. (2009). *Acta Cryst. E* **65**, o597.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Westrip, S. P. (2009). *publCIF*. In preparation.

supplementary materials

Acta Cryst. (2009). E65, o599 [doi:10.1107/S1600536809006163]

n-Undecanyl 2-(4-chloroanilino)-4,4-dimethyl-6-oxocyclohex-1-enecarbodithioate

E. S. H. El Ashry, M. R. Amer, M. R. Shah and S. W. Ng

Comment

(type here to add)

Experimental

To a solution of (4-chlorophenylamino)-5,5-dimethyl-cyclohex-2-en-1-one (0.1 mol) in DMSO (20 ml) and sodium hydroxide (0.4 g) in water (1 ml), carbon disulfphide (0.3 mol) was added in the course of 30 minutes. The mixture was stirred for 20 min below 283 K, and then 1-bromoundecane (0.1 mol) was added dropwise at room temperature for 30 min. The reaction mixture was left for 24 h and then diluted with water (200 ml) and acidified with 10% hydrochloric acid. The resulting precipitate was collected by filtration, dried and purified on silica gel column (40% ethyl acetate in hexane) to give yellow crystals (48% yield; mp 368 K).

Refinement

Carbon-bound H-atoms were placed in calculated positions (C—H 0.93 to 0.99 Å) and were included in the refinement in the riding model approximation, with $U(H)$ set to 1.2 to 1.5 $U(C)$. The methyl groups were allowed to rotate but not to tip.

The amino H-atom was located in a difference Fourier map, and was refined with a distance restraint of N—H 0.88±0.01 Å; its isotropic displacement parameter was freely refined.

Figures




Fig. 1. Anisotropic displacement ellipsoid plot (Barbour, 2001) plot of $C_{26}H_{38}ClNO_2S$ at the 70% probability level; hydrogen atoms are drawn as spheres of arbitrary radius.

n-Undecanyl 2-(4-chloroanilino)-4,4-dimethyl-6-oxocyclohex-1-enecarbodithioate

Crystal data

$C_{26}H_{38}ClNO_2S$

$Z = 2$

$M_r = 480.14$

$F_{000} = 516$

Triclinic, $P\bar{1}$

$D_x = 1.194 \text{ Mg m}^{-3}$

Hall symbol: -P 1

Mo $K\alpha$ radiation

$a = 7.9865 (2) \text{ \AA}$

$\lambda = 0.71073 \text{ \AA}$

Cell parameters from 1537 reflections

supplementary materials

$b = 11.8015 (4) \text{ \AA}$	$\theta = 2.5\text{--}22.0^\circ$
$c = 14.9676 (4) \text{ \AA}$	$\mu = 0.32 \text{ mm}^{-1}$
$\alpha = 97.522 (2)^\circ$	$T = 100 \text{ K}$
$\beta = 94.004 (3)^\circ$	Irregular block, yellow
$\gamma = 106.134 (2)^\circ$	$0.20 \times 0.15 \times 0.10 \text{ mm}$
$V = 1335.13 (7) \text{ \AA}^3$	

Data collection

Bruker SMART APEX diffractometer	6108 independent reflections
Radiation source: fine-focus sealed tube	3899 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.051$
$T = 100 \text{ K}$	$\theta_{\text{max}} = 27.5^\circ$
ω scans	$\theta_{\text{min}} = 1.4^\circ$
Absorption correction: Multi-scan (SADABS; Sheldrick, 1996)	$h = -10 \rightarrow 10$
$T_{\text{min}} = 0.892, T_{\text{max}} = 0.969$	$k = -15 \rightarrow 15$
12760 measured reflections	$l = -19 \rightarrow 19$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.052$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.157$	$w = 1/[\sigma^2(F_o^2) + (0.074P)^2]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.01$	$(\Delta/\sigma)_{\text{max}} = 0.001$
6108 reflections	$\Delta\rho_{\text{max}} = 0.33 \text{ e \AA}^{-3}$
287 parameters	$\Delta\rho_{\text{min}} = -0.36 \text{ e \AA}^{-3}$
1 restraint	Extinction correction: none
Primary atom site location: structure-invariant direct methods	

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Cl1	0.19825 (13)	0.41121 (9)	1.03994 (6)	0.0563 (3)
S1	0.63724 (9)	0.24958 (6)	0.39696 (5)	0.02316 (18)
S2	0.39872 (10)	0.15853 (6)	0.52890 (5)	0.0290 (2)
N1	0.4552 (3)	0.3457 (2)	0.68108 (16)	0.0254 (5)
H1	0.416 (4)	0.2718 (14)	0.651 (2)	0.050 (11)*
O1	0.7692 (3)	0.47573 (16)	0.43665 (13)	0.0286 (5)
C1	0.7341 (3)	0.4897 (2)	0.51557 (18)	0.0212 (6)
C2	0.8197 (4)	0.6096 (2)	0.57245 (18)	0.0244 (6)
H2A	0.7635	0.6678	0.5514	0.029*

H2B	0.9444	0.6354	0.5613	0.029*
C3	0.8122 (3)	0.6156 (2)	0.67421 (18)	0.0230 (6)
C4	0.6254 (3)	0.5507 (2)	0.68855 (18)	0.0221 (6)
H4A	0.6187	0.5495	0.7542	0.027*
H4B	0.5464	0.5960	0.6681	0.027*
C5	0.5613 (3)	0.4250 (2)	0.63920 (18)	0.0230 (6)
C6	0.6116 (3)	0.3932 (2)	0.55220 (18)	0.0218 (6)
C7	0.9411 (4)	0.5570 (3)	0.71535 (19)	0.0300 (7)
H7A	1.0607	0.5995	0.7053	0.045*
H7B	0.9125	0.4734	0.6865	0.045*
H7C	0.9329	0.5603	0.7806	0.045*
C8	0.8553 (4)	0.7455 (3)	0.7191 (2)	0.0335 (7)
H8A	0.7681	0.7812	0.6950	0.050*
H8B	0.9721	0.7897	0.7063	0.050*
H8C	0.8532	0.7491	0.7847	0.050*
C9	0.5498 (3)	0.2748 (2)	0.49969 (18)	0.0228 (6)
C10	0.5423 (4)	0.0905 (2)	0.36058 (19)	0.0275 (6)
H10A	0.5528	0.0470	0.4120	0.033*
H10B	0.4163	0.0727	0.3390	0.033*
C11	0.6397 (4)	0.0507 (3)	0.28414 (18)	0.0278 (6)
H11A	0.5910	-0.0368	0.2659	0.033*
H11B	0.7645	0.0673	0.3077	0.033*
C12	0.6303 (4)	0.1104 (3)	0.20027 (19)	0.0297 (7)
H12A	0.5057	0.0963	0.1776	0.036*
H12B	0.6836	0.1976	0.2176	0.036*
C13	0.7230 (4)	0.0651 (3)	0.12447 (19)	0.0297 (7)
H13A	0.6651	-0.0213	0.1047	0.036*
H13B	0.8458	0.0746	0.1482	0.036*
C14	0.7234 (4)	0.1293 (3)	0.04282 (19)	0.0319 (7)
H14A	0.6006	0.1169	0.0176	0.038*
H14B	0.7767	0.2160	0.0632	0.038*
C15	0.8222 (4)	0.0882 (3)	-0.03191 (19)	0.0334 (7)
H15A	0.7756	0.0004	-0.0489	0.040*
H15B	0.9474	0.1070	-0.0084	0.040*
C16	0.8086 (4)	0.1459 (3)	-0.11602 (19)	0.0305 (7)
H16A	0.8572	0.2336	-0.0990	0.037*
H16B	0.6832	0.1285	-0.1386	0.037*
C17	0.9041 (4)	0.1038 (3)	-0.19232 (19)	0.0312 (7)
H17A	1.0307	0.1254	-0.1709	0.037*
H17B	0.8603	0.0156	-0.2069	0.037*
C18	0.8816 (4)	0.1564 (3)	-0.27808 (19)	0.0294 (7)
H18A	0.9328	0.2442	-0.2644	0.035*
H18B	0.7547	0.1396	-0.2972	0.035*
C19	0.9668 (4)	0.1080 (3)	-0.35609 (19)	0.0323 (7)
H19A	1.0946	0.1280	-0.3383	0.039*
H19B	0.9193	0.0198	-0.3683	0.039*
C20	0.9360 (5)	0.1577 (3)	-0.4419 (2)	0.0421 (8)
H20A	0.9942	0.1245	-0.4899	0.063*
H20B	0.9841	0.2449	-0.4305	0.063*

supplementary materials

H20C	0.8098	0.1360	-0.4610	0.063*
C21	0.3962 (4)	0.3664 (3)	0.76804 (19)	0.0264 (6)
C22	0.2856 (4)	0.4365 (3)	0.7835 (2)	0.0357 (8)
H22	0.2508	0.4750	0.7366	0.043*
C23	0.2256 (4)	0.4507 (3)	0.8673 (2)	0.0420 (8)
H23	0.1499	0.4992	0.8783	0.050*
C24	0.2753 (4)	0.3946 (3)	0.9345 (2)	0.0347 (7)
C25	0.3846 (5)	0.3240 (3)	0.9201 (2)	0.0427 (8)
H25	0.4186	0.2856	0.9673	0.051*
C26	0.4448 (4)	0.3095 (3)	0.8357 (2)	0.0375 (8)
H26	0.5196	0.2603	0.8246	0.045*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cl1	0.0583 (6)	0.0751 (7)	0.0282 (4)	0.0072 (5)	0.0165 (4)	0.0020 (4)
S1	0.0251 (4)	0.0190 (4)	0.0240 (4)	0.0041 (3)	0.0038 (3)	0.0034 (3)
S2	0.0293 (4)	0.0188 (4)	0.0367 (4)	0.0013 (3)	0.0120 (3)	0.0042 (3)
N1	0.0269 (13)	0.0211 (13)	0.0272 (13)	0.0033 (10)	0.0097 (10)	0.0046 (11)
O1	0.0343 (12)	0.0214 (10)	0.0270 (11)	0.0011 (8)	0.0093 (9)	0.0049 (8)
C1	0.0183 (14)	0.0177 (13)	0.0273 (15)	0.0042 (11)	0.0019 (11)	0.0047 (11)
C2	0.0249 (15)	0.0189 (14)	0.0279 (15)	0.0030 (11)	0.0029 (12)	0.0057 (12)
C3	0.0183 (14)	0.0236 (14)	0.0250 (14)	0.0032 (11)	0.0022 (11)	0.0029 (12)
C4	0.0216 (14)	0.0201 (14)	0.0246 (14)	0.0058 (11)	0.0032 (11)	0.0032 (11)
C5	0.0196 (14)	0.0222 (14)	0.0283 (15)	0.0070 (11)	-0.0006 (11)	0.0073 (12)
C6	0.0198 (14)	0.0220 (14)	0.0235 (14)	0.0059 (11)	0.0011 (11)	0.0047 (11)
C7	0.0241 (15)	0.0379 (18)	0.0269 (15)	0.0068 (13)	0.0003 (12)	0.0074 (13)
C8	0.0330 (17)	0.0254 (16)	0.0369 (17)	0.0019 (13)	0.0055 (14)	0.0002 (13)
C9	0.0191 (14)	0.0242 (14)	0.0277 (15)	0.0091 (11)	0.0026 (11)	0.0065 (12)
C10	0.0314 (16)	0.0199 (15)	0.0272 (15)	0.0008 (12)	0.0024 (12)	0.0041 (12)
C11	0.0332 (17)	0.0244 (15)	0.0253 (15)	0.0097 (12)	0.0011 (12)	0.0004 (12)
C12	0.0337 (17)	0.0267 (16)	0.0270 (16)	0.0070 (13)	0.0015 (13)	0.0031 (12)
C13	0.0298 (16)	0.0285 (16)	0.0271 (16)	0.0060 (13)	-0.0014 (12)	-0.0013 (13)
C14	0.0371 (18)	0.0296 (16)	0.0280 (16)	0.0092 (13)	0.0035 (13)	0.0024 (13)
C15	0.0361 (18)	0.0376 (18)	0.0279 (16)	0.0133 (14)	0.0041 (14)	0.0039 (14)
C16	0.0324 (17)	0.0280 (16)	0.0305 (16)	0.0094 (13)	0.0029 (13)	0.0017 (13)
C17	0.0295 (17)	0.0364 (18)	0.0276 (16)	0.0101 (13)	0.0026 (13)	0.0043 (13)
C18	0.0297 (16)	0.0261 (16)	0.0322 (16)	0.0072 (13)	0.0042 (13)	0.0058 (13)
C19	0.0274 (16)	0.0376 (18)	0.0333 (17)	0.0079 (13)	0.0072 (13)	0.0117 (14)
C20	0.044 (2)	0.051 (2)	0.0373 (19)	0.0168 (17)	0.0141 (16)	0.0157 (16)
C21	0.0242 (15)	0.0273 (16)	0.0250 (15)	0.0036 (12)	0.0038 (12)	0.0029 (12)
C22	0.0321 (17)	0.053 (2)	0.0291 (16)	0.0205 (15)	0.0067 (13)	0.0119 (15)
C23	0.0337 (18)	0.065 (2)	0.0345 (18)	0.0235 (17)	0.0114 (14)	0.0077 (17)
C24	0.0323 (17)	0.0421 (19)	0.0229 (15)	0.0004 (14)	0.0097 (13)	0.0004 (14)
C25	0.060 (2)	0.040 (2)	0.0296 (17)	0.0152 (17)	0.0037 (16)	0.0102 (15)
C26	0.049 (2)	0.0312 (18)	0.0371 (18)	0.0181 (15)	0.0081 (15)	0.0079 (14)

Geometric parameters (Å, °)

C11—C24	1.740 (3)	C12—H12B	0.9900
S1—C9	1.757 (3)	C13—C14	1.520 (4)
S1—C10	1.811 (3)	C13—H13A	0.9900
S2—C9	1.685 (3)	C13—H13B	0.9900
N1—C5	1.332 (3)	C14—C15	1.517 (4)
N1—C21	1.429 (3)	C14—H14A	0.9900
N1—H1	0.885 (10)	C14—H14B	0.9900
O1—C1	1.234 (3)	C15—C16	1.519 (4)
C1—C6	1.474 (4)	C15—H15A	0.9900
C1—C2	1.504 (4)	C15—H15B	0.9900
C2—C3	1.522 (4)	C16—C17	1.522 (4)
C2—H2A	0.9900	C16—H16A	0.9900
C2—H2B	0.9900	C16—H16B	0.9900
C3—C4	1.520 (4)	C17—C18	1.518 (4)
C3—C8	1.525 (4)	C17—H17A	0.9900
C3—C7	1.529 (4)	C17—H17B	0.9900
C4—C5	1.499 (4)	C18—C19	1.519 (4)
C4—H4A	0.9900	C18—H18A	0.9900
C4—H4B	0.9900	C18—H18B	0.9900
C5—C6	1.423 (4)	C19—C20	1.514 (4)
C6—C9	1.446 (4)	C19—H19A	0.9900
C7—H7A	0.9800	C19—H19B	0.9900
C7—H7B	0.9800	C20—H20A	0.9800
C7—H7C	0.9800	C20—H20B	0.9800
C8—H8A	0.9800	C20—H20C	0.9800
C8—H8B	0.9800	C21—C26	1.376 (4)
C8—H8C	0.9800	C21—C22	1.380 (4)
C10—C11	1.524 (4)	C22—C23	1.380 (4)
C10—H10A	0.9900	C22—H22	0.9500
C10—H10B	0.9900	C23—C24	1.368 (4)
C11—C12	1.525 (4)	C23—H23	0.9500
C11—H11A	0.9900	C24—C25	1.373 (5)
C11—H11B	0.9900	C25—C26	1.390 (4)
C12—C13	1.516 (4)	C25—H25	0.9500
C12—H12A	0.9900	C26—H26	0.9500
C9—S1—C10	102.95 (13)	C12—C13—H13A	108.9
C5—N1—C21	127.5 (2)	C14—C13—H13A	108.9
C5—N1—H1	116 (2)	C12—C13—H13B	108.9
C21—N1—H1	116 (2)	C14—C13—H13B	108.9
O1—C1—C6	121.4 (2)	H13A—C13—H13B	107.7
O1—C1—C2	117.2 (2)	C15—C14—C13	114.0 (3)
C6—C1—C2	121.4 (2)	C15—C14—H14A	108.8
C1—C2—C3	116.1 (2)	C13—C14—H14A	108.8
C1—C2—H2A	108.3	C15—C14—H14B	108.8
C3—C2—H2A	108.3	C13—C14—H14B	108.8
C1—C2—H2B	108.3	H14A—C14—H14B	107.7

supplementary materials

C3—C2—H2B	108.3	C14—C15—C16	113.2 (3)
H2A—C2—H2B	107.4	C14—C15—H15A	108.9
C4—C3—C2	107.3 (2)	C16—C15—H15A	108.9
C4—C3—C8	108.7 (2)	C14—C15—H15B	108.9
C2—C3—C8	109.7 (2)	C16—C15—H15B	108.9
C4—C3—C7	110.5 (2)	H15A—C15—H15B	107.7
C2—C3—C7	110.7 (2)	C15—C16—C17	113.9 (2)
C8—C3—C7	109.9 (2)	C15—C16—H16A	108.8
C5—C4—C3	113.6 (2)	C17—C16—H16A	108.8
C5—C4—H4A	108.8	C15—C16—H16B	108.8
C3—C4—H4A	108.8	C17—C16—H16B	108.8
C5—C4—H4B	108.8	H16A—C16—H16B	107.7
C3—C4—H4B	108.8	C18—C17—C16	113.7 (2)
H4A—C4—H4B	107.7	C18—C17—H17A	108.8
N1—C5—C6	122.3 (3)	C16—C17—H17A	108.8
N1—C5—C4	116.4 (2)	C18—C17—H17B	108.8
C6—C5—C4	121.3 (2)	C16—C17—H17B	108.8
C5—C6—C9	124.1 (2)	H17A—C17—H17B	107.7
C5—C6—C1	116.2 (2)	C17—C18—C19	113.5 (2)
C9—C6—C1	119.7 (2)	C17—C18—H18A	108.9
C3—C7—H7A	109.5	C19—C18—H18A	108.9
C3—C7—H7B	109.5	C17—C18—H18B	108.9
H7A—C7—H7B	109.5	C19—C18—H18B	108.9
C3—C7—H7C	109.5	H18A—C18—H18B	107.7
H7A—C7—H7C	109.5	C20—C19—C18	112.5 (3)
H7B—C7—H7C	109.5	C20—C19—H19A	109.1
C3—C8—H8A	109.5	C18—C19—H19A	109.1
C3—C8—H8B	109.5	C20—C19—H19B	109.1
H8A—C8—H8B	109.5	C18—C19—H19B	109.1
C3—C8—H8C	109.5	H19A—C19—H19B	107.8
H8A—C8—H8C	109.5	C19—C20—H20A	109.5
H8B—C8—H8C	109.5	C19—C20—H20B	109.5
C6—C9—S2	125.1 (2)	H20A—C20—H20B	109.5
C6—C9—S1	117.8 (2)	C19—C20—H20C	109.5
S2—C9—S1	117.15 (16)	H20A—C20—H20C	109.5
C11—C10—S1	108.50 (19)	H20B—C20—H20C	109.5
C11—C10—H10A	110.0	C26—C21—C22	120.1 (3)
S1—C10—H10A	110.0	C26—C21—N1	118.4 (3)
C11—C10—H10B	110.0	C22—C21—N1	121.4 (3)
S1—C10—H10B	110.0	C23—C22—C21	119.8 (3)
H10A—C10—H10B	108.4	C23—C22—H22	120.1
C10—C11—C12	114.8 (2)	C21—C22—H22	120.1
C10—C11—H11A	108.6	C24—C23—C22	119.9 (3)
C12—C11—H11A	108.6	C24—C23—H23	120.1
C10—C11—H11B	108.6	C22—C23—H23	120.1
C12—C11—H11B	108.6	C23—C24—C25	121.1 (3)
H11A—C11—H11B	107.5	C23—C24—Cl1	120.0 (3)
C13—C12—C11	113.2 (2)	C25—C24—Cl1	119.0 (3)
C13—C12—H12A	108.9	C24—C25—C26	119.1 (3)

C11—C12—H12A	108.9	C24—C25—H25	120.4
C13—C12—H12B	108.9	C26—C25—H25	120.4
C11—C12—H12B	108.9	C21—C26—C25	120.0 (3)
H12A—C12—H12B	107.7	C21—C26—H26	120.0
C12—C13—C14	113.5 (2)	C25—C26—H26	120.0
O1—C1—C2—C3	−165.2 (2)	C10—S1—C9—C6	174.7 (2)
C6—C1—C2—C3	15.6 (4)	C10—S1—C9—S2	−5.8 (2)
C1—C2—C3—C4	−45.9 (3)	C9—S1—C10—C11	−166.72 (19)
C1—C2—C3—C8	−163.9 (2)	S1—C10—C11—C12	−60.6 (3)
C1—C2—C3—C7	74.7 (3)	C10—C11—C12—C13	−177.9 (2)
C2—C3—C4—C5	55.4 (3)	C11—C12—C13—C14	−176.6 (3)
C8—C3—C4—C5	174.0 (2)	C12—C13—C14—C15	177.6 (3)
C7—C3—C4—C5	−65.3 (3)	C13—C14—C15—C16	175.3 (3)
C21—N1—C5—C6	179.8 (3)	C14—C15—C16—C17	−178.9 (3)
C21—N1—C5—C4	−0.5 (4)	C15—C16—C17—C18	176.8 (3)
C3—C4—C5—N1	144.5 (2)	C16—C17—C18—C19	−176.1 (2)
C3—C4—C5—C6	−35.7 (3)	C17—C18—C19—C20	177.6 (3)
N1—C5—C6—C9	2.0 (4)	C5—N1—C21—C26	−116.3 (3)
C4—C5—C6—C9	−177.8 (2)	C5—N1—C21—C22	67.0 (4)
N1—C5—C6—C1	−178.1 (2)	C26—C21—C22—C23	0.8 (5)
C4—C5—C6—C1	2.1 (4)	N1—C21—C22—C23	177.4 (3)
O1—C1—C6—C5	−171.1 (2)	C21—C22—C23—C24	−0.3 (5)
C2—C1—C6—C5	8.2 (4)	C22—C23—C24—C25	0.0 (5)
O1—C1—C6—C9	8.8 (4)	C22—C23—C24—Cl1	−179.4 (3)
C2—C1—C6—C9	−172.0 (2)	C23—C24—C25—C26	−0.1 (5)
C5—C6—C9—S2	4.6 (4)	Cl1—C24—C25—C26	179.3 (3)
C1—C6—C9—S2	−175.2 (2)	C22—C21—C26—C25	−0.9 (5)
C5—C6—C9—S1	−175.8 (2)	N1—C21—C26—C25	−177.6 (3)
C1—C6—C9—S1	4.3 (3)	C24—C25—C26—C21	0.6 (5)

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
N1—H1···S2	0.89 (1)	2.10 (2)	2.873 (2)	146 (3)

supplementary materials

Fig. 1

