

Tetra(chlorido/iodido)(1,10-phenanthroline)platinum(IV) hemi[di(chlorine/iodine)]

Nam-Ho Kim and Kwang Ha*

School of Applied Chemical Engineering, The Research Institute of Catalysis, Chonnam National University, Gwangju 500-757, Republic of Korea
Correspondence e-mail: hakwang@chonnam.ac.kr


Received 14 February 2009; accepted 3 March 2009

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.010$ Å; disorder in main residue; R factor = 0.031; wR factor = 0.062; data-to-parameter ratio = 16.7.

The asymmetric unit of the title compound, $[\text{PtCl}_{3.66}\text{I}_{0.34}(\text{C}_{12}\text{H}_8\text{N}_2)] \cdot 0.5(\text{Cl}_{0.13}\text{I}_{1.87})$, contains a neutral Pt^{IV} complex and one half of a halogen molecule. The Pt^{IV} ion is six-coordinated in a distorted octahedral environment by two N atoms of the 1,10-phenanthroline ligand and Cl or I atoms. The refinement of the structure and the EDX analysis indicate that the compound is a solid solution in which there is some substitution of Cl for I and *vice versa*. The chemical formula of the pure state of the compound would have been $[\text{PtCl}_4(\text{C}_{12}\text{H}_8\text{N}_2)] \cdot 0.5\text{I}_2$. In the analysed crystal, two Cl atoms are partially (*ca* 25% and 9%) replaced by I atoms, and the I_2 molecule has a minor component modelled as ICl. As a result of the disorder, the different *trans* effects of the N and Cl/I atoms are not distinct. The complex displays intermolecular π - π interactions between the six-membered rings, with a centroid-centroid distance of 3.771 (4) Å. There are also weak intramolecular $\text{C}-\text{H} \cdots \text{Cl}$ hydrogen bonds.

Related literature

For details of some other Pt-phenanthroline complexes, see: Buse *et al.* (1977); Fanizzi *et al.* (1996); Kim *et al.* (2009*a,b*). For related Pt-bipyridine complexes, see: Hambley (1986); Hojjat Kashani *et al.* (2008). For bond-length data, see: Orpen *et al.* (1989).



Experimental

Crystal data

$[\text{PtCl}_{3.66}\text{I}_{0.34}(\text{C}_{12}\text{H}_8\text{N}_2)] \cdot 0.5(\text{Cl}_{0.13}\text{I}_{1.87})$	$V = 3180.8$ (19) Å ³
$M_r = 669.26$	$Z = 8$
Orthorhombic, <i>Pbca</i>	Mo $K\alpha$ radiation
$a = 14.215$ (5) Å	$\mu = 11.92$ mm ⁻¹
$b = 12.733$ (5) Å	$T = 293$ K
$c = 17.575$ (6) Å	$0.25 \times 0.17 \times 0.15$ mm

Data collection

Bruker SMART 1000 CCD diffractometer	17485 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 2000)	3246 independent reflections
$T_{\min} = 0.111$, $T_{\max} = 0.168$	2169 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.049$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.031$	8 restraints
$wR(F^2) = 0.062$	H-atom parameters constrained
$S = 0.86$	$\Delta\rho_{\text{max}} = 1.29$ e Å ⁻³
3246 reflections	$\Delta\rho_{\text{min}} = -0.50$ e Å ⁻³
194 parameters	

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H} \cdots A$	$D-\text{H}$	$\text{H} \cdots A$	$D \cdots A$	$D-\text{H} \cdots A$
$\text{Cl1}-\text{H1} \cdots \text{Cl2}$	0.93	2.73	3.320 (8)	122
$\text{Cl10}-\text{H10} \cdots \text{Cl1}$	0.93	2.66	3.240 (7)	121

Data collection: *SMART* (Bruker, 2000); cell refinement: *SAINT* (Bruker, 2000); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2009); software used to prepare material for publication: *SHELXL97*.

This work was supported by a Korea Research Foundation grant funded by the Korean Government (MOEHRD) (KRF-2007-412-J02001).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: PK2156).

References

- Bruker (2000). *SADABS*, *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Buse, K. D., Keller, H. J. & Pritzkow, H. (1977). *Inorg. Chem.* **16**, 1072–1076.
- Fanizzi, F. P., Natile, G., Lanfranchi, M., Tiripicchio, A., Laschi, F. & Zanello, P. (1996). *Inorg. Chem.* **35**, 3173–3182.
- Hambley, T. W. (1986). *Acta Cryst.* **C42**, 49–51.
- Hojjat Kashani, L., Amani, V., Yousefi, M. & Khavasi, H. R. (2008). *Acta Cryst.* **E64**, m905–m906.
- Kim, N.-H., Hwang, I.-C. & Ha, K. (2009*a*). *Acta Cryst.* **E65**, m224.
- Kim, N.-H., Hwang, I.-C. & Ha, K. (2009*b*). *Acta Cryst.* **E65**, m230.
- Orpen, G., Brammer, L., Allen, F. H., Kennard, O., Watson, D. G. & Taylor, R. (1989). *J. Chem. Soc. Dalton Trans.* pp. S1–83.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

supplementary materials

Acta Cryst. (2009). E65, m379 [doi:10.1107/S1600536809007703]

Tetra(chlorido/iodido)(1,10-phenanthroline)platinum(IV) hemi[di(chlorine/iodine)]

N.-H. Kim and K. Ha

Comment

The asymmetric unit of the title compound contains a neutral Pt^{IV} complex and one half-molecule of iodine which includes some Cl atoms (*ca* 6%). The Pt^{IV} ion is six-coordinated in a distorted octahedral environment by two N atoms of the 1,10-phenanthroline ligand and Cl or I atoms. The chemical formula of the pure state of the title compound would have been [PtCl₄(C₁₂H₈N₂)]_{0.5}I₂. In the particular crystal of the compound used, two Cl atoms (Cl3 and Cl4) are partially (*ca* 25% and 9%, respectively) displaced by the I atoms (I3 and I4) through the substitution reaction between the Cl⁻ and I⁻ ligand, and the I₂ molecule also appears to have a minor component, that is I—Cl (Fig. 1 and 2). The chemical formula which resulted from the refinement of the structure was [PtCl_{3.66}I_{0.34}(C₁₂H₈N₂)]_{0.5}(Cl_{0.13}I_{1.87}), and in this case the ratio of the Cl atom to I atom is 2.91:1. An EDX analysis of the compound, however, gave a ratio of Cl:I = 2.47:1. Accordingly, the exact composition may very well be variable, and likely dependent on the exact conditions present during crystal formation. Even though these data are slightly different, they indicate clearly that the crystals are a solid solution in which there was some substitution of Cl for I and *vice versa*.

As a result of the different *trans* effects of the N and Cl atoms, the Pt—Cl bonds *trans* to the N atom are in general slightly shorter than bond lengths to mutually *trans* Cl atoms (Kim *et al.* 2009*a* and 2009*b*). But the *trans* effects of the N and Cl/I atoms in the crystal are not distinct owing to the disordered atoms. The Pt—I distance is restrained to the value given in table 9.6.3.3 of the International Tables Vol. C (Orpen *et al.*, 1989) (2.658 Å). The main contributor to the distortion from a true octahedral structure is the tight N1—Pt1—N2 chelate angle (81.3 (2)°), which result in non-linear *trans* axes (<Cl1—Pt1—N1 = 174.14 (16)° and <Cl2—Pt1—N2 = 175.97 (17)°). The complex displays intermolecular π - π interactions between the six-membered rings, with a shortest centroid-centroid distance of 3.771 (4) Å and with a dihedral angle between the ring planes of 2.1 (3)°. There are also weak intramolecular C—H \cdots Cl hydrogen bonds (Table 1).

The iodine molecule was presumably formed as a consequence of the oxidation of the iodide ion by the Pt⁴⁺ ion, and crystallized with the partially substituted complex. The bond distance between the I atoms is 2.708 (2) Å.

Experimental

To a solution of [PtCl₄(C₁₂H₈N₂)]_{0.5}H₂O (0.0821 g, 0.153 mmol) in H₂O (20 ml) was added KI (0.1318 g, 0.794 mmol), and stirred for 2 h at room temperature. The precipitate was separated by filtration and washed with water (20 ml) and MeOH (20 ml) and dried under vacuum, to give a dark brown powder (0.0846 g). Black crystals suitable for X-ray analysis were isolated from an acetone solution of the reaction products. EDX analysis (%atom): Cl 62.30%, I 25.18%, Pt 12.52%.

Refinement

H atoms were positioned geometrically and allowed to ride on their respective parent atoms [C—H = 0.93 Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$]. The disordered Cl5 atom was refined isotropically. Eight restraints instructions were used for the refinement using the following *SHELXL97* (Sheldrick, 2008) commands: EADP Cl3 I3 and Cl4 I4, SIMU 0.010 I1 Cl5, BIND I1 Cl5a and Cl5 I1a, FREE Cl5 Cl5a, *DFIX* 2.658 0.010 Pt1 I3 and Pt1 I4.

Figures




Fig. 1. The disordered structure of the title compound, with displacement ellipsoids drawn at the 30% probability level for non-H atoms [Symmetry code: (a) 1 - x, 1 - y, -z]. The minor bonds are drawn with dashed lines.




Fig. 2. View of a packing detail of the title compound. For the sake of clarity, only the major disorder component is shown.

Chloridoiodido(1,10-phenanthroline)platinum(IV) hemi[di(chloride/iodide)]

Crystal data

[PtCl_{3.66}I_{0.34}(C₁₂H₈N₂)]·0.5(Cl_{0.13}I_{1.87})

$M_r = 669.26$

Orthorhombic, *Pbca*

Hall symbol: -P 2ac 2ab

$a = 14.215 (5) \text{ \AA}$

$b = 12.733 (5) \text{ \AA}$

$c = 17.575 (6) \text{ \AA}$

$V = 3180.8 (19) \text{ \AA}^3$

$Z = 8$

$F_{000} = 2424$

$D_x = 2.795 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation

$\lambda = 0.71073 \text{ \AA}$

Cell parameters from 973 reflections

$\theta = 2.4\text{--}24.5^\circ$

$\mu = 11.92 \text{ mm}^{-1}$

$T = 293 \text{ K}$

Plate, black

$0.25 \times 0.17 \times 0.15 \text{ mm}$

Data collection

Bruker SMART 1000 CCD
diffractometer

Radiation source: fine-focus sealed tube

3246 independent reflections

2169 reflections with $I > 2\sigma(I)$

Monochromator: graphite $R_{\text{int}} = 0.049$
 $T = 293$ K $\theta_{\text{max}} = 26.4^\circ$
 φ and ω scans $\theta_{\text{min}} = 2.3^\circ$
 Absorption correction: Multi-scan (SADABS; Bruker, 2000) $h = -15 \rightarrow 17$
 $T_{\text{min}} = 0.111$, $T_{\text{max}} = 0.168$ $k = -10 \rightarrow 15$
 17485 measured reflections $l = -17 \rightarrow 21$

Refinement

Refinement on F^2 Secondary atom site location: difference Fourier map
 Least-squares matrix: full Hydrogen site location: inferred from neighbouring sites
 $R[F^2 > 2\sigma(F^2)] = 0.031$ H-atom parameters constrained
 $wR(F^2) = 0.062$ $w = 1/[\sigma^2(F_o^2) + (0.0244P)^2]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $S = 0.86$ $(\Delta/\sigma)_{\text{max}} = 0.001$
 3246 reflections $\Delta\rho_{\text{max}} = 1.29 \text{ e } \text{\AA}^{-3}$
 194 parameters $\Delta\rho_{\text{min}} = -0.50 \text{ e } \text{\AA}^{-3}$
 8 restraints Extinction correction: none
 Primary atom site location: structure-invariant direct methods

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Pt1	-0.134096 (18)	0.30394 (2)	0.177609 (15)	0.03991 (10)	
Cl1	-0.17047 (13)	0.14177 (15)	0.22685 (10)	0.0521 (5)	
Cl2	-0.28869 (12)	0.36077 (15)	0.19550 (10)	0.0512 (5)	
Cl3	-0.0903 (14)	0.3607 (17)	0.2949 (8)	0.0494 (13)	0.746 (3)
I3	-0.0759 (11)	0.3695 (13)	0.3094 (6)	0.0494 (13)	0.254 (3)
Cl4	-0.1730 (7)	0.2478 (8)	0.0572 (3)	0.0518 (8)	0.913 (3)
I4	-0.177 (2)	0.238 (2)	0.0398 (9)	0.0518 (8)	0.087 (3)
N1	-0.0875 (4)	0.4446 (4)	0.1355 (3)	0.0393 (14)	
N2	0.0029 (4)	0.2642 (4)	0.1585 (3)	0.0388 (13)	
C1	-0.1341 (5)	0.5328 (6)	0.1283 (4)	0.0517 (19)	

supplementary materials

H1	-0.1978	0.5341	0.1403	0.062*	
C2	-0.0911 (6)	0.6243 (6)	0.1031 (4)	0.058 (2)	
H2	-0.1256	0.6863	0.0999	0.070*	
C3	0.0015 (6)	0.6235 (6)	0.0831 (4)	0.055 (2)	
H3	0.0305	0.6846	0.0660	0.066*	
C4	0.0535 (5)	0.5285 (6)	0.0887 (4)	0.0438 (18)	
C5	0.1496 (5)	0.5174 (6)	0.0688 (4)	0.0484 (19)	
H5	0.1820	0.5741	0.0481	0.058*	
C6	0.1946 (5)	0.4255 (6)	0.0796 (4)	0.050 (2)	
H6	0.2581	0.4210	0.0672	0.061*	
C7	0.1485 (5)	0.3347 (5)	0.1093 (4)	0.0403 (17)	
C8	0.1899 (5)	0.2382 (6)	0.1246 (4)	0.052 (2)	
H8	0.2534	0.2282	0.1141	0.062*	
C9	0.1393 (5)	0.1590 (6)	0.1546 (4)	0.0494 (19)	
H9	0.1678	0.0947	0.1647	0.059*	
C10	0.0446 (5)	0.1735 (5)	0.1703 (4)	0.0447 (18)	
H10	0.0098	0.1177	0.1897	0.054*	
C11	0.0538 (5)	0.3460 (6)	0.1273 (4)	0.0384 (17)	
C12	0.0055 (5)	0.4415 (5)	0.1167 (3)	0.0368 (16)	
I1	0.56522 (7)	0.45323 (7)	0.04476 (5)	0.0912 (4)	0.936 (3)
Cl5	0.558 (3)	0.496 (4)	0.021 (2)	0.062 (9)*	0.064 (3)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Pt1	0.04138 (16)	0.03167 (16)	0.04668 (17)	-0.00138 (13)	0.00129 (14)	0.00057 (14)
Cl1	0.0561 (11)	0.0392 (11)	0.0608 (12)	-0.0060 (9)	0.0047 (9)	0.0072 (9)
Cl2	0.0490 (11)	0.0431 (12)	0.0616 (12)	0.0050 (9)	0.0083 (9)	0.0040 (9)
Cl3	0.064 (4)	0.045 (3)	0.039 (4)	-0.005 (2)	-0.007 (2)	-0.014 (3)
I3	0.064 (4)	0.045 (3)	0.039 (4)	-0.005 (2)	-0.007 (2)	-0.014 (3)
Cl4	0.0649 (15)	0.052 (2)	0.038 (3)	-0.0073 (14)	-0.006 (3)	-0.004 (2)
I4	0.0649 (15)	0.052 (2)	0.038 (3)	-0.0073 (14)	-0.006 (3)	-0.004 (2)
N1	0.044 (4)	0.029 (4)	0.045 (3)	-0.001 (3)	-0.001 (3)	-0.002 (3)
N2	0.037 (3)	0.032 (3)	0.046 (3)	0.000 (3)	-0.005 (3)	-0.001 (3)
C1	0.053 (5)	0.048 (5)	0.055 (5)	0.001 (4)	-0.002 (4)	-0.007 (4)
C2	0.057 (5)	0.031 (5)	0.086 (6)	0.006 (4)	-0.009 (5)	-0.004 (4)
C3	0.071 (6)	0.034 (5)	0.060 (5)	-0.011 (4)	-0.003 (4)	0.003 (4)
C4	0.052 (5)	0.036 (5)	0.043 (4)	-0.006 (4)	-0.007 (4)	-0.003 (3)
C5	0.050 (5)	0.044 (5)	0.051 (5)	-0.017 (4)	-0.006 (4)	0.007 (4)
C6	0.045 (5)	0.052 (6)	0.054 (5)	-0.012 (4)	0.006 (4)	-0.011 (4)
C7	0.044 (5)	0.036 (4)	0.041 (4)	-0.001 (3)	-0.002 (3)	-0.006 (3)
C8	0.046 (5)	0.053 (5)	0.056 (5)	0.007 (4)	0.006 (4)	-0.004 (4)
C9	0.047 (5)	0.046 (5)	0.055 (5)	0.009 (4)	-0.007 (4)	-0.010 (4)
C10	0.050 (5)	0.032 (5)	0.052 (5)	-0.006 (3)	0.002 (4)	-0.002 (3)
C11	0.040 (4)	0.037 (4)	0.038 (4)	-0.007 (3)	-0.008 (3)	-0.001 (3)
C12	0.042 (4)	0.034 (4)	0.034 (4)	-0.002 (3)	-0.004 (3)	0.002 (3)
I1	0.1295 (7)	0.0635 (6)	0.0805 (6)	-0.0192 (5)	0.0312 (5)	-0.0058 (4)

Geometric parameters (Å, °)

Pt1—N2	2.040 (5)	C4—C12	1.391 (9)
Pt1—N1	2.048 (5)	C4—C5	1.417 (9)
Pt1—Cl3	2.272 (9)	C5—C6	1.346 (9)
Pt1—Cl1	2.2977 (19)	C5—H5	0.9300
Pt1—Cl4	2.301 (4)	C6—C7	1.428 (9)
Pt1—Cl2	2.3347 (19)	C6—H6	0.9300
Pt1—I3	2.598 (7)	C7—C8	1.389 (9)
Pt1—I4	2.635 (9)	C7—C11	1.391 (8)
N1—C1	1.309 (8)	C8—C9	1.346 (10)
N1—C12	1.364 (8)	C8—H8	0.9300
N2—C10	1.316 (8)	C9—C10	1.387 (9)
N2—C11	1.381 (8)	C9—H9	0.9300
C1—C2	1.388 (10)	C10—H10	0.9300
C1—H1	0.9300	C11—C12	1.408 (9)
C2—C3	1.364 (10)	I1—Cl5 ⁱ	2.19 (4)
C2—H2	0.9300	I1—I1 ⁱ	2.708 (2)
C3—C4	1.421 (10)	Cl5—I1 ⁱ	2.19 (4)
C3—H3	0.9300		
N2—Pt1—N1	81.3 (2)	N1—C1—H1	119.0
N2—Pt1—Cl3	88.1 (6)	C2—C1—H1	119.0
N1—Pt1—Cl3	87.8 (6)	C3—C2—C1	120.0 (7)
N2—Pt1—Cl1	93.11 (17)	C3—C2—H2	120.0
N1—Pt1—Cl1	174.14 (16)	C1—C2—H2	120.0
Cl3—Pt1—Cl1	90.3 (6)	C2—C3—C4	119.4 (7)
N2—Pt1—Cl4	90.0 (3)	C2—C3—H3	120.3
N1—Pt1—Cl4	91.0 (3)	C4—C3—H3	120.3
Cl3—Pt1—Cl4	177.9 (6)	C12—C4—C5	118.7 (7)
Cl1—Pt1—Cl4	90.8 (3)	C12—C4—C3	116.6 (7)
N2—Pt1—Cl2	175.97 (17)	C5—C4—C3	124.7 (7)
N1—Pt1—Cl2	94.69 (16)	C6—C5—C4	120.7 (7)
Cl3—Pt1—Cl2	92.1 (5)	C6—C5—H5	119.7
Cl1—Pt1—Cl2	90.91 (7)	C4—C5—H5	119.7
Cl4—Pt1—Cl2	89.7 (3)	C5—C6—C7	122.5 (7)
N2—Pt1—I3	85.6 (4)	C5—C6—H6	118.8
N1—Pt1—I3	86.5 (4)	C7—C6—H6	118.8
Cl3—Pt1—I3	2.7 (9)	C8—C7—C11	117.2 (7)
Cl1—Pt1—I3	91.4 (4)	C8—C7—C6	126.4 (7)
Cl4—Pt1—I3	175.2 (4)	C11—C7—C6	116.3 (6)
Cl2—Pt1—I3	94.6 (4)	C9—C8—C7	120.8 (7)
N2—Pt1—I4	89.4 (8)	C9—C8—H8	119.6
N1—Pt1—I4	91.3 (7)	C7—C8—H8	119.6
Cl3—Pt1—I4	177.5 (9)	C8—C9—C10	119.8 (7)
Cl1—Pt1—I4	90.4 (7)	C8—C9—H9	120.1
Cl4—Pt1—I4	0.7 (10)	C10—C9—H9	120.1
Cl2—Pt1—I4	90.3 (7)	N2—C10—C9	121.5 (7)

supplementary materials

I3—Pt1—I4	174.8 (8)	N2—C10—H10	119.2
C1—N1—C12	119.5 (6)	C9—C10—H10	119.2
C1—N1—Pt1	128.4 (5)	N2—C11—C7	121.3 (6)
C12—N1—Pt1	112.1 (4)	N2—C11—C12	116.6 (6)
C10—N2—C11	119.3 (6)	C7—C11—C12	122.0 (6)
C10—N2—Pt1	128.5 (5)	N1—C12—C4	122.5 (6)
C11—N2—Pt1	112.2 (4)	N1—C12—C11	117.7 (6)
N1—C1—C2	121.9 (7)	C4—C12—C11	119.7 (6)
N2—Pt1—N1—C1	-176.8 (6)	C3—C4—C5—C6	176.6 (7)
Cl3—Pt1—N1—C1	-88.4 (8)	C4—C5—C6—C7	1.6 (11)
Cl4—Pt1—N1—C1	93.3 (6)	C5—C6—C7—C8	-177.6 (7)
Cl2—Pt1—N1—C1	3.6 (6)	C5—C6—C7—C11	-0.4 (10)
I3—Pt1—N1—C1	-90.8 (7)	C11—C7—C8—C9	1.1 (10)
I4—Pt1—N1—C1	93.9 (9)	C6—C7—C8—C9	178.3 (7)
N2—Pt1—N1—C12	0.2 (4)	C7—C8—C9—C10	0.1 (11)
Cl3—Pt1—N1—C12	88.7 (7)	C11—N2—C10—C9	2.1 (9)
Cl4—Pt1—N1—C12	-89.7 (5)	Pt1—N2—C10—C9	-179.7 (5)
Cl2—Pt1—N1—C12	-179.4 (4)	C8—C9—C10—N2	-1.7 (10)
I3—Pt1—N1—C12	86.3 (5)	C10—N2—C11—C7	-0.9 (9)
I4—Pt1—N1—C12	-89.0 (8)	Pt1—N2—C11—C7	-179.4 (5)
N1—Pt1—N2—C10	-179.6 (6)	C10—N2—C11—C12	-179.4 (6)
Cl3—Pt1—N2—C10	92.4 (8)	Pt1—N2—C11—C12	2.1 (7)
Cl1—Pt1—N2—C10	2.2 (5)	C8—C7—C11—N2	-0.7 (9)
Cl4—Pt1—N2—C10	-88.6 (6)	C6—C7—C11—N2	-178.2 (6)
I3—Pt1—N2—C10	93.4 (7)	C8—C7—C11—C12	177.7 (6)
I4—Pt1—N2—C10	-88.2 (9)	C6—C7—C11—C12	0.3 (9)
N1—Pt1—N2—C11	-1.3 (4)	C1—N1—C12—C4	-1.0 (9)
Cl3—Pt1—N2—C11	-89.3 (7)	Pt1—N1—C12—C4	-178.3 (5)
Cl1—Pt1—N2—C11	-179.5 (4)	C1—N1—C12—C11	178.2 (6)
Cl4—Pt1—N2—C11	89.7 (5)	Pt1—N1—C12—C11	0.9 (7)
I3—Pt1—N2—C11	-88.3 (6)	C5—C4—C12—N1	-178.4 (6)
I4—Pt1—N2—C11	90.1 (8)	C3—C4—C12—N1	2.3 (9)
C12—N1—C1—C2	-1.2 (10)	C5—C4—C12—C11	2.4 (9)
Pt1—N1—C1—C2	175.7 (5)	C3—C4—C12—C11	-176.9 (6)
N1—C1—C2—C3	1.8 (11)	N2—C11—C12—N1	-2.0 (9)
C1—C2—C3—C4	-0.4 (11)	C7—C11—C12—N1	179.5 (6)
C2—C3—C4—C12	-1.6 (10)	N2—C11—C12—C4	177.2 (6)
C2—C3—C4—C5	179.2 (7)	C7—C11—C12—C4	-1.3 (10)
C12—C4—C5—C6	-2.6 (10)		

Symmetry codes: (i) $-x+1, -y+1, -z$.

Hydrogen-bond geometry ($\text{\AA}, ^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C1—H1 \cdots Cl2	0.93	2.73	3.320 (8)	122
C10—H10 \cdots Cl1	0.93	2.66	3.240 (7)	121

Fig. 1




Fig. 2

