

catena-Poly[[dimethylformamide- κ O)-copper(II)]-bis(μ -4-nitrophenyl-cyanamido- $\kappa^2 N^1:N^3$)]

Hossein Chiniforoshan,^{a*} Soghra Jalilpour,^a Bahare Shirinfar^a and Hamid Reza Khavasi^b

^aDepartment of Chemistry, Isfahan University of Technology, Isfahan 84156-38111, Iran, and ^bDepartment of Chemistry, Shahid Beheshti University, G.C., Evin, Tehran 1983963113, Iran

Correspondence e-mail: chinif@cc.iut.ac.ir


Received 9 January 2009; accepted 4 March 2009

Key indicators: single-crystal X-ray study; $T = 120$ K; mean $\sigma(C-C) = 0.002$ Å; R factor = 0.033; wR factor = 0.091; data-to-parameter ratio = 17.9.

In the title compound, $[Cu(C_7H_4N_3O_2)_2(C_3H_7NO)]$, the Cu^{II} atom is five-coordinated in a distorted square-pyramidal geometry, with the N atoms in equatorial positions and the dimethylformamide O atom in an axial position. The dihedral angle between adjacent benzene rings is 70.33 (12)°.


Related literature

The phenylcyanamide molecule can function as bridging ligand and can coordinate to two different metallic centers by means of the nitrile and amine N atoms ($\mu_{1,3}$ bonding mode), forming di- and polynuclear complexes, see: Ainscough *et al.* (1991); Brader *et al.* (1990); Crutchley (2001); Escuer *et al.* (2004). For the magnetic properties of coordination polymers, see: Grosshenny *et al.* (1996). For the preparation of 4-NO₂-phenylcyanamide used in the synthesis, see: Crutchley & Naklicki (1989).



Experimental

Crystal data



$M_r = 460.91$

Monoclinic, $P2_1/c$

$a = 21.5103$ (12) Å

$b = 8.7883$ (5) Å

$c = 9.9195$ (5) Å

$\beta = 101.746$ (4)°

$V = 1835.91$ (17) Å³

$Z = 4$

Mo $K\alpha$ radiation

$\mu = 1.24$ mm⁻¹

$T = 120$ K

$0.50 \times 0.23 \times 0.15$ mm

Data collection

Stoe IPDS-II diffractometer

Absorption correction: numerical
with shape of crystal determined
optically

$T_{min} = 0.720$, $T_{max} = 0.832$

13057 measured reflections

4874 independent reflections

4496 reflections with $I > 2\sigma(I)$

$R_{int} = 0.050$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.033$

$wR(F^2) = 0.091$

$S = 1.08$

4874 reflections

273 parameters

H-atom parameters constrained

$\Delta\rho_{\text{max}} = 0.99$ e Å⁻³

$\Delta\rho_{\text{min}} = -0.91$ e Å⁻³

Table 1
Selected geometric parameters (Å, °).

N2—Cu1	2.0862 (13)	Cu1—N1 ⁱⁱ	1.9748 (13)
N5—Cu1	2.0599 (12)	Cu1—O5	2.1443 (12)
Cu1—N4 ⁱ	1.9648 (13)		
N4 ⁱ —Cu1—N1 ⁱⁱ	154.42 (6)	N4 ⁱ —Cu1—O5	102.75 (5)
N4 ⁱ —Cu1—N5	90.85 (5)	N1 ⁱⁱ —Cu1—O5	102.83 (5)
N1 ⁱⁱ —Cu1—N2	90.41 (5)	N5—Cu1—O5	92.25 (5)
N5—Cu1—N2	172.66 (5)	N2—Cu1—O5	95.08 (5)

Symmetry codes: (i) $x, -y - \frac{3}{2}, z - \frac{1}{2}$; (ii) $x, -y - \frac{3}{2}, z + \frac{1}{2}$.

Data collection: *X-AREA* (Stoe & Cie, 2005); cell refinement: *X-AREA*; data reduction: *X-AREA*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

The authors acknowledge financial support from Isfahan University of Technology.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BQ2120).

References

- Ainscough, E. W., Baker, E. N., Brader, M. L. & Brodie, A. M. (1991). *J. Chem. Soc. Dalton Trans.* pp. 1243–1249.
- Brader, M. L., Ainscough, E. W., Baker, E. N. & Ingham, A. M. (1990). *J. Chem. Soc. Dalton Trans.* pp. 2785–2792.
- Crutchley, R. J. (2001). *Coord. Chem. Rev.* **219**, 125–155.
- Crutchley, R. J. & Naklicki, M. L. (1989). *Inorg. Chem.* **28**, 1955–1958.
- Escuer, A., Mautner, F. A., Sanz, N. & Vicente, R. (2004). *Polyhedron*, **23**, 1409–1417.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Grosshenny, V., Harriman, A., Romero, F. M. & Ziessel, R. (1996). *J. Chem. Phys.* **100**, 17472–17484.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Stoe & Cie (2005). *X-AREA*, *X-RED* and *X-SHAPE*. Stoe & Cie, Darmstadt, Germany.

supplementary materials

Acta Cryst. (2009). E65, m386 [doi:10.1107/S160053680900796X]

catena-Poly[[(dimethylformamide- κO)copper(II)]-bis(μ -4-nitrophenylcyanamido- $\kappa^2 N^1:N^3$)]

H. Chiniforoshan, S. Jalilpour, B. Shirinfar and H. R. Khavasi

Comment

Phenylcyanamide ligands are used in the construction of transition metal coordination complexes. The study of coordination polymeric materials holds great interest. The phenylcyanamide can function as bridging ligand and can coordinate to two different metallic centers by means of the nitrile and amine nitrogen ($\mu_{1,3}$ bonding mode), forming di- and polynuclear complexes (Brader *et al.*, 1990; Crutchley *et al.*, 2001; Ainscough *et al.*, 1991; Escuer *et al.*, 2004). It can modify the solubility and crystallinity of resulting compounds and there is the different coordination in complexes. We are attempting to construct conductive inorganic polymer chains that are cross-linked by cyanamide groups to a coordination complex. Coordination polymers also hold promise as novel materials because of their magnetic properties (Grosshenny *et al.*, 1996). More recently various aromatic cyanamide complexes have been studied by x-ray crystallography.

In the molecule of the title compound, (I), (Fig. 1) the selected bond lengths and angles are listed in Table 1. In this molecule, the $\{\text{Cu}(4-\text{NO}_2\text{-pcyd})_2(\text{DMF})\}_n$ one-dimensional chain coordination polymer bridged by 4-NO₂-phenylcyanamide. Each copper atom has a distorted square pyramidal geometry, that nitrogen atoms are in equatorial position and oxygen atom from DMF molecule is in axial position (Table 1.). The dihedral angle between adjacent phenyl rings in the polymeric chain is 70.33 (12)°.

Experimental

The 4-NO₂-phenylcyanamide (Crutchley *et al.*, 1989) (0.163 gr, 0.5 mmol) dissolved in methanol (30 ml) was added slowly to a solution of copper(II) acetate monohydrate (0.998 gr, 1 mmol) in methanol (30 ml). The mixture was stirred for 4 h. The solid filtered and crystals suitable for X-ray structure determination were obtained by dissolving in DMF then diffused by n-Hexane, after 1 week.

Refinement

All of the H atoms were positioned geometrically, with C—H = 0.93 Å for aromatic and aldehyde H atoms and with C—H = 0.93 Å for methyl H atoms, and constrained to ride on their parent atoms, with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$.

supplementary materials

Figures




Fig. 1. View of (I) with 30% probability displacement ellipsoids. Symmetry code (i): $x, -y - 3/2, z - 1/2$.

catena-Poly[[dimethylformamide- κO)copper(II)]- bis(μ -4-nitrophenylcyanamido- $\kappa^2 N^1:N^3$)]

Crystal data

[Cu(C ₇ H ₄ N ₃ O ₂) ₂ (C ₃ H ₇ NO)]	$F_{000} = 940$
$M_r = 460.91$	$D_x = 1.668 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
Hall symbol: -P 2ybc	$\lambda = 0.71073 \text{ \AA}$
$a = 21.5103 (12) \text{ \AA}$	Cell parameters from 1359 reflections
$b = 8.7883 (5) \text{ \AA}$	$\theta = 3.0\text{--}29.3^\circ$
$c = 9.9195 (5) \text{ \AA}$	$\mu = 1.24 \text{ mm}^{-1}$
$\beta = 101.746 (4)^\circ$	$T = 120 \text{ K}$
$V = 1835.91 (17) \text{ \AA}^3$	Prism, violet
$Z = 4$	$0.5 \times 0.23 \times 0.15 \text{ mm}$

Data collection

Stoe IPDS-II diffractometer	$R_{\text{int}} = 0.050$
rotation method scans	$\theta_{\text{max}} = 29.3^\circ$
Absorption correction: numerical shape of crystal determined optically	$\theta_{\text{min}} = 3.0^\circ$
$T_{\text{min}} = 0.720, T_{\text{max}} = 0.832$	$h = -29 \rightarrow 24$
13057 measured reflections	$k = -10 \rightarrow 12$
4874 independent reflections	$l = -13 \rightarrow 13$
4496 reflections with $I > 2\sigma(I)$	

Refinement

Refinement on F^2	H-atom parameters constrained
Least-squares matrix: full	$w = 1/[\sigma^2(F_o^2) + (0.051P)^2 + 0.8628P]$ where $P = (F_o^2 + 2F_c^2)/3$
$R[F^2 > 2\sigma(F^2)] = 0.033$	$(\Delta/\sigma)_{\text{max}} = 0.017$

$wR(F^2) = 0.091$ $\Delta\rho_{\max} = 0.99 \text{ e \AA}^{-3}$
 $S = 1.08$ $\Delta\rho_{\min} = -0.91 \text{ e \AA}^{-3}$
4874 reflections Extinction correction: none
273 parameters

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.18470 (7)	-0.78476 (17)	0.48486 (14)	0.0134 (3)
C2	0.11109 (7)	-0.85879 (17)	0.61210 (15)	0.0134 (3)
C3	0.06225 (7)	-0.77637 (19)	0.52815 (15)	0.0170 (3)
H3	0.0717	-0.7132	0.4599	0.02*
C4	-0.00005 (7)	-0.78755 (19)	0.54535 (16)	0.0187 (3)
H4	-0.0323	-0.7327	0.4893	0.022*
C5	-0.01325 (7)	-0.8821 (2)	0.64786 (16)	0.0179 (3)
C6	0.03397 (8)	-0.96827 (18)	0.73088 (16)	0.0177 (3)
H6	0.0239	-1.0325	0.7978	0.021*
C7	0.09602 (7)	-0.95722 (17)	0.71271 (15)	0.0155 (3)
H7	0.1278	-1.015	0.7671	0.019*
C8	0.31962 (7)	-0.74093 (18)	1.03158 (14)	0.0140 (3)
C9	0.39374 (7)	-0.82569 (17)	0.90993 (15)	0.0142 (3)
C10	0.41040 (7)	-0.94717 (18)	0.83254 (15)	0.0156 (3)
H10	0.3799	-1.0178	0.7932	0.019*
C11	0.47252 (7)	-0.96186 (19)	0.81480 (15)	0.0179 (3)
H11	0.4839	-1.0419	0.7634	0.021*
C12	0.51748 (8)	-0.8551 (2)	0.87497 (16)	0.0186 (3)
C13	0.50288 (7)	-0.7369 (2)	0.95631 (16)	0.0192 (3)
H13	0.534	-0.6688	0.9982	0.023*
C14	0.44059 (7)	-0.72297 (19)	0.97349 (15)	0.0170 (3)
H14	0.4299	-0.6448	1.0277	0.02*
C15	0.23643 (8)	-1.18081 (18)	0.70530 (16)	0.0164 (3)
H15	0.2162	-1.1469	0.6187	0.02*
C16	0.27640 (9)	-1.3914 (2)	0.85538 (18)	0.0239 (3)
H16A	0.2471	-1.4544	0.8914	0.029*
H16B	0.2907	-1.3101	0.9187	0.029*
H16C	0.3121	-1.4514	0.8431	0.029*
C17	0.22428 (9)	-1.4360 (2)	0.61151 (18)	0.0242 (3)
H17C	0.195	-1.5073	0.6375	0.029*
H17B	0.2605	-1.4896	0.5928	0.029*
H17A	0.2038	-1.382	0.5305	0.029*
N1	0.19773 (6)	-0.73209 (16)	0.38708 (13)	0.0159 (2)

supplementary materials

N2	0.17467 (6)	-0.84608 (14)	0.59891 (13)	0.0133 (2)
N3	-0.07813 (7)	-0.8905 (2)	0.67009 (16)	0.0245 (3)
N4	0.30602 (6)	-0.68183 (17)	1.12639 (13)	0.0169 (3)
N5	0.32972 (6)	-0.80866 (16)	0.92140 (12)	0.0136 (2)
N6	0.58231 (7)	-0.8682 (2)	0.85134 (15)	0.0249 (3)
N7	0.24485 (6)	-1.32833 (16)	0.72334 (14)	0.0166 (3)
Cu1	0.252435 (8)	-0.842098 (19)	0.763487 (17)	0.01030 (7)
O1	-0.11924 (6)	-0.8137 (2)	0.59671 (16)	0.0360 (3)
O2	-0.08874 (8)	-0.9731 (2)	0.7622 (2)	0.0497 (5)
O3	0.61904 (7)	-0.7620 (2)	0.88752 (16)	0.0380 (4)
O4	0.59667 (7)	-0.9839 (2)	0.79478 (16)	0.0379 (4)
O5	0.25389 (6)	-1.08324 (13)	0.79693 (11)	0.0182 (2)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0091 (6)	0.0152 (6)	0.0141 (6)	-0.0002 (5)	-0.0015 (5)	-0.0010 (5)
C2	0.0118 (6)	0.0143 (6)	0.0137 (6)	-0.0026 (5)	0.0017 (5)	-0.0023 (5)
C3	0.0150 (7)	0.0208 (7)	0.0146 (6)	0.0011 (5)	0.0017 (5)	0.0014 (5)
C4	0.0131 (7)	0.0228 (8)	0.0189 (6)	0.0024 (6)	0.0005 (5)	-0.0020 (6)
C5	0.0116 (6)	0.0208 (7)	0.0219 (7)	-0.0027 (6)	0.0047 (5)	-0.0059 (6)
C6	0.0162 (7)	0.0176 (7)	0.0204 (7)	-0.0046 (5)	0.0062 (5)	-0.0007 (5)
C7	0.0139 (6)	0.0158 (6)	0.0163 (6)	-0.0025 (5)	0.0019 (5)	0.0004 (5)
C8	0.0089 (6)	0.0183 (7)	0.0135 (6)	-0.0012 (5)	-0.0009 (5)	0.0010 (5)
C9	0.0115 (6)	0.0182 (7)	0.0125 (6)	0.0023 (5)	0.0015 (5)	0.0023 (5)
C10	0.0134 (6)	0.0177 (7)	0.0145 (6)	0.0020 (5)	0.0005 (5)	-0.0002 (5)
C11	0.0163 (7)	0.0224 (7)	0.0147 (6)	0.0060 (6)	0.0029 (5)	0.0012 (5)
C12	0.0116 (6)	0.0271 (8)	0.0173 (7)	0.0029 (6)	0.0035 (5)	0.0051 (6)
C13	0.0133 (6)	0.0246 (8)	0.0190 (7)	-0.0025 (6)	0.0019 (5)	0.0015 (6)
C14	0.0144 (7)	0.0204 (7)	0.0159 (6)	0.0002 (5)	0.0025 (5)	-0.0024 (5)
C15	0.0183 (7)	0.0142 (7)	0.0169 (6)	-0.0004 (5)	0.0039 (5)	0.0002 (5)
C16	0.0293 (8)	0.0163 (7)	0.0251 (8)	0.0047 (6)	0.0031 (6)	0.0053 (6)
C17	0.0303 (9)	0.0170 (7)	0.0265 (8)	-0.0018 (6)	0.0089 (7)	-0.0079 (6)
N1	0.0104 (5)	0.0230 (7)	0.0139 (5)	-0.0003 (5)	0.0014 (4)	0.0018 (5)
N2	0.0100 (5)	0.0181 (6)	0.0110 (5)	-0.0018 (4)	0.0006 (4)	0.0006 (4)
N3	0.0151 (6)	0.0283 (8)	0.0312 (7)	-0.0031 (6)	0.0074 (5)	-0.0066 (6)
N4	0.0105 (5)	0.0263 (7)	0.0130 (5)	-0.0018 (5)	-0.0001 (4)	-0.0031 (5)
N5	0.0093 (5)	0.0191 (6)	0.0119 (5)	0.0002 (4)	0.0007 (4)	-0.0023 (4)
N6	0.0147 (6)	0.0396 (9)	0.0211 (6)	0.0052 (6)	0.0049 (5)	0.0051 (6)
N7	0.0183 (6)	0.0130 (6)	0.0186 (6)	-0.0008 (4)	0.0039 (5)	-0.0012 (4)
Cu1	0.00955 (10)	0.01191 (10)	0.00912 (10)	-0.00041 (6)	0.00118 (6)	-0.00033 (5)
O1	0.0144 (6)	0.0544 (10)	0.0386 (8)	0.0052 (6)	0.0040 (5)	-0.0007 (7)
O2	0.0246 (7)	0.0595 (11)	0.0711 (12)	0.0005 (7)	0.0246 (8)	0.0261 (10)
O3	0.0161 (6)	0.0527 (10)	0.0463 (8)	-0.0059 (6)	0.0087 (6)	-0.0026 (7)
O4	0.0244 (7)	0.0487 (9)	0.0443 (8)	0.0094 (7)	0.0154 (6)	-0.0049 (7)
O5	0.0248 (6)	0.0119 (5)	0.0169 (5)	-0.0014 (4)	0.0017 (4)	-0.0005 (4)

Geometric parameters (Å, °)

C1—N1	1.160 (2)	C13—C14	1.390 (2)
C1—N2	1.3099 (18)	C13—H13	0.93
C2—C3	1.402 (2)	C14—H14	0.93
C2—N2	1.4046 (19)	C15—O5	1.250 (2)
C2—C7	1.408 (2)	C15—N7	1.316 (2)
C3—C4	1.388 (2)	C15—H15	0.93
C3—H3	0.93	C16—N7	1.457 (2)
C4—C5	1.387 (2)	C16—H16A	0.96
C4—H4	0.93	C16—H16B	0.96
C5—C6	1.394 (2)	C16—H16C	0.96
C5—N3	1.458 (2)	C17—N7	1.457 (2)
C6—C7	1.386 (2)	C17—H17C	0.96
C6—H6	0.93	C17—H17B	0.96
C7—H7	0.93	C17—H17A	0.96
C8—N4	1.163 (2)	N1—Cu1 ⁱ	1.9748 (13)
C8—N5	1.3010 (19)	N2—Cu1	2.0862 (13)
C9—C14	1.403 (2)	N3—O2	1.224 (2)
C9—C10	1.403 (2)	N3—O1	1.227 (2)
C9—N5	1.4126 (19)	N4—Cu1 ⁱⁱ	1.9648 (13)
C10—C11	1.389 (2)	N5—Cu1	2.0599 (12)
C10—H10	0.93	N6—O3	1.229 (2)
C11—C12	1.392 (2)	N6—O4	1.230 (2)
C11—H11	0.93	Cu1—N4 ⁱ	1.9648 (13)
C12—C13	1.390 (2)	Cu1—N1 ⁱⁱ	1.9748 (13)
C12—N6	1.465 (2)	Cu1—O5	2.1443 (12)
N1—C1—N2	175.56 (15)	N7—C16—H16A	109.5
C3—C2—N2	121.98 (13)	N7—C16—H16B	109.5
C3—C2—C7	119.06 (14)	H16A—C16—H16B	109.5
N2—C2—C7	118.96 (13)	N7—C16—H16C	109.5
C4—C3—C2	121.05 (14)	H16A—C16—H16C	109.5
C4—C3—H3	119.5	H16B—C16—H16C	109.5
C2—C3—H3	119.5	N7—C17—H17C	109.5
C5—C4—C3	118.64 (14)	N7—C17—H17B	109.5
C5—C4—H4	120.7	H17C—C17—H17B	109.5
C3—C4—H4	120.7	N7—C17—H17A	109.5
C4—C5—C6	121.70 (14)	H17C—C17—H17A	109.5
C4—C5—N3	119.07 (15)	H17B—C17—H17A	109.5
C6—C5—N3	119.22 (15)	C1—N1—Cu1 ⁱ	158.01 (12)
C7—C6—C5	119.36 (14)	C1—N2—C2	116.60 (12)
C7—C6—H6	120.3	C1—N2—Cu1	114.97 (10)
C5—C6—H6	120.3	C2—N2—Cu1	124.69 (9)
C6—C7—C2	120.15 (14)	O2—N3—O1	123.22 (16)
C6—C7—H7	119.9	O2—N3—C5	118.16 (16)
C2—C7—H7	119.9	O1—N3—C5	118.63 (16)
N4—C8—N5	175.15 (15)	C8—N4—Cu1 ⁱⁱ	152.54 (13)

supplementary materials

C14—C9—C10	119.68 (14)	C8—N5—C9	116.64 (12)
C14—C9—N5	121.22 (13)	C8—N5—Cu1	117.20 (10)
C10—C9—N5	119.09 (14)	C9—N5—Cu1	124.94 (9)
C11—C10—C9	119.96 (15)	O3—N6—O4	123.70 (16)
C11—C10—H10	120	O3—N6—C12	118.03 (16)
C9—C10—H10	120	O4—N6—C12	118.26 (16)
C10—C11—C12	119.05 (14)	C15—N7—C17	121.62 (15)
C10—C11—H11	120.5	C15—N7—C16	121.55 (14)
C12—C11—H11	120.5	C17—N7—C16	116.82 (14)
C13—C12—C11	122.23 (14)	N4 ⁱ —Cu1—N1 ⁱⁱ	154.42 (6)
C13—C12—N6	119.11 (15)	N4 ⁱ —Cu1—N5	90.85 (5)
C11—C12—N6	118.65 (15)	N1 ⁱⁱ —Cu1—N5	88.36 (5)
C14—C13—C12	118.31 (15)	N4 ⁱ —Cu1—N2	87.13 (5)
C14—C13—H13	120.8	N1 ⁱⁱ —Cu1—N2	90.41 (5)
C12—C13—H13	120.8	N5—Cu1—N2	172.66 (5)
C13—C14—C9	120.69 (14)	N4 ⁱ —Cu1—O5	102.75 (5)
C13—C14—H14	119.7	N1 ⁱⁱ —Cu1—O5	102.83 (5)
C9—C14—H14	119.7	N5—Cu1—O5	92.25 (5)
O5—C15—N7	124.33 (15)	N2—Cu1—O5	95.08 (5)
O5—C15—H15	117.8	C15—O5—Cu1	124.92 (11)
N7—C15—H15	117.8		
N2—C2—C3—C4	−178.27 (14)	C14—C9—N5—C8	−27.7 (2)
C7—C2—C3—C4	1.7 (2)	C10—C9—N5—C8	152.76 (14)
C2—C3—C4—C5	0.1 (2)	C14—C9—N5—Cu1	139.25 (12)
C3—C4—C5—C6	−1.6 (2)	C10—C9—N5—Cu1	−40.26 (19)
C3—C4—C5—N3	177.78 (15)	C13—C12—N6—O3	−10.4 (2)
C4—C5—C6—C7	1.3 (2)	C11—C12—N6—O3	169.52 (16)
N3—C5—C6—C7	−178.12 (14)	C13—C12—N6—O4	170.58 (16)
C5—C6—C7—C2	0.6 (2)	C11—C12—N6—O4	−9.5 (2)
C3—C2—C7—C6	−2.0 (2)	O5—C15—N7—C17	−178.97 (15)
N2—C2—C7—C6	177.94 (14)	O5—C15—N7—C16	−0.3 (3)
C14—C9—C10—C11	−2.5 (2)	C8—N5—Cu1—N4 ⁱ	147.21 (12)
N5—C9—C10—C11	177.06 (13)	C9—N5—Cu1—N4 ⁱ	−19.70 (13)
C9—C10—C11—C12	0.2 (2)	C8—N5—Cu1—N1 ⁱⁱ	−7.22 (12)
C10—C11—C12—C13	2.2 (2)	C9—N5—Cu1—N1 ⁱⁱ	−174.13 (13)
C10—C11—C12—N6	−177.70 (14)	C8—N5—Cu1—O5	−110.00 (12)
C11—C12—C13—C14	−2.2 (2)	C9—N5—Cu1—O5	83.09 (13)
N6—C12—C13—C14	177.69 (14)	C1—N2—Cu1—N4 ⁱ	−21.30 (11)
C12—C13—C14—C9	−0.1 (2)	C2—N2—Cu1—N4 ⁱ	−178.68 (12)
C10—C9—C14—C13	2.5 (2)	C1—N2—Cu1—N1 ⁱⁱ	133.23 (11)
N5—C9—C14—C13	−177.06 (14)	C2—N2—Cu1—N1 ⁱⁱ	−24.15 (12)
C3—C2—N2—C1	−16.4 (2)	C1—N2—Cu1—O5	−123.85 (11)
C7—C2—N2—C1	163.60 (14)	C2—N2—Cu1—O5	78.76 (12)
C3—C2—N2—Cu1	140.62 (12)	N7—C15—O5—Cu1	171.10 (11)
C7—C2—N2—Cu1	−39.34 (18)	N4 ⁱ —Cu1—O5—C15	−57.95 (14)

supplementary materials

C4—C5—N3—O2	−179.12 (18)	N1 ⁱⁱ —Cu1—O5—C15	121.84 (13)
C6—C5—N3—O2	0.3 (3)	N5—Cu1—O5—C15	−149.33 (13)
C4—C5—N3—O1	0.3 (2)	N2—Cu1—O5—C15	30.25 (13)
C6—C5—N3—O1	179.73 (16)		
Symmetry codes: (i) $x, -y-3/2, z-1/2$; (ii) $x, -y-3/2, z+1/2$.			

supplementary materials

Fig. 1

