

Diaquabis(*N,N*-diethylnicotinamide- κN^1)bis(4-formylbenzoato- κO)cobalt(II)

Mustafa Sertçelik,^a Barış Tercan,^b Ertan Şahin,^c Hacalı Necefoğlu^a and Tuncer Hökelek^{d*}

^aKafkas University, Department of Chemistry, 63100 Kars, Turkey, ^bKarabük University, Department of Physics, 78050 Karabük, Turkey, ^cAtatürk University, Department of Chemistry, 22240 Erzurum, Turkey, and ^dHacettepe University, Department of Physics, 06800 Beytepe, Ankara, Turkey

Correspondence e-mail: merzifon@hacettepe.edu.tr


Received 25 February 2009; accepted 6 March 2009

Key indicators: single-crystal X-ray study; $T = 294$ K; mean $\sigma(C-C) = 0.008$ Å; R factor = 0.069; wR factor = 0.195; data-to-parameter ratio = 15.3.

In the crystal structure of the title Co^{II} complex, [Co(C₈H₅O₃)₂(C₁₀H₁₄N₂O)₂(H₂O)₂], the metal centre is located on an inversion center and is coordinated by two 4-formylbenzoate (FOB), two diethylnicotinamide (DENA) ligands and two water molecules in a slightly distorted CoO₄N₂ octahedral geometry. In the crystal structure, O—H···O hydrogen bonds link the molecules into infinite chains. $\pi-\pi$ contacts between the parallel pyridine rings of neighboring DENA ligands [centroid–centroid distance = 3.652 (3) Å] further stabilize the crystal structure.

Related literature

For general background, see: Antolini *et al.* (1982); Bigoli *et al.* (1972); Nadzhafov *et al.* (1981); Shnulin *et al.* (1981). For related structures, see: Hökelek *et al.* (1995, 1997, 2007, 2008); Hökelek & Necefoğlu (1996, 1997, 2007); Sertçelik *et al.* (2009a, 2009b).

**Experimental***Crystal data*

[Co(C ₈ H ₅ O ₃) ₂ (C ₁₀ H ₁₄ N ₂ O) ₂ (H ₂ O) ₂]	$\beta = 78.608$ (3)°
$M_r = 749.67$	$\gamma = 68.022$ (2)°
Triclinic, $P\bar{1}$	$V = 918.64$ (5) Å ³
$a = 7.2962$ (2) Å	$Z = 1$
$b = 8.6863$ (3) Å	Mo $K\alpha$ radiation
$c = 15.9453$ (5) Å	$\mu = 0.53$ mm ⁻¹
$\alpha = 85.433$ (2)°	$T = 294$ K
	0.35 × 0.25 × 0.15 mm

Data collection

Rigaku R-AXIS RAPID-S diffractometer	19487 measured reflections
Absorption correction: multi-scan (Blessing, 1995)	3755 independent reflections
$T_{\min} = 0.853$, $T_{\max} = 0.926$	3016 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.074$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.069$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.195$	$\Delta\rho_{\max} = 1.02$ e Å ⁻³
$S = 1.07$	$\Delta\rho_{\min} = -0.33$ e Å ⁻³
3755 reflections	
246 parameters	
3 restraints	

Table 1
Selected bond lengths (Å).

Co1—O1	2.088 (3)	Co1—N1	2.163 (3)
Co1—O5	2.121 (3)		

Table 2
Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
O5—H51···O4 ⁱ	0.94 (4)	1.86 (4)	2.787 (5)	174 (6)
O5—H52···O2 ⁱⁱ	0.92 (2)	1.73 (4)	2.646 (5)	168 (6)

Symmetry codes: (i) $-x, -y, -z$; (ii) $-x + 1, -y, -z$.

Data collection: *CrystalClear* (Rigaku/MSC, 2005); cell refinement: *CrystalClear*; data reduction: *CrystalClear*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *Mercury* (Macrae *et al.*, 2006); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

The authors are indebted to the Department of Chemistry, Atatürk University, Erzurum, Turkey, for the use of the X-ray diffractometer purchased under grant No. 2003/219 of the University Research Fund.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: XU2489).

References

- Antolini, L., Battaglia, L. P., Corradi, A. B., Marcotrigiano, G., Menabue, L., Pellacani, G. C. & Saladini, M. (1982). *Inorg. Chem.* **21**, 1391–1395.
- Bigoli, F., Braibanti, A., Pellinghelli, M. A. & Tiripicchio, A. (1972). *Acta Cryst. B* **28**, 962–966.
- Blessing, R. H. (1995). *Acta Cryst. A* **51**, 33–38.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Hökelek, T., Budak, K. & Necefoğlu, H. (1997). *Acta Cryst. C* **53**, 1049–1051.
- Hökelek, T., Çaylak, N. & Necefoğlu, H. (2007). *Acta Cryst. E* **63**, m2561–m2562.
- Hökelek, T., Çaylak, N. & Necefoğlu, H. (2008). *Acta Cryst. E* **64**, m505–m506.
- Hökelek, T. & Necefoğlu, H. (1996). *Acta Cryst. C* **52**, 1128–1131.
- Hökelek, T. & Necefoğlu, H. (1997). *Acta Cryst. C* **53**, 187–189.
- Hökelek, T. & Necefoğlu, H. (2007). *Acta Cryst. E* **63**, m821–m823.
- Hökelek, T., Necefoğlu, H. & Balci, M. (1995). *Acta Cryst. C* **51**, 2020–2023.
- Macrae, C. F., Edgington, P. R., McCabe, P., Pidcock, E., Shields, G. P., Taylor, R., Towler, M. & van de Streek, J. (2006). *J. Appl. Cryst.* **39**, 453–457.
- Nadzhafov, G. N., Shnulin, A. N. & Mamedov, Kh. S. (1981). *Zh. Strukt. Khim.* **22**, 124–128.
- Rigaku/MSC (2005). *CrystalClear*. Rigaku/MSC, The Woodlands, Texas, USA.
- Sertçelik, M., Tercan, B., Şahin, E., Necefoğlu, H. & Hökelek, T. (2009a). *Acta Cryst. E* **65**, m326–m327.
- Sertçelik, M., Tercan, B., Şahin, E., Necefoğlu, H. & Hökelek, T. (2009b). *Acta Cryst. E* **65**, m324–m325.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Shnulin, A. N., Nadzhafov, G. N., Amiraslanov, I. R., Usibaliev, B. T. & Mamedov, Kh. S. (1981). *Koord. Khim.* **7**, 1409–1416.

supplementary materials

Diaqua bis(*N,N*-diethylnicotinamide- κN^1)bis(4-formylbenzoato- κO)cobalt(II)**M. Sertçelik, B. Tercan, E. Sahin, H. Necefoglu and T. Hökelek****Comment**

Transition metal complexes with biochemically active ligands frequently show interesting physical and/or chemical properties, as a result they may find applications in biological systems (Antolini *et al.*, 1982). The structural functions and co-ordination relationships of the arylcarboxylate ion in transition metal complexes of benzoic acid derivatives change depending on the nature and position of the substituent groups on the benzene ring, the nature of the additional ligand molecule or solvent, and the medium of the synthesis (Nadzhafov *et al.*, 1981; Shnulin *et al.*, 1981). The nicotinic acid derivative *N,N*-diethylnicotinamide (DENA) is an important respiratory stimulant (Bigoli *et al.*, 1972).

The structure determination of the title compound, (I), a cobalt complex with two formylbenzoate (FOB), two diethyl-nicotinamide (DENA) ligands and two water molecules, was undertaken in order to determine the properties of the ligands and also to compare the results obtained with those reported previously.

Compound (I) is a monomeric complex, with the Co atom on a centre of symmetry. It contains two FOB, two DENA ligands and two water molecules (Fig. 1). All ligands are monodentate. The four O atoms (O1, O5, and the symmetry-related atoms, O1', O5') in the equatorial plane around the Co atom form a slightly distorted square-planar arrangement, while the slightly distorted octahedral coordination is completed by the two N atoms of the DENA ligands (N1, N1') in the axial positions (Table 1 and Fig. 1).

The near equality of the C1—O1 [1.262 (5) Å] and C1—O2 [1.257 (5) Å] bonds in the carboxylate group indicates a delocalized bonding arrangement, rather than localized single and double bonds, and may be compared with the corresponding distances: 1.263 (4) and 1.249 (4) Å in [Ni(DENA)₂(C₈H₅O₃)₂(H₂O)₂], (II), (Sertçelik *et al.*, 2009a), 1.262 (3) and 1.249 (3) Å in [Mn(DENA)₂(C₈H₅O₃)₂(H₂O)₂], (III), (Sertçelik *et al.*, 2009b), 1.256 (6) and 1.245 (6) Å in [Mn(DENA)₂(C₇H₄ClO₂)₂(H₂O)₂], (IV), (Hökelek *et al.*, 2008), 1.265 (6) and 1.275 (6) Å in [Mn(C₉H₁₀NO₂)₂(H₂O)₄]. 2(H₂O), (V), Hökelek & Necefoglu, 2007), 1.260 (4) and 1.252 (4) Å in [Zn(DENA)₂(C₇H₄FO₂)₂(H₂O)₂], (VI), (Hökelek *et al.*, 2007), 1.259 (9) and 1.273 (9) Å in Cu₂(DENA)₂(C₆H₅COO)₄, (VII), (Hökelek *et al.*, 1995), 1.279 (4) and 1.246 (4) Å in [Zn₂(DENA)₂(C₇H₅O₃)₄]. 2H₂O, (VIII), (Hökelek & Necefoglu, 1996), 1.251 (6) and 1.254 (7) Å in [Co(DENA)₂(C₇H₅O₃)₂(H₂O)₂], (IX), (Hökelek & Necefoglu, 1997), 1.278 (3) and 1.246 (3) Å in [Cu(DENA)₂(C₇H₄NO₄)₂(H₂O)₂], (X), (Hökelek *et al.*, 1997). This may be due to the intermolecular O—H···O hydrogen bonding of the carboxylate O atoms (Table 2). In (I), the average Co—O bond length is 2.105 (3) Å and the Co atom is displaced out of the least-squares plane of the carboxylate group (O1/C1/O2) by 0.768 (1) Å. The dihedral angle between the planar carboxylate group and the benzene ring A (C2—C7) is 4.02 (35)°, while that between rings A and B (N1/C9—C13) is 79.61 (14)°.

In the crystal structure, intermolecular O—H···O hydrogen bonds (Table 1) link the molecules into infinite chains (Fig. 2), in which they may be effective in the stabilization of the structure. The π — π contact between the DENA rings, Cg1—Cg1ⁱ

supplementary materials

[symmetry code: (i) $2 - x, 1 - y, -z$, where $Cg1$ is centroid of the ring B (N1/C9—C13)] may further stabilize the structure, with centroid-centroid distance of 3.652 (3) Å.

Experimental

The title compound was prepared by the reaction of $\text{CoSO}_4 \cdot \text{H}_2\text{O}$ (1.73 g, 10 mmol) in H_2O (50 ml) and DENA (3.56 g, 20 mmol) in H_2O (15 ml) with sodium 4-formylbenzoate (3.44 g, 20 mmol) in H_2O (50 ml). The mixture was filtered and set aside to crystallize at ambient temperature for several days, giving red single crystals.

Refinement

H atoms of water molecule and formyl group were located in difference Fourier maps and refined isotropically, with restraints of $O5-H51 = 0.94$ (4), $O5-H52 = 0.92$ (2) Å, $H51-O5-H52 = 105$ (3)° and $U_{\text{iso}}(\text{H}) = 0.09$ (2) and 0.075 (18) Å² (for H_2O); $C8-H81 = 1.04$ (6) Å and $U_{\text{iso}}(\text{H}) = 0.086$ (19) Å² (for formyl group). The remaining H atoms were positioned geometrically with C—H = 0.93, 0.97 and 0.96 Å, for aromatic, methylene and methyl H atoms and constrained to ride on their parent atoms, with $U_{\text{iso}}(\text{H}) = xU_{\text{eq}}(\text{C})$, where $x = 1.5$ for methyl H atoms, and $x = 1.2$ for all other H atoms.

Figures




Fig. 1. The molecular structure of the title molecule with the atom-numbering scheme. Displacement ellipsoids are drawn at the 50% probability level. Hydrogen bonds are shown as dashed lines. Primed atoms are generated by the symmetry operator ($1 -x, -y, -z$).



Fig. 2. A partial packing diagram of (I) viewed down the a axis, showing hydrogen bonds (dotted lines) linking the molecules into chains, where b and c axes are vertical and horizontal, respectively. H atoms not involved in hydrogen bonding are omitted.

Diaquabis(*N,N*-diethylnicotinamide- κN^1)bis(4- formylbenzoato- κO)cobalt(II)

Crystal data

$[\text{Co}(\text{C}_8\text{H}_5\text{O}_3)_2(\text{C}_{10}\text{H}_{14}\text{N}_2\text{O})_2(\text{H}_2\text{O})_2]$	$Z = 1$
$M_r = 749.67$	$F_{000} = 393$
Triclinic, $P\bar{1}$	$D_x = 1.355 \text{ Mg m}^{-3}$
Hall symbol: -P 1	Mo $K\alpha$ radiation
$a = 7.2962$ (2) Å	$\lambda = 0.71073$ Å
$b = 8.6863$ (3) Å	Cell parameters from 4172 reflections
$c = 15.9453$ (5) Å	$\theta = 2.5\text{--}26.4^\circ$
$\alpha = 85.433$ (2)°	$\mu = 0.53 \text{ mm}^{-1}$
$\beta = 78.608$ (3)°	$T = 294$ K
$\gamma = 68.022$ (2)°	Prism, red
	$0.35 \times 0.25 \times 0.15$ mm

$V = 918.64 (5) \text{ \AA}^3$

Data collection

Rigaku R-AXIS RAPID-S diffractometer	3755 independent reflections
Radiation source: fine-focus sealed tube	3016 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.074$
$T = 294 \text{ K}$	$\theta_{\text{max}} = 26.4^\circ$
ω scans	$\theta_{\text{min}} = 2.5^\circ$
Absorption correction: multi-scan (Blessing, 1995)	$h = -9 \rightarrow 9$
$T_{\text{min}} = 0.853, T_{\text{max}} = 0.926$	$k = -10 \rightarrow 10$
19487 measured reflections	$l = -19 \rightarrow 19$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.069$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.195$	$w = 1/[\sigma^2(F_o^2) + (0.0999P)^2 + 0.5936P]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.07$	$(\Delta/\sigma)_{\text{max}} < 0.001$
3755 reflections	$\Delta\rho_{\text{max}} = 1.02 \text{ e \AA}^{-3}$
246 parameters	$\Delta\rho_{\text{min}} = -0.33 \text{ e \AA}^{-3}$
3 restraints	Extinction correction: none
Primary atom site location: structure-invariant direct methods	

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Co1	0.5000	0.0000	0.0000	0.0386 (3)
O1	0.4778 (5)	-0.1144 (4)	-0.10591 (18)	0.0473 (7)

supplementary materials

O2	0.7579 (5)	-0.1290 (4)	-0.1979 (2)	0.0555 (8)
O3	0.0578 (8)	-0.1907 (7)	-0.4559 (3)	0.0958 (15)
O4	-0.2309 (5)	0.3315 (4)	-0.1246 (2)	0.0585 (9)
O5	0.2256 (5)	-0.0208 (4)	0.0630 (2)	0.0506 (8)
H51	0.224 (10)	-0.126 (4)	0.080 (3)	0.09 (2)*
H52	0.214 (9)	0.033 (5)	0.113 (2)	0.075 (18)*
N1	0.3188 (5)	0.2363 (4)	-0.0485 (2)	0.0419 (8)
N2	-0.1116 (7)	0.4197 (6)	-0.2512 (3)	0.0609 (11)
C1	0.5813 (6)	-0.1268 (5)	-0.1802 (3)	0.0422 (9)
C2	0.4825 (6)	-0.1361 (5)	-0.2530 (3)	0.0418 (9)
C3	0.5824 (7)	-0.1374 (6)	-0.3370 (3)	0.0496 (11)
H3	0.7097	-0.1317	-0.3480	0.060*
C4	0.4938 (8)	-0.1471 (7)	-0.4037 (3)	0.0570 (12)
H4	0.5613	-0.1474	-0.4596	0.068*
C5	0.3035 (8)	-0.1566 (6)	-0.3882 (3)	0.0546 (11)
C6	0.2016 (7)	-0.1530 (6)	-0.3045 (3)	0.0510 (11)
H6	0.0741	-0.1583	-0.2934	0.061*
C7	0.2906 (7)	-0.1417 (6)	-0.2383 (3)	0.0463 (10)
H7	0.2213	-0.1377	-0.1824	0.056*
C8	0.2121 (10)	-0.1698 (9)	-0.4608 (4)	0.0750 (16)
H81	0.294 (9)	-0.169 (7)	-0.522 (4)	0.086 (19)*
C9	0.3415 (6)	0.3797 (5)	-0.0385 (3)	0.0439 (10)
H9	0.4366	0.3784	-0.0070	0.053*
C10	0.2310 (7)	0.5285 (6)	-0.0724 (3)	0.0494 (10)
H10	0.2502	0.6255	-0.0636	0.059*
C11	0.0905 (7)	0.5306 (5)	-0.1200 (3)	0.0478 (10)
H11	0.0157	0.6287	-0.1449	0.057*
C12	0.0637 (6)	0.3839 (5)	-0.1299 (3)	0.0426 (9)
C13	0.1798 (6)	0.2402 (5)	-0.0936 (3)	0.0419 (9)
H13	0.1613	0.1421	-0.1005	0.050*
C14	-0.1028 (7)	0.3743 (6)	-0.1700 (3)	0.0468 (10)
C15	0.0391 (10)	0.4689 (8)	-0.3107 (3)	0.0732 (16)
H15A	0.1301	0.4851	-0.2785	0.088*
H15B	-0.0288	0.5745	-0.3370	0.088*
C16	0.1577 (13)	0.3470 (13)	-0.3789 (6)	0.136 (4)
H16A	0.2601	0.3820	-0.4123	0.203*
H16B	0.0709	0.3392	-0.4151	0.203*
H16C	0.2193	0.2404	-0.3536	0.203*
C17	-0.2911 (10)	0.4212 (8)	-0.2844 (4)	0.0754 (16)
H17A	-0.4114	0.4706	-0.2420	0.090*
H17B	-0.3049	0.4888	-0.3359	0.090*
C18	-0.2719 (12)	0.2533 (8)	-0.3035 (5)	0.097 (2)
H18A	-0.3828	0.2589	-0.3290	0.146*
H18B	-0.2721	0.1894	-0.2515	0.146*
H18C	-0.1483	0.2014	-0.3424	0.146*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Co1	0.0376 (5)	0.0409 (5)	0.0385 (5)	-0.0145 (3)	-0.0104 (3)	0.0022 (3)
O1	0.0497 (17)	0.0557 (18)	0.0396 (16)	-0.0214 (15)	-0.0118 (13)	0.0004 (13)
O2	0.0445 (18)	0.069 (2)	0.0525 (18)	-0.0191 (16)	-0.0098 (14)	-0.0061 (16)
O3	0.100 (3)	0.148 (5)	0.068 (3)	-0.070 (3)	-0.032 (2)	0.002 (3)
O4	0.0547 (19)	0.070 (2)	0.062 (2)	-0.0353 (18)	-0.0136 (16)	0.0081 (17)
O5	0.0472 (18)	0.059 (2)	0.0508 (18)	-0.0255 (16)	-0.0102 (14)	0.0031 (15)
N1	0.0381 (18)	0.0404 (19)	0.049 (2)	-0.0148 (15)	-0.0127 (15)	0.0019 (15)
N2	0.065 (3)	0.077 (3)	0.054 (2)	-0.036 (2)	-0.023 (2)	0.010 (2)
C1	0.043 (2)	0.038 (2)	0.044 (2)	-0.0119 (18)	-0.0106 (18)	-0.0002 (17)
C2	0.044 (2)	0.039 (2)	0.041 (2)	-0.0131 (18)	-0.0086 (17)	0.0002 (17)
C3	0.044 (2)	0.055 (3)	0.049 (2)	-0.018 (2)	-0.0040 (19)	-0.001 (2)
C4	0.065 (3)	0.071 (3)	0.037 (2)	-0.028 (3)	-0.006 (2)	-0.001 (2)
C5	0.061 (3)	0.062 (3)	0.045 (2)	-0.026 (2)	-0.014 (2)	0.000 (2)
C6	0.049 (3)	0.061 (3)	0.048 (3)	-0.025 (2)	-0.012 (2)	0.001 (2)
C7	0.047 (2)	0.050 (2)	0.041 (2)	-0.020 (2)	-0.0047 (18)	0.0025 (18)
C8	0.083 (4)	0.106 (5)	0.050 (3)	-0.046 (4)	-0.017 (3)	-0.003 (3)
C9	0.041 (2)	0.047 (2)	0.048 (2)	-0.0174 (19)	-0.0145 (18)	0.0010 (18)
C10	0.054 (3)	0.044 (2)	0.056 (3)	-0.023 (2)	-0.013 (2)	0.003 (2)
C11	0.049 (2)	0.041 (2)	0.051 (3)	-0.016 (2)	-0.011 (2)	0.0079 (19)
C12	0.041 (2)	0.049 (2)	0.038 (2)	-0.0166 (19)	-0.0089 (17)	0.0038 (18)
C13	0.042 (2)	0.041 (2)	0.044 (2)	-0.0156 (18)	-0.0107 (17)	0.0020 (17)
C14	0.044 (2)	0.049 (3)	0.049 (2)	-0.018 (2)	-0.0126 (19)	0.0054 (19)
C15	0.087 (4)	0.084 (4)	0.057 (3)	-0.040 (3)	-0.015 (3)	0.005 (3)
C16	0.109 (6)	0.178 (9)	0.122 (7)	-0.066 (7)	0.029 (5)	-0.069 (7)
C17	0.082 (4)	0.075 (4)	0.080 (4)	-0.032 (3)	-0.040 (3)	0.017 (3)
C18	0.134 (7)	0.086 (5)	0.088 (5)	-0.046 (5)	-0.049 (5)	-0.001 (4)

Geometric parameters (\AA , $^\circ$)

Co1—O1	2.088 (3)	C7—C6	1.370 (6)
Co1—O1 ⁱ	2.088 (3)	C7—H7	0.9300
Co1—O5	2.121 (3)	C8—H81	1.04 (6)
Co1—O5 ⁱ	2.121 (3)	C9—H9	0.9300
Co1—N1	2.163 (3)	C10—C9	1.377 (6)
Co1—N1 ⁱ	2.163 (3)	C10—C11	1.385 (6)
O1—C1	1.262 (5)	C10—H10	0.9300
O2—C1	1.257 (5)	C11—H11	0.9300
O3—C8	1.193 (7)	C12—C11	1.385 (6)
O4—C14	1.219 (5)	C12—C13	1.378 (6)
O5—H51	0.94 (4)	C13—H13	0.9300
O5—H52	0.92 (2)	C14—N2	1.330 (6)
N1—C9	1.341 (5)	C14—C12	1.511 (6)
N1—C13	1.343 (5)	C15—C16	1.476 (9)
N2—C15	1.473 (7)	C15—H15A	0.9700

supplementary materials

N2—C17	1.501 (7)	C15—H15B	0.9700
C2—C1	1.504 (6)	C16—H16A	0.9600
C2—C3	1.391 (6)	C16—H16B	0.9600
C2—C7	1.392 (6)	C16—H16C	0.9600
C3—C4	1.371 (6)	C17—C18	1.463 (8)
C3—H3	0.9300	C17—H17A	0.9700
C4—C5	1.394 (7)	C17—H17B	0.9700
C4—H4	0.9300	C18—H18A	0.9600
C5—C6	1.391 (6)	C18—H18B	0.9600
C5—C8	1.477 (7)	C18—H18C	0.9600
C6—H6	0.9300		
O1 ⁱ —Co1—O1	180.00 (17)	O3—C8—C5	126.1 (6)
O1 ⁱ —Co1—O5 ⁱ	88.16 (12)	O3—C8—H81	116 (3)
O1—Co1—O5 ⁱ	91.84 (12)	C5—C8—H81	117 (3)
O1 ⁱ —Co1—O5	91.84 (12)	N1—C9—C10	123.3 (4)
O1—Co1—O5	88.16 (12)	N1—C9—H9	118.3
O1 ⁱ —Co1—N1	91.25 (12)	C10—C9—H9	118.3
O1—Co1—N1	88.75 (12)	C9—C10—C11	118.5 (4)
O1 ⁱ —Co1—N1 ⁱ	88.75 (12)	C9—C10—H10	120.7
O1—Co1—N1 ⁱ	91.25 (12)	C11—C10—H10	120.7
O5 ⁱ —Co1—O5	180.00 (18)	C10—C11—H11	120.6
O5 ⁱ —Co1—N1	93.36 (12)	C12—C11—C10	118.8 (4)
O5—Co1—N1	86.64 (12)	C12—C11—H11	120.6
O5 ⁱ —Co1—N1 ⁱ	86.64 (12)	C11—C12—C14	123.6 (4)
O5—Co1—N1 ⁱ	93.36 (12)	C13—C12—C11	119.1 (4)
N1—Co1—N1 ⁱ	180.0 (2)	C13—C12—C14	116.9 (4)
C1—O1—Co1	127.5 (3)	N1—C13—C12	122.6 (4)
Co1—O5—H51	119 (4)	N1—C13—H13	118.7
Co1—O5—H52	97 (3)	C12—C13—H13	118.7
H52—O5—H51	105 (3)	O4—C14—N2	122.1 (4)
C9—N1—Co1	123.6 (3)	O4—C14—C12	117.9 (4)
C9—N1—C13	117.7 (4)	N2—C14—C12	120.0 (4)
C13—N1—Co1	118.6 (3)	N2—C15—C16	113.8 (6)
C14—N2—C15	125.4 (4)	N2—C15—H15A	108.8
C14—N2—C17	116.9 (4)	N2—C15—H15B	108.8
C15—N2—C17	117.6 (4)	C16—C15—H15A	108.8
O1—C1—C2	116.9 (4)	C16—C15—H15B	108.8
O2—C1—O1	125.1 (4)	H15A—C15—H15B	107.7
O2—C1—C2	118.0 (4)	C15—C16—H16A	109.5
C3—C2—C1	120.0 (4)	C15—C16—H16B	109.5
C3—C2—C7	118.7 (4)	C15—C16—H16C	109.5
C7—C2—C1	121.3 (4)	H16A—C16—H16B	109.5
C2—C3—H3	119.9	H16A—C16—H16C	109.5
C4—C3—C2	120.3 (4)	H16B—C16—H16C	109.5
C4—C3—H3	119.9	N2—C17—H17A	109.4
C3—C4—C5	120.5 (4)	N2—C17—H17B	109.4

C3—C4—H4	119.7	C18—C17—N2	111.3 (5)
C5—C4—H4	119.7	C18—C17—H17A	109.4
C4—C5—C8	119.7 (5)	C18—C17—H17B	109.4
C6—C5—C4	119.5 (4)	H17A—C17—H17B	108.0
C6—C5—C8	120.7 (5)	C17—C18—H18A	109.5
C5—C6—H6	120.2	C17—C18—H18B	109.5
C7—C6—C5	119.5 (4)	C17—C18—H18C	109.5
C7—C6—H6	120.2	H18A—C18—H18B	109.5
C2—C7—H7	119.3	H18A—C18—H18C	109.5
C6—C7—C2	121.4 (4)	H18B—C18—H18C	109.5
C6—C7—H7	119.3		
O5 ⁱ —Co1—O1—C1	-11.3 (4)	C1—C2—C3—C4	179.5 (4)
O5—Co1—O1—C1	168.7 (4)	C7—C2—C3—C4	-1.1 (7)
N1—Co1—O1—C1	82.0 (4)	C1—C2—C7—C6	-178.9 (4)
N1 ⁱ —Co1—O1—C1	-98.0 (4)	C3—C2—C7—C6	1.8 (7)
O1 ⁱ —Co1—N1—C9	33.5 (3)	C2—C3—C4—C5	-0.3 (7)
O1—Co1—N1—C9	-146.5 (3)	C3—C4—C5—C6	1.2 (8)
O1 ⁱ —Co1—N1—C13	-148.7 (3)	C3—C4—C5—C8	-179.0 (5)
O1—Co1—N1—C13	31.3 (3)	C4—C5—C6—C7	-0.6 (8)
O5 ⁱ —Co1—N1—C9	-54.7 (3)	C8—C5—C6—C7	179.6 (5)
O5—Co1—N1—C9	125.3 (3)	C4—C5—C8—O3	174.2 (7)
O5 ⁱ —Co1—N1—C13	123.0 (3)	C6—C5—C8—O3	-6.0 (10)
O5—Co1—N1—C13	-57.0 (3)	C2—C7—C6—C5	-1.0 (7)
Co1—O1—C1—O2	27.6 (6)	C11—C10—C9—N1	-0.6 (7)
Co1—O1—C1—C2	-151.1 (3)	C9—C10—C11—C12	1.5 (7)
Co1—N1—C9—C10	177.3 (3)	C13—C12—C11—C10	-1.2 (6)
C13—N1—C9—C10	-0.5 (6)	C14—C12—C11—C10	171.5 (4)
Co1—N1—C13—C12	-177.2 (3)	C11—C12—C13—N1	0.1 (6)
C9—N1—C13—C12	0.7 (6)	C14—C12—C13—N1	-173.1 (4)
C14—N2—C15—C16	109.8 (7)	O4—C14—C12—C11	-114.3 (5)
C17—N2—C15—C16	-70.5 (8)	O4—C14—C12—C13	58.6 (6)
C14—N2—C17—C18	-77.7 (7)	N2—C14—C12—C11	62.5 (6)
C15—N2—C17—C18	102.6 (6)	N2—C14—C12—C13	-124.6 (5)
C3—C2—C1—O1	175.2 (4)	O4—C14—N2—C15	-178.4 (5)
C3—C2—C1—O2	-3.6 (6)	O4—C14—N2—C17	1.9 (7)
C7—C2—C1—O1	-4.1 (6)	C12—C14—N2—C15	5.0 (8)
C7—C2—C1—O2	177.1 (4)	C12—C14—N2—C17	-174.7 (4)

Symmetry codes: (i) $-x+1, -y, -z$.

Hydrogen-bond geometry (\AA , $^\circ$)

$D\cdots H$	$D—H$	$H\cdots A$	$D\cdots A$	$D—H\cdots A$
O5—H51 ⁱⁱ —O4 ⁱⁱ	0.94 (4)	1.86 (4)	2.787 (5)	174 (6)
O5—H52 ⁱ —O2 ⁱ	0.92 (2)	1.73 (4)	2.646 (5)	168 (6)

Symmetry codes: (ii) $-x, -y, -z$; (i) $-x+1, -y, -z$.

supplementary materials

Fig. 1




Fig. 2

