

N-[(2-Hydroxy-1-naphthyl)(3-nitrophenyl)methyl]acetamide

M. Nizam Mohideen,^a A. Subbiah Pandi,^{b*}
N. Panneer Selvam^c and P. T. Perumal^c

^aDepartment of Physics, The New College (Autonomous), Chennai 600 014, India,

^bDepartment of Physics, Presidency College (Autonomous), Chennai 600 005, India, and ^cOrganic Chemistry Division, Central Leather Research Institute, Chennai 600 020, India

Correspondence e-mail: a_spandian@yahoo.com


Received 9 January 2009; accepted 28 February 2009

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.046; wR factor = 0.144; data-to-parameter ratio = 17.0.

The title compound, $\text{C}_{19}\text{H}_{16}\text{N}_2\text{O}_4$, is of interest as a precursor to biologically active substituted quinolines and related compounds. The dihedral angle between the naphthalene ring system and the benzene ring is $81.9(1)^\circ$. The crystal structure is stabilized by $\text{N}-\text{H}\cdots\text{O}$ intermolecular hydrogen bonds, linking the molecules into pairs around a center of symmetry. The crystal structure is further stabilized by intermolecular $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds, which link the molecules into chains running along a axis. An intramolecular $\text{C}-\text{H}\cdots\text{O}$ short contact is also present.

Related literature

For N -(substituted phenyl)acetamides as precursors for the synthesis of heterocyclic compounds, see: Wen *et al.* (2005, 2006). For multicomponent reactions, see: Devi & Bhuyan (2004); Domling & Ugi (2000). For the properties and potential applications of amide-type compounds and their metal ion complexes, see: Saravanan Kumar *et al.* (2005); Yin *et al.* (2004). For related structures, see: Mosslemin *et al.* (2007); Zia-ur-Rehman *et al.* (2008). For bond-length data, see: Allen *et al.* (1987); Liu & Li (2004). For hydrogen-bond motifs, see: Bernstein *et al.* (1995).



Experimental

Crystal data

$\text{C}_{19}\text{H}_{16}\text{N}_2\text{O}_4$	$\gamma = 82.600(3)^\circ$
$M_r = 336.34$	$V = 820.27(14)\text{ \AA}^3$
Triclinic, $P\bar{1}$	$Z = 2$
$a = 7.5261(4)\text{ \AA}$	Mo $K\alpha$ radiation
$b = 8.8635(5)\text{ \AA}$	$\mu = 0.10\text{ mm}^{-1}$
$c = 13.3008(7)\text{ \AA}$	$T = 293\text{ K}$
$\alpha = 74.720(3)^\circ$	$0.4 \times 0.2 \times 0.1\text{ mm}$
$\beta = 73.754(3)^\circ$	

Data collection

Bruker Kappa APEXII CCD diffractometer	9406 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 2004)	3864 independent reflections
$T_{\min} = 0.974$, $T_{\max} = 0.990$	2227 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.026$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.046$	227 parameters
$wR(F^2) = 0.144$	H-atom parameters constrained
$S = 0.99$	$\Delta\rho_{\max} = 0.21\text{ e \AA}^{-3}$
3864 reflections	$\Delta\rho_{\min} = -0.17\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O1—H1 \cdots O4 ⁱ	0.82	1.87	2.646 (2)	158
N1—H1A \cdots O2 ⁱⁱ	0.86	2.35	3.167 (2)	160
C11—H11 \cdots O4	0.98	2.28	2.739 (2)	107

Symmetry codes: (i) $x + 1, y, z$; (ii) $-x + 2, -y + 1, -z + 1$.

Data collection: *APEX2* (Bruker, 2004); cell refinement: *APEX2* and *SAINT* (Bruker, 2004); data reduction: *SAINT* and *XPREP* (Bruker, 2004); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *SHELXL97* and *PLATON* (Spek, 2009)'.

The authors thank Dr Babu Vargheese, SAIF, IIT, Madras, India, for his help with collecting the X-ray intensity data. MNM and ASP thank Dr J. Jothi Kumar, Principal of Presidency College, Chennai, India, for providing the computer and internet facilities.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: JH2076).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N. L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
Bruker (2004). *APEX2*, *SAINT*, *XPREP* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
Devi, I. & Bhuyan, P. J. (2004). *Tetrahedron Lett.* **45**, 8625–8627.
Domling, A. & Ugi, I. (2000). *Angew. Chem. Int. Ed.* **39**, 3168–3210.
Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
Liu, W.-Y. & Li, Y.-Z. (2004). *Acta Cryst. E60*, o694–o695.
Mosslemin, M. H., Arab-Salmanabadi, S. & Masoudi, M. (2007). *Acta Cryst. E63*, o444–o445.

- Saravanakumar, D., Sengottuvelan, N. & Kandaswamy, M. (2005). *Inorg. Chem. Commun.* **8**, 386–389.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.
- Wen, Y.-H., Li, X.-M., Xu, L.-L., Tang, X.-F. & Zhang, S.-S. (2006). *Acta Cryst. E* **62**, o4427–o4428.
- Wen, Y.-H., Zhang, S.-S., Yu, B.-H., Li, X.-M. & Liu, Q. (2005). *Acta Cryst. E* **61**, o347–o348.
- Yin, Z., Yang, Z., He, J. & Cheng, J. P. (2004). *Tetrahedron Lett.* **45**, 6803–6806.
- Zia-ur-Rehman, M., Akbar, N., Arshad, M. N. & Khan, I. U. (2008). *Acta Cryst. E* **64**, o2092.

supplementary materials

Acta Cryst. (2009). E65, o714-o715 [doi:10.1107/S1600536809007442]

N-[(2-Hydroxy-1-naphthyl)(3-nitrophenyl)methyl]acetamide

M. NizamMohideen, A. SubbiahPandi, N. Panneer Selvam and P. T. Perumal

Comment

N-(substituted phenyl)acetamides are well known for their importance as intermediates in organic synthesis. They are used as precursors for the synthesis of many heterocyclic compounds (Wen *et al.*, 2005, 2006). Multi-component reactions (MCRs) have attracted considerable attention in terms of the saving of both energy and raw materials (Devi & Bhuyan, 2004). They have merits over multi-step reactions in several aspects, including the simplicity of a one-pot procedure, possible structural variations and in building up complex molecules (Domling & Ugi, 2000). Much attention has been focused on amide-type compounds and their metal ion complexes for their properties and potential applications including molecular recognition, ion electrodes, photochemistry and topological structures in ion extraction, biochemistry, catalysis and magnetism (Saravanakumar *et al.*, 2005; Yin *et al.*, 2004). The amide linkage [-NHC(O)-] is known to be strong enough to form and maintain protein architectures and has been utilized to create various molecular devices for a spectrum of purposes in organic chemistry. In order to obtain fundamental information about this phenomenon, an X-ray crystal structure analysis of (I) was undertaken.

The conformation of (I), together with the atom-numbering scheme, is shown in Fig. 1. In the structure, all bond lengths and angles are within normal ranges (Allen *et al.*, 1987), and comparable with those in previously reported structure (Mosslemin *et al.*, 2007). The bond distance of C18=O4 is 1.222 (2) Å, which is typical for double bonds (Liu & Li., 2004). The nitro group is slightly twisted out of the plane of the benzene ring, as indicated by O2—N2—C16—C15 and O3—N2—C16—C15 torsion angles of -162.9 (2) and 14.4 (3)°, respectively, and comparable with those in previously reported structure (Zia-ur-Rehman *et al.*, 2008).

The naphthalene ring is planar, the maximum deviation from the least squares plane being -0.027 (1) Å for atom C7. Atom O1 deviating by 0.05 (1) Å from the least squares plane of the naphthalene ring. The dihedral angle between the naphthalene and benzene ring is 81.9 (1)°. The dihedral angle between the fused rings is 1.1 (1)°.

The crystal structure is stabilized by N—H···O intermolecular hydrogen bonds (Table 1, Fig 2.) that generate centrosymmetric hydrogenbonded dimers with a cyclic $R^2_{2}(16)$ ring system (Bernstein, *et al.*, 1995). The crystal structure is further stabilized by intermolecular O—H···O hydrogen bonds link the molecules into chains running along *a* axis.

Experimental

A mixture of 3-nitrobenzaldehyde (10 mmol), β-naphthol (10 mmol) and iodine (0.4 mmol, 4 mol%) were mixed in acetone (5 ml). To that suspension acetyl chloride (2.8 mmol, 0.2 ml) was added and the reaction mixture was stirred at room temperature for 3 h. After the completion of the reaction (as monitored by TLC), saturated sodium thiosulfate solution (5 ml) was added. The precipitated solid was filtered and dried. The dried sample was washed with diethyl ether (2 x 10 ml) and again dried. Single crystals of the title compound suitable for X-ray diffraction were obtained by slow evaporation of a solution in Ethanol.

supplementary materials

Refinement

All H atoms were positioned geometrically, with N—H = 0.86 and C—H = 0.93, 0.98 and 0.96 Å aromatic, methylene and methyl H, respectively, and constrained to ride on their parent atoms, with $U_{\text{iso}}(\text{H}) = xU_{\text{eq}}(\text{C, N})$, where $x = 1.5$ for methyl H, and $x = 1.2$ for all other H atoms.

Figures




Fig. 1. The molecular configuration and atom-numbering scheme for (I). Displacement ellipsoids are drawn at the 30% probability level.




Fig. 2. Part of the crystal structure of (I), showing the $R^2_2(16)$ rings. For the sake of clarity, H atoms not participating in the hydrogen bonding have been omitted. Hydrogen bonding is shown as dashed lines. [Symmetry codes: (*) $-x + 2, -y + 1, -z + 1$]

N-[(2-Hydroxy-1-naphthyl)(3-nitrophenyl)methyl]acetamide

Crystal data

$\text{C}_{19}\text{H}_{16}\text{N}_2\text{O}_4$	$Z = 2$
$M_r = 336.34$	$F_{000} = 352$
Triclinic, $P\bar{1}$	$D_x = 1.362 \text{ Mg m}^{-3}$
Hall symbol: -P 1	Mo $K\alpha$ radiation
$a = 7.5261 (4) \text{ \AA}$	$\lambda = 0.71073 \text{ \AA}$
$b = 8.8635 (5) \text{ \AA}$	Cell parameters from 3864 reflections
$c = 13.3008 (7) \text{ \AA}$	$\theta = 2.5\text{--}25^\circ$
$\alpha = 74.720 (3)^\circ$	$\mu = 0.10 \text{ mm}^{-1}$
$\beta = 73.754 (3)^\circ$	$T = 293 \text{ K}$
$\gamma = 82.600 (3)^\circ$	Needle, colourless
$V = 820.27 (14) \text{ \AA}^3$	$0.4 \times 0.2 \times 0.1 \text{ mm}$

Data collection

Bruker Kappa APEXII CCD diffractometer	3864 independent reflections
Radiation source: fine-focus sealed tube	2227 reflections with $I > 2\sigma(I)$

Monochromator: graphite	$R_{\text{int}} = 0.026$
$T = 293 \text{ K}$	$\theta_{\text{max}} = 28.4^\circ$
ω and φ scan	$\theta_{\text{min}} = 2.6^\circ$
Absorption correction: Multi-scan (SADABS; Bruker, 2004)	$h = -9 \rightarrow 10$
$T_{\text{min}} = 0.974, T_{\text{max}} = 0.990$	$k = -11 \rightarrow 11$
9406 measured reflections	$l = -17 \rightarrow 17$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.046$	H-atom parameters constrained
$wR(F^2) = 0.144$	$w = 1/[\sigma^2(F_o^2) + (0.0612P)^2 + 0.220P]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 0.99$	$(\Delta/\sigma)_{\text{max}} = 0.009$
3864 reflections	$\Delta\rho_{\text{max}} = 0.21 \text{ e \AA}^{-3}$
227 parameters	$\Delta\rho_{\text{min}} = -0.17 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	1.11375 (16)	0.78003 (16)	0.26550 (11)	0.0490 (4)
H1	1.2259	0.7587	0.2525	0.074*
O2	0.9758 (4)	0.5649 (2)	0.62604 (17)	0.1025 (7)
O3	0.8781 (4)	0.6954 (3)	0.74538 (16)	0.1150 (8)
O4	0.46636 (19)	0.7392 (2)	0.26984 (17)	0.0801 (6)
N1	0.74868 (19)	0.71417 (18)	0.29573 (12)	0.0401 (4)
H1A	0.8443	0.6528	0.3045	0.048*
N2	0.8965 (3)	0.6794 (3)	0.65559 (16)	0.0667 (5)
C1	1.0777 (2)	0.9135 (2)	0.19395 (14)	0.0385 (4)
C2	1.2195 (2)	0.9945 (3)	0.11175 (16)	0.0502 (5)
H2	1.3425	0.9575	0.1056	0.060*

supplementary materials

C3	1.1769 (3)	1.1262 (3)	0.04174 (17)	0.0536 (5)
H3	1.2720	1.1793	-0.0117	0.064*
C4	0.9922 (3)	1.1848 (2)	0.04788 (15)	0.0473 (5)
C5	0.9457 (4)	1.3208 (3)	-0.02594 (17)	0.0632 (6)
H5	1.0402	1.3758	-0.0786	0.076*
C6	0.7679 (4)	1.3729 (3)	-0.0220 (2)	0.0732 (7)
H6	0.7408	1.4619	-0.0721	0.088*
C7	0.6253 (4)	1.2929 (3)	0.05736 (19)	0.0634 (6)
H7	0.5029	1.3285	0.0595	0.076*
C8	0.6626 (3)	1.1628 (2)	0.13215 (17)	0.0495 (5)
H8	0.5649	1.1126	0.1852	0.059*
C9	0.8474 (2)	1.1027 (2)	0.13058 (14)	0.0384 (4)
C10	0.8943 (2)	0.9665 (2)	0.20492 (14)	0.0353 (4)
C11	0.7483 (2)	0.8774 (2)	0.29803 (14)	0.0356 (4)
H11	0.6276	0.9264	0.2889	0.043*
C12	0.7637 (2)	0.8970 (2)	0.40573 (14)	0.0366 (4)
C13	0.7214 (2)	1.0446 (2)	0.42690 (16)	0.0448 (5)
H13	0.6865	1.1272	0.3751	0.054*
C14	0.7299 (3)	1.0718 (3)	0.52306 (17)	0.0508 (5)
H14	0.6998	1.1715	0.5356	0.061*
C15	0.7827 (3)	0.9515 (3)	0.60021 (16)	0.0500 (5)
H15	0.7880	0.9677	0.6656	0.060*
C16	0.8275 (3)	0.8069 (2)	0.57761 (15)	0.0452 (5)
C17	0.8192 (2)	0.7764 (2)	0.48202 (14)	0.0411 (4)
H17	0.8504	0.6766	0.4697	0.049*
C18	0.6061 (2)	0.6567 (2)	0.28056 (16)	0.0457 (5)
C19	0.6235 (3)	0.4876 (3)	0.2787 (2)	0.0719 (7)
H19A	0.5450	0.4293	0.3438	0.108*
H19B	0.7499	0.4483	0.2734	0.108*
H19C	0.5866	0.4764	0.2176	0.108*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0284 (6)	0.0571 (9)	0.0628 (9)	0.0062 (6)	-0.0192 (6)	-0.0126 (7)
O2	0.163 (2)	0.0682 (13)	0.0971 (15)	0.0310 (13)	-0.0825 (15)	-0.0215 (12)
O3	0.189 (3)	0.1062 (17)	0.0557 (12)	0.0012 (16)	-0.0525 (14)	-0.0122 (11)
O4	0.0403 (8)	0.0771 (12)	0.1474 (17)	0.0087 (8)	-0.0497 (10)	-0.0471 (12)
N1	0.0309 (7)	0.0380 (9)	0.0574 (9)	0.0045 (6)	-0.0211 (7)	-0.0146 (7)
N2	0.0861 (14)	0.0639 (14)	0.0559 (12)	-0.0130 (11)	-0.0321 (10)	-0.0056 (11)
C1	0.0318 (8)	0.0457 (11)	0.0440 (10)	0.0000 (8)	-0.0138 (7)	-0.0177 (9)
C2	0.0313 (9)	0.0701 (14)	0.0556 (12)	-0.0066 (9)	-0.0079 (8)	-0.0277 (11)
C3	0.0516 (12)	0.0623 (14)	0.0462 (11)	-0.0172 (10)	-0.0023 (9)	-0.0161 (11)
C4	0.0581 (12)	0.0454 (12)	0.0416 (10)	-0.0068 (9)	-0.0100 (9)	-0.0170 (9)
C5	0.0878 (18)	0.0488 (13)	0.0467 (12)	-0.0081 (12)	-0.0087 (12)	-0.0080 (11)
C6	0.104 (2)	0.0478 (14)	0.0603 (15)	0.0171 (14)	-0.0250 (14)	-0.0070 (12)
C7	0.0736 (15)	0.0513 (13)	0.0669 (15)	0.0205 (12)	-0.0295 (12)	-0.0161 (12)
C8	0.0502 (11)	0.0455 (12)	0.0554 (12)	0.0086 (9)	-0.0204 (9)	-0.0145 (10)

C9	0.0424 (9)	0.0368 (10)	0.0410 (10)	0.0010 (8)	-0.0135 (8)	-0.0162 (8)
C10	0.0314 (8)	0.0386 (10)	0.0418 (10)	-0.0003 (7)	-0.0128 (7)	-0.0165 (8)
C11	0.0268 (8)	0.0343 (10)	0.0496 (10)	0.0039 (7)	-0.0154 (7)	-0.0131 (8)
C12	0.0238 (7)	0.0404 (10)	0.0462 (10)	-0.0016 (7)	-0.0079 (7)	-0.0126 (8)
C13	0.0374 (9)	0.0424 (11)	0.0554 (12)	0.0021 (8)	-0.0107 (8)	-0.0162 (9)
C14	0.0437 (10)	0.0500 (12)	0.0622 (13)	-0.0030 (9)	-0.0055 (9)	-0.0280 (11)
C15	0.0436 (10)	0.0636 (14)	0.0466 (11)	-0.0122 (10)	-0.0046 (9)	-0.0229 (11)
C16	0.0413 (10)	0.0504 (12)	0.0440 (11)	-0.0097 (9)	-0.0106 (8)	-0.0084 (9)
C17	0.0389 (9)	0.0405 (11)	0.0468 (11)	-0.0037 (8)	-0.0121 (8)	-0.0133 (9)
C18	0.0350 (9)	0.0511 (12)	0.0568 (12)	-0.0047 (8)	-0.0161 (8)	-0.0175 (10)
C19	0.0677 (15)	0.0586 (15)	0.103 (2)	-0.0135 (12)	-0.0270 (14)	-0.0323 (15)

Geometric parameters (Å, °)

O1—C1	1.359 (2)	C7—H7	0.9300
O1—H1	0.8200	C8—C9	1.420 (3)
O2—N2	1.209 (3)	C8—H8	0.9300
O3—N2	1.207 (3)	C9—C10	1.420 (2)
O4—C18	1.222 (2)	C10—C11	1.520 (2)
N1—C18	1.329 (2)	C11—C12	1.525 (2)
N1—C11	1.455 (2)	C11—H11	0.9800
N1—H1A	0.8600	C12—C17	1.378 (2)
N2—C16	1.468 (3)	C12—C13	1.389 (3)
C1—C10	1.380 (2)	C13—C14	1.382 (3)
C1—C2	1.407 (3)	C13—H13	0.9300
C2—C3	1.353 (3)	C14—C15	1.375 (3)
C2—H2	0.9300	C14—H14	0.9300
C3—C4	1.408 (3)	C15—C16	1.372 (3)
C3—H3	0.9300	C15—H15	0.9300
C4—C5	1.414 (3)	C16—C17	1.387 (3)
C4—C9	1.428 (3)	C17—H17	0.9300
C5—C6	1.349 (3)	C18—O4	1.222 (2)
C5—H5	0.9300	C18—C19	1.494 (3)
C6—C7	1.391 (4)	C19—H19A	0.9600
C6—H6	0.9300	C19—H19B	0.9600
C7—C8	1.366 (3)	C19—H19C	0.9600
C1—O1—H1	109.5	C9—C10—C11	121.98 (14)
C18—N1—C11	122.19 (15)	N1—C11—C10	112.60 (14)
C18—N1—H1A	118.9	N1—C11—C12	112.92 (14)
C11—N1—H1A	118.9	C10—C11—C12	111.04 (13)
O3—N2—O2	122.6 (2)	N1—C11—H11	106.6
O3—N2—C16	118.8 (2)	C10—C11—H11	106.6
O2—N2—C16	118.6 (2)	C12—C11—H11	106.6
O1—C1—C10	116.89 (16)	C17—C12—C13	118.70 (17)
O1—C1—C2	122.07 (15)	C17—C12—C11	123.28 (16)
C10—C1—C2	121.03 (17)	C13—C12—C11	118.02 (16)
C3—C2—C1	120.01 (18)	C14—C13—C12	121.61 (19)
C3—C2—H2	120.0	C14—C13—H13	119.2
C1—C2—H2	120.0	C12—C13—H13	119.2

supplementary materials

C2—C3—C4	121.59 (19)	C15—C14—C13	120.02 (19)
C2—C3—H3	119.2	C15—C14—H14	120.0
C4—C3—H3	119.2	C13—C14—H14	120.0
C3—C4—C5	122.1 (2)	C16—C15—C14	117.86 (18)
C3—C4—C9	118.75 (18)	C16—C15—H15	121.1
C5—C4—C9	119.13 (19)	C14—C15—H15	121.1
C6—C5—C4	121.8 (2)	C15—C16—C17	123.22 (19)
C6—C5—H5	119.1	C15—C16—N2	118.79 (18)
C4—C5—H5	119.1	C17—C16—N2	117.91 (19)
C5—C6—C7	119.7 (2)	C12—C17—C16	118.56 (18)
C5—C6—H6	120.2	C12—C17—H17	120.7
C7—C6—H6	120.2	C16—C17—H17	120.7
C8—C7—C6	121.0 (2)	O4—C18—N1	120.94 (18)
C8—C7—H7	119.5	O4—C18—N1	120.94 (18)
C6—C7—H7	119.5	O4—C18—C19	121.76 (18)
C7—C8—C9	121.4 (2)	O4—C18—C19	121.76 (18)
C7—C8—H8	119.3	N1—C18—C19	117.29 (17)
C9—C8—H8	119.3	C18—C19—H19A	109.5
C10—C9—C8	123.85 (17)	C18—C19—H19B	109.5
C10—C9—C4	119.07 (16)	H19A—C19—H19B	109.5
C8—C9—C4	117.07 (17)	C18—C19—H19C	109.5
C1—C10—C9	119.52 (16)	H19A—C19—H19C	109.5
C1—C10—C11	118.50 (15)	H19B—C19—H19C	109.5
O1—C1—C2—C3	179.40 (18)	C1—C10—C11—N1	−58.74 (19)
C10—C1—C2—C3	−0.2 (3)	C9—C10—C11—N1	122.18 (17)
C1—C2—C3—C4	−0.7 (3)	C1—C10—C11—C12	69.00 (19)
C2—C3—C4—C5	−178.9 (2)	C9—C10—C11—C12	−110.08 (17)
C2—C3—C4—C9	0.3 (3)	N1—C11—C12—C17	15.6 (2)
C3—C4—C5—C6	177.5 (2)	C10—C11—C12—C17	−111.95 (18)
C9—C4—C5—C6	−1.7 (3)	N1—C11—C12—C13	−165.13 (14)
C4—C5—C6—C7	0.9 (4)	C10—C11—C12—C13	67.30 (18)
C5—C6—C7—C8	0.6 (4)	C17—C12—C13—C14	−1.4 (3)
C6—C7—C8—C9	−1.2 (3)	C11—C12—C13—C14	179.28 (16)
C7—C8—C9—C10	−178.80 (19)	C12—C13—C14—C15	0.6 (3)
C7—C8—C9—C4	0.4 (3)	C13—C14—C15—C16	0.6 (3)
C3—C4—C9—C10	1.0 (3)	C14—C15—C16—C17	−1.0 (3)
C5—C4—C9—C10	−179.76 (18)	C14—C15—C16—N2	175.85 (18)
C3—C4—C9—C8	−178.19 (18)	O3—N2—C16—C15	14.4 (3)
C5—C4—C9—C8	1.0 (3)	O2—N2—C16—C15	−162.9 (2)
O1—C1—C10—C9	−178.11 (15)	O3—N2—C16—C17	−168.6 (2)
C2—C1—C10—C9	1.5 (3)	O2—N2—C16—C17	14.1 (3)
O1—C1—C10—C11	2.8 (2)	C13—C12—C17—C16	1.1 (2)
C2—C1—C10—C11	−177.60 (16)	C11—C12—C17—C16	−179.69 (16)
C8—C9—C10—C1	177.26 (17)	C15—C16—C17—C12	0.1 (3)
C4—C9—C10—C1	−1.9 (2)	N2—C16—C17—C12	−176.74 (16)
C8—C9—C10—C11	−3.7 (3)	C11—N1—C18—O4	−1.4 (3)
C4—C9—C10—C11	177.17 (16)	C11—N1—C18—O4	−1.4 (3)
C18—N1—C11—C10	−114.89 (18)	C11—N1—C18—C19	179.39 (19)
C18—N1—C11—C12	118.37 (18)		

Hydrogen-bond geometry (Å, °)

<i>D—H···A</i>	<i>D—H</i>	<i>H···A</i>	<i>D···A</i>	<i>D—H···A</i>
O1—H1···O4 ⁱ	0.82	1.87	2.646 (2)	158
N1—H1A···O2 ⁱⁱ	0.86	2.35	3.167 (2)	160
C11—H11···O4	0.98	2.28	2.739 (2)	107

Symmetry codes: (i) $x+1, y, z$; (ii) $-x+2, -y+1, -z+1$.

supplementary materials

Fig. 1




Fig. 2

