

2-Ethyl-5-nitroaniline**Yan Chen, Zheng Fang* and Ping Wei**

College of Life Science and Pharmaceutical Engineering, Nanjing University of Technology, Xinmofan Road No. 5 Nanjing, Nanjing 210009, People's Republic of China
Correspondence e-mail: fzcpu@163.com


Received 16 March 2009; accepted 17 March 2009

Key indicators: single-crystal X-ray study; $T = 298\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.055; wR factor = 0.153; data-to-parameter ratio = 13.5.

The molecule of the title compound, $\text{C}_8\text{H}_{10}\text{N}_2\text{O}_2$, is nearly planar [maximum deviation of $0.163(3)\text{ \AA}$ for one of the O atoms of the NO_2 group]. In the crystal structure, weak intermolecular $\text{N}-\text{H}\cdots\text{N}$ and $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds link the molecules into chains, forming $R_2^2(10)$ ring motifs.

Related literature

For a related structure, see: Corwin (1955). For bond-length data, see: Allen *et al.* (1987). For ring motifs, see: Bernstein *et al.* (1995).

**Experimental***Crystal data*

$\text{C}_8\text{H}_{10}\text{N}_2\text{O}_2$
 $M_r = 166.18$
Monoclinic, $C2/c$
 $a = 23.037(5)\text{ \AA}$
 $b = 3.9540(8)\text{ \AA}$

$c = 18.393(4)\text{ \AA}$
 $\beta = 104.51(3)^\circ$
 $V = 1621.9(6)\text{ \AA}^3$
 $Z = 8$
Mo $K\alpha$ radiation

$\mu = 0.10\text{ mm}^{-1}$
 $T = 298\text{ K}$

$0.20 \times 0.10 \times 0.10\text{ mm}$

Data collection

Enraf–Nonius CAD-4
diffractometer
Absorption correction: ψ scan
(North *et al.*, 1968)
 $T_{\min} = 0.980$, $T_{\max} = 0.990$
2937 measured reflections

1474 independent reflections
906 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.056$
3 standard reflections
frequency: 120 min
intensity decay: 1%

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.055$
 $wR(F^2) = 0.153$
 $S = 1.01$
1474 reflections

109 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.23\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.21\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N}2-\text{H}2\text{B}\cdots\text{N}2^{\text{i}}$	0.86	2.62	3.423 (3)	156
$\text{C}7-\text{H}7\text{A}\cdots\text{O}1^{\text{ii}}$	0.93	2.60	3.417 (4)	147

Symmetry codes: (i) $-x + \frac{1}{2}, y - \frac{1}{2}, -z + \frac{1}{2}$; (ii) $-x + \frac{1}{2}, -y + \frac{3}{2}, -z$.

Data collection: *CAD-4 Software* (Enraf–Nonius, 1989); cell refinement: *CAD-4 Software*; data reduction: *XCAD4* (Harms & Wocadlo, 1995); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997) and *PLATON* (Spek, 2009); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008).

The authors thank the Center of Testing and Analysis, Nanjing University, for support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HK2644).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Corwin, H. (1955). *J. Org. Chem.* **20**, 1026–1029.
- Enraf–Nonius (1989). *CAD-4 Software*. Enraf–Nonius, Delft. The Netherlands.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Harms, K. & Wocadlo, S. (1995). *XCAD4*. University of Marburg, Germany.
- North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst. A* **24**, 351–359.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.

supplementary materials

Acta Cryst. (2009). E65, o827 [doi:10.1107/S1600536809009878]

2-Ethyl-5-nitroaniline

Y. Chen, Z. Fang and P. Wei

Comment

Some derivatives of aniline are important chemical materials. We report herein the crystal structure of the title compound.

In the molecule of the title compound (Fig. 1), the bond lengths (Allen *et al.*, 1987) and angles are within normal ranges. Ring A (C3-C8) is, of course, planar. Atoms C1, C2, N1, N2, O1 and O2 are -0.067 (3), -0.028 (2), -0.035 (3), -0.055 (3), 0.054 (3) and -0.163 (3) Å away from the ring plane of A, respectively.

In the crystal structure, weak intermolecular N-H···N and C-H···O hydrogen bonds (Table 1) link the molecules into chains, forming $R_2^2(10)$ ring motifs (Fig. 2) (Bernstein *et al.*, 1995), in which they may be effective in the stabilization of the structure.

Experimental

For the preparation of the title compound, 2-ethylaniline (12.1 g) was dissolved in concentrated sulfuric acid (50 ml). The mixture was cooled to 273 K, and nitric acid (6.37 ml) was added in small portions. The mixture was stirred at 295 K for 0.5 h. Then, it was poured into a large volume of ice, used sodium hydroxide to neutralize excess acid, filtered, and dried. The compound was crystallized from cyclohexane (Corwin, 1955). Crystals suitable for X-ray analysis were obtained by slow evaporation of a methanol solution.

Refinement

H atoms were positioned geometrically, with N-H = 0.86 Å (for NH₂) and C-H = 0.93, 0.97 and 0.96 Å for aromatic, methylene and methyl H, respectively, and constrained to ride on their parent atoms, with $U_{\text{iso}}(\text{H}) = xU_{\text{eq}}(\text{C}, \text{N})$, where $x = 1.5$ for methyl H and $x = 1.2$ for all other H atoms.

Figures




Fig. 1. The molecular structure of the title molecule, with the atom-numbering scheme. Displacement ellipsoids are drawn at the 30% probability level.

supplementary materials




Fig. 2. A partial packing diagram of the title compound. Hydrogen bonds are shown as dashed lines.

2-Ethyl-5-nitroaniline

Crystal data

C ₈ H ₁₀ N ₂ O ₂	$F_{000} = 704$
$M_r = 166.18$	$D_x = 1.361 \text{ Mg m}^{-3}$
Monoclinic, C2/c	Mo $K\alpha$ radiation
Hall symbol: -C 2yc	$\lambda = 0.71073 \text{ \AA}$
$a = 23.037 (5) \text{ \AA}$	Cell parameters from 25 reflections
$b = 3.9540 (8) \text{ \AA}$	$\theta = 10\text{--}13^\circ$
$c = 18.393 (4) \text{ \AA}$	$\mu = 0.10 \text{ mm}^{-1}$
$\beta = 104.51 (3)^\circ$	$T = 298 \text{ K}$
$V = 1621.9 (6) \text{ \AA}^3$	Block, colorless
$Z = 8$	$0.20 \times 0.10 \times 0.10 \text{ mm}$

Data collection

Enraf–Nonius CAD-4 diffractometer	$R_{\text{int}} = 0.056$
Radiation source: fine-focus sealed tube	$\theta_{\text{max}} = 25.3^\circ$
Monochromator: graphite	$\theta_{\text{min}} = 1.8^\circ$
$T = 298 \text{ K}$	$h = -27 \rightarrow 27$
$\omega/2\theta$ scans	$k = 0 \rightarrow 4$
Absorption correction: ψ scan (North <i>et al.</i> , 1968)	$l = -22 \rightarrow 22$
$T_{\text{min}} = 0.980$, $T_{\text{max}} = 0.990$	3 standard reflections
2937 measured reflections	every 120 min
1474 independent reflections	intensity decay: 1%
906 reflections with $I > 2\sigma(I)$	

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.055$	H-atom parameters constrained
$wR(F^2) = 0.153$	$w = 1/[\sigma^2(F_o^2) + (0.06P)^2 + P]$
$S = 1.01$	where $P = (F_o^2 + 2F_c^2)/3$
	$(\Delta/\sigma)_{\text{max}} < 0.001$

1474 reflections $\Delta\rho_{\max} = 0.23 \text{ e \AA}^{-3}$
 109 parameters $\Delta\rho_{\min} = -0.21 \text{ e \AA}^{-3}$
 Primary atom site location: structure-invariant direct Extinction correction: none
 methods

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.18034 (10)	0.6924 (7)	-0.09151 (11)	0.0931 (9)
O2	0.08462 (10)	0.6656 (7)	-0.12713 (11)	0.0921 (8)
N1	0.13189 (12)	0.6134 (7)	-0.08113 (12)	0.0598 (6)
N2	0.23320 (8)	0.2602 (7)	0.17155 (11)	0.0643 (8)
H2A	0.2663	0.3390	0.1651	0.077*
H2B	0.2322	0.1704	0.2138	0.077*
C1	0.06424 (11)	-0.1027 (8)	0.20919 (14)	0.0592 (8)
H1A	0.0687	-0.1949	0.2586	0.089*
H1B	0.0479	-0.2721	0.1723	0.089*
H1C	0.0377	0.0881	0.2025	0.089*
C2	0.12500 (10)	0.0091 (7)	0.19996 (12)	0.0489 (7)
H2C	0.1413	0.1738	0.2388	0.059*
H2D	0.1515	-0.1852	0.2087	0.059*
C3	0.12627 (10)	0.1609 (6)	0.12529 (12)	0.0406 (6)
C4	0.07514 (10)	0.1952 (7)	0.06693 (13)	0.0458 (6)
H4A	0.0388	0.1174	0.0738	0.055*
C5	0.07620 (11)	0.3397 (7)	-0.00061 (13)	0.0479 (7)
H5A	0.0413	0.3608	-0.0389	0.058*
C6	0.13008 (10)	0.4519 (7)	-0.01007 (12)	0.0445 (6)
C7	0.18252 (10)	0.4228 (7)	0.04537 (13)	0.0476 (7)
H7A	0.2185	0.4993	0.0372	0.057*
C8	0.18102 (10)	0.2780 (7)	0.11353 (12)	0.0442 (6)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0812 (15)	0.132 (2)	0.0759 (14)	-0.0154 (16)	0.0390 (12)	0.0189 (15)
O2	0.0889 (15)	0.121 (2)	0.0588 (12)	-0.0001 (15)	0.0045 (11)	0.0277 (14)

supplementary materials

N1	0.0728 (16)	0.0621 (16)	0.0473 (12)	-0.0051 (14)	0.0201 (12)	-0.0025 (12)
N2	0.0382 (11)	0.101 (2)	0.0502 (12)	-0.0073 (13)	0.0052 (9)	-0.0046 (14)
C1	0.0702 (18)	0.0533 (18)	0.0597 (16)	-0.0030 (15)	0.0266 (13)	0.0046 (14)
C2	0.0550 (15)	0.0431 (16)	0.0493 (14)	0.0009 (12)	0.0142 (12)	-0.0057 (12)
C3	0.0433 (13)	0.0346 (14)	0.0441 (13)	0.0014 (11)	0.0115 (10)	-0.0080 (11)
C4	0.0402 (13)	0.0443 (16)	0.0529 (14)	-0.0059 (12)	0.0117 (11)	-0.0044 (13)
C5	0.0455 (14)	0.0474 (16)	0.0471 (14)	-0.0007 (13)	0.0044 (11)	-0.0016 (13)
C6	0.0526 (14)	0.0411 (15)	0.0415 (13)	0.0020 (12)	0.0148 (11)	-0.0082 (12)
C7	0.0438 (13)	0.0515 (17)	0.0519 (14)	-0.0063 (13)	0.0202 (11)	-0.0094 (13)
C8	0.0394 (13)	0.0479 (16)	0.0448 (13)	0.0022 (12)	0.0097 (10)	-0.0105 (12)

Geometric parameters (\AA , $^\circ$)

O1—N1	1.219 (3)	C2—H2C	0.9700
O2—N1	1.218 (3)	C2—H2D	0.9700
N1—C6	1.465 (3)	C3—C4	1.387 (3)
N2—C8	1.395 (3)	C3—C8	1.410 (3)
N2—H2A	0.8600	C4—C5	1.373 (3)
N2—H2B	0.8600	C4—H4A	0.9300
C1—C2	1.517 (3)	C5—C6	1.370 (3)
C1—H1A	0.9600	C5—H5A	0.9300
C1—H1B	0.9600	C6—C7	1.377 (3)
C1—H1C	0.9600	C7—C8	1.387 (3)
C2—C3	1.506 (3)	C7—H7A	0.9300
O2—N1—O1	122.8 (2)	C4—C3—C8	117.9 (2)
O2—N1—C6	118.3 (2)	C4—C3—C2	122.5 (2)
O1—N1—C6	118.9 (2)	C8—C3—C2	119.56 (19)
C8—N2—H2A	120.0	C5—C4—C3	122.5 (2)
C8—N2—H2B	120.0	C5—C4—H4A	118.8
H2A—N2—H2B	120.0	C3—C4—H4A	118.8
C2—C1—H1A	109.5	C6—C5—C4	118.2 (2)
C2—C1—H1B	109.5	C6—C5—H5A	120.9
H1A—C1—H1B	109.5	C4—C5—H5A	120.9
C2—C1—H1C	109.5	C5—C6—C7	122.3 (2)
H1A—C1—H1C	109.5	C5—C6—N1	118.9 (2)
H1B—C1—H1C	109.5	C7—C6—N1	118.8 (2)
C3—C2—C1	116.59 (19)	C6—C7—C8	119.2 (2)
C3—C2—H2C	108.1	C6—C7—H7A	120.4
C1—C2—H2C	108.1	C8—C7—H7A	120.4
C3—C2—H2D	108.1	C7—C8—N2	120.0 (2)
C1—C2—H2D	108.1	C7—C8—C3	120.0 (2)
H2C—C2—H2D	107.3	N2—C8—C3	119.9 (2)
C1—C2—C3—C4	-0.7 (4)	O1—N1—C6—C7	5.9 (4)
C1—C2—C3—C8	178.4 (2)	C5—C6—C7—C8	-0.6 (4)
C8—C3—C4—C5	-0.5 (4)	N1—C6—C7—C8	178.4 (2)
C2—C3—C4—C5	178.7 (3)	C6—C7—C8—N2	-177.1 (2)
C3—C4—C5—C6	0.3 (4)	C6—C7—C8—C3	0.4 (4)
C4—C5—C6—C7	0.3 (4)	C4—C3—C8—C7	0.1 (3)
C4—C5—C6—N1	-178.7 (2)	C2—C3—C8—C7	-179.0 (2)

O2—N1—C6—C5	5.6 (4)	C4—C3—C8—N2	177.6 (2)
O1—N1—C6—C5	−175.1 (3)	C2—C3—C8—N2	−1.5 (4)
O2—N1—C6—C7	−173.4 (2)		

Hydrogen-bond geometry (\AA , °)

$D\text{—H}\cdots A$	$D\text{—H}$	$\text{H}\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
N2—H2B···N2 ⁱ	0.86	2.62	3.423 (3)	156
C7—H7A···O1 ⁱⁱ	0.93	2.60	3.417 (4)	147

Symmetry codes: (i) $-x+1/2, y-1/2, -z+1/2$; (ii) $-x+1/2, -y+3/2, -z$.

supplementary materials

Fig. 1




Fig. 2

