

## 5-Benzenesulfonamido-2-chlorobenzoic acid

Muhammad Nadeem Arshad,<sup>a</sup> M. Nawaz Tahir,<sup>b\*</sup>  
Islam Ullah Khan,<sup>a</sup> Muhammad Shafiq<sup>a</sup> and  
Hafiz Muhammad Adeel Sharif<sup>a</sup>

<sup>a</sup>Department of Chemistry, Government College University, Lahore, Pakistan, and

<sup>b</sup>Department of Physics, University of Sargodha, Sargodha, Pakistan

Correspondence e-mail: dmntahir\_uos@yahoo.com


Received 9 March 2009; accepted 17 March 2009

Key indicators: single-crystal X-ray study;  $T = 296$  K; mean  $\sigma(\text{C}-\text{C}) = 0.003$  Å;  $R$  factor = 0.039;  $wR$  factor = 0.108; data-to-parameter ratio = 18.0.

In the title compound,  $\text{C}_{13}\text{H}_{10}\text{ClNO}_4\text{S}$ , the dihedral angle between the aromatic ring planes is  $87.07(6)^\circ$  and an intramolecular  $\text{C}-\text{H}\cdots\text{O}$  interaction occurs. In the crystal, inversion dimers linked by two  $\text{O}-\text{H}\cdots\text{O}$  hydrogen bonds arise from the carboxyl groups.  $\text{N}-\text{H}\cdots\text{O}$  hydrogen bonds link the dimers into chains and short  $\text{C}-\text{Cl}\cdots\pi$  and  $\text{S}-\text{O}\cdots\pi$  contacts are also seen.

## Related literature

For related structures: see: Arshad *et al.* (2008); Arshad, Khan *et al.* (2009); Arshad, Tahir *et al.* (2009). For chemical background, see: Bouchain *et al.* (2003). For graph-set theory, see: Bernstein *et al.* (1995).



## Experimental

## Crystal data

$\text{C}_{13}\text{H}_{10}\text{ClNO}_4\text{S}$

$M_r = 311.73$

Monoclinic,  $P2_1/c$

$a = 11.7139(4)$  Å

$b = 5.3957(2)$  Å

$c = 20.7565(8)$  Å

$\beta = 91.483(2)^\circ$

$V = 1311.47(8)$  Å<sup>3</sup>

$Z = 4$

Mo  $K\alpha$  radiation

$\mu = 0.46$  mm<sup>-1</sup>

$T = 296$  K

$0.24 \times 0.18 \times 0.15$  mm

## Data collection

Bruker Kappa APEXII CCD

diffractometer

Absorption correction: multi-scan

(*SADABS*; Bruker, 2005)

$T_{\min} = 0.939$ ,  $T_{\max} = 0.940$

14693 measured reflections

3269 independent reflections

2513 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.029$

## Refinement

$R[F^2 > 2\sigma(F^2)] = 0.039$

$wR(F^2) = 0.108$

$S = 1.02$

3269 reflections

182 parameters

H-atom parameters constrained

$\Delta\rho_{\text{max}} = 0.37$  e Å<sup>-3</sup>

$\Delta\rho_{\text{min}} = -0.36$  e Å<sup>-3</sup>

## Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{O1}-\text{H1O}\cdots\text{O2}^i$	0.82	1.83	2.648 (2)	178
$\text{N1}-\text{H1N}\cdots\text{O3}^{ii}$	0.86	2.16	2.898 (2)	144
$\text{C4}-\text{H4}\cdots\text{O4}$	0.93	2.41	3.052 (3)	126
$\text{C6}-\text{C11}\cdots\text{CgB}^{iii}$	1.73 (1)	3.81 (1)	4.605 (2)	106 (1)
$\text{S1}-\text{O4}\cdots\text{CgA}^{iv}$	1.43 (1)	3.14 (1)	4.2532 (9)	134 (1)

Symmetry codes: (i)  $-x, -y + 3, -z$ ; (ii)  $-x, y + \frac{1}{2}, -z + \frac{1}{2}$ ; (iii)  $x, -y + \frac{3}{2}, z - \frac{1}{2}$ ; (iv)  $x, y - 1, z$ . *CgA* and *CgB* are the centroids of the C1-C6 and C8-C13 benzene rings, respectively.

Data collection: *APEX2* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997) and *PLATON* (Spek, 2009); software used to prepare material for publication: *WinGX* (Farrugia, 1999) and *PLATON*.

MNA gratefully acknowledges the Higher Education Commission, Islamabad, Pakistan, for providing him with a Scholarship under the Indigenous PhD Program (PIN 042-120607-PS2-183).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB2925).

## References

- Arshad, M. N., Khan, I. U., Shafiq, M. & Mukhtar, A. (2009). *Acta Cryst.* **E65**, o549.
- Arshad, M. N., Tahir, M. N., Khan, I. U., Shafiq, M. & Siddiqui, W. A. (2008). *Acta Cryst.* **E64**, o2045.
- Arshad, M. N., Tahir, M. N., Khan, I. U., Siddiqui, W. A. & Shafiq, M. (2009). *Acta Cryst.* **E65**, o281.
- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555-1573.
- Bouchain, G., Leit, S., Frechette, S., Khalil, E. A., Lavoie, R., Moradei, O., Woo, S. H., Fournel, M., Yan, P. T., Kalita, Y. A., Trachy-Bourget, M. C., Beaulieu, C., Li, Z., Robert, M. F., MacLeod, A. R., Besterman, J. M. & Delorme, D. (2003). *J. Med. Chem.* **46**, 820-830.
- Bruker (2005). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2007). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837-838.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112-122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148-155.

**supplementary materials**

*Acta Cryst.* (2009). E65, o831 [ doi:10.1107/S1600536809009787 ]

## 5-Benzenesulfonamido-2-chlorobenzoic acid

M. N. Arshad, M. N. Tahir, I. U. Khan, M. Shafiq and H. M. A. Sharif

### Comment

Sulfonamide derivatives have been used as antibacterial agents. Recently these type of derivatives (Bouchain *et al.*, 2003) have been reported as antitumor agents. As part of our ongoing studies of sulfonamide (Arshad, Khan *et al.*, 2009) and thiazine related heterocycles (Arshad *et al.*, 2008), we now report the crystal structure of the title compound, (I), (Fig 1).

The crystal structure of 2-chloro-5-(2-iodobenzenesulfonamido)benzoic acid (Arshad, Tahir *et al.*, 2009), (II), has been reported recently. The title compound differs from (II) as there is no iodine atom on the phenylsulfonyl moiety. Therefore, (II) is the best structure with which the bond distances *etc* can be compared. The title compound consists of dimers due to the carboxylic moiety, forming  $R_2^2(8)$  ring motifs (Bernstein *et al.*, 1995), (Fig 2). These dimers link each other through the N—H $\cdots$ O type of intermolecular H-bonding where the acceptor is the SO<sub>2</sub> moiety (Table 1). The benzene rings A (C1—C6) and B (C8—C13) are oriented at a dihedral angle of 87.07 (6)°. The molecules are stabilized due to intra as well as intermolecular H-bonding and the  $\pi$ -interactions (Table 1).

### Experimental

5-Amino-2-chlorobenzoic acid (1 g, 5.27 mmol) was dissolved in distilled water (10 ml). The pH of the solution was maintained at 8–9 using 1M, Na<sub>2</sub>CO<sub>3</sub> solution. Benzene sulfonyl chloride (0.932 g, 5.27 mmol) was then added to the solution, which was stirred at room temperature until the consumption of all the benzene sulfonyl chloride. During the reaction the pH was again strictly maintained at 8–9 using 1M, Na<sub>2</sub>CO<sub>3</sub>. On completion of the reaction the pH was adjusted 1–2, using 1 N HCl under vigorous stirring. The precipitates obtained were filtered off, washed with distilled water and dried. Colourless prisms of (I) were obtained by recrystallization from methanol.

### Refinement

The H-atoms were positioned geometrically, with O—H = 0.82 Å for hydroxy, N—H = 0.86 Å for amine and C—H = 0.93 Å for aromatic H, and constrained to ride on their parent atoms, with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C}, \text{N}, \text{O})$ .

### Figures




Fig. 1. View of (I) with displacement ellipsoids drawn at the 50% probability level. H-atoms are shown by small circles of arbitrary radii. The dotted lines show the intramolecular H-bonds.




Fig. 2. The partial packing diagram of (I) which shows that the molecules are dimerized and linked to each other.

## 5-Benzenesulfonamido-2-chlorobenzoic acid

### Crystal data

$C_{13}H_{10}ClNO_4S$

$M_r = 311.73$

Monoclinic,  $P2_1/c$

Hall symbol: -P 2ybc

$a = 11.7139$  (4) Å

$b = 5.3957$  (2) Å

$c = 20.7565$  (8) Å

$\beta = 91.483$  (2)°

$V = 1311.47$  (8) Å<sup>3</sup>

$Z = 4$

$F_{000} = 640$

$D_x = 1.579$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation

$\lambda = 0.71073$  Å

Cell parameters from 2234 reflections

$\theta = 2.1$ – $27.0$ °

$\mu = 0.46$  mm<sup>-1</sup>

$T = 296$  K

Prismatic, colorless

$0.24 \times 0.18 \times 0.15$  mm

### Data collection

Bruker Kappa APEXII CCD  
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

Detector resolution: 7.40 pixels mm<sup>-1</sup>

$T = 296$  K

$\omega$  scans

Absorption correction: multi-scan  
(SADABS; Bruker, 2005)

$T_{\min} = 0.939$ ,  $T_{\max} = 0.940$

14693 measured reflections

3269 independent reflections

2513 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.029$

$\theta_{\text{max}} = 28.3$ °

$\theta_{\text{min}} = 2.6$ °

$h = -15 \rightarrow 15$

$k = -7 \rightarrow 7$

$l = -27 \rightarrow 27$

### Refinement

Refinement on  $F^2$

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.039$

$wR(F^2) = 0.108$

$S = 1.02$

3269 reflections

182 parameters

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.048P)^2 + 0.7097P]$$

where  $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\text{max}} < 0.001$

$\Delta\rho_{\text{max}} = 0.37$  e Å<sup>-3</sup>

$\Delta\rho_{\text{min}} = -0.36$  e Å<sup>-3</sup>

Primary atom site location: structure-invariant direct methods Extinction correction: none

*Special details*

**Geometry.** Bond distances, angles *etc.* have been calculated using the rounded fractional coordinates. All su's are estimated from the variances of the (full) variance-covariance matrix. The cell e.s.d.'s are taken into account in the estimation of distances, angles and torsion angles

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Cl1	0.34738 (5)	1.16021 (13)	-0.01036 (3)	0.0671 (2)
S1	0.18337 (4)	0.37106 (8)	0.24025 (2)	0.0317 (1)
O1	-0.00951 (12)	1.2492 (3)	0.05816 (8)	0.0530 (5)
O2	0.12244 (13)	1.3665 (3)	-0.00925 (8)	0.0557 (5)
O3	0.08966 (12)	0.2790 (3)	0.27620 (7)	0.0460 (5)
O4	0.25750 (12)	0.2023 (3)	0.20902 (7)	0.0420 (4)
N1	0.12278 (13)	0.5510 (3)	0.18667 (8)	0.0370 (5)
C1	0.16292 (15)	1.0292 (3)	0.06222 (9)	0.0320 (5)
C2	0.11598 (15)	0.8798 (3)	0.10932 (9)	0.0311 (5)
C3	0.17784 (15)	0.6911 (3)	0.13923 (9)	0.0313 (5)
C4	0.28911 (17)	0.6482 (4)	0.12081 (10)	0.0420 (6)
C5	0.33651 (18)	0.7953 (4)	0.07447 (11)	0.0465 (7)
C6	0.27543 (17)	0.9845 (4)	0.04525 (10)	0.0385 (6)
C7	0.09147 (16)	1.2312 (4)	0.03343 (9)	0.0344 (6)
C8	0.26803 (15)	0.5589 (3)	0.29150 (9)	0.0322 (5)
C9	0.21566 (19)	0.7353 (4)	0.32856 (11)	0.0484 (7)
C10	0.2813 (2)	0.8817 (5)	0.36915 (13)	0.0645 (9)
C11	0.3978 (2)	0.8482 (5)	0.37292 (13)	0.0655 (10)
C12	0.4499 (2)	0.6723 (5)	0.33606 (12)	0.0558 (8)
C13	0.38461 (17)	0.5253 (4)	0.29468 (10)	0.0432 (7)
H1N	0.04957	0.56188	0.18730	0.0444*
H1O	-0.04283	1.36994	0.04279	0.0636*
H2	0.04096	0.90730	0.12110	0.0374*
H4	0.33146	0.52045	0.13970	0.0503*
H5	0.41138	0.76657	0.06261	0.0558*
H9	0.13680	0.75516	0.32615	0.0580*
H10	0.24711	1.00298	0.39397	0.0773*
H11	0.44182	0.94605	0.40084	0.0786*
H12	0.52867	0.65190	0.33884	0.0670*
H13	0.41898	0.40565	0.26941	0.0519*

## supplementary materials

---

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C11	0.0512 (3)	0.0760 (4)	0.0756 (4)	0.0223 (3)	0.0284 (3)	0.0383 (3)
S1	0.0291 (2)	0.0308 (2)	0.0350 (3)	-0.0010 (2)	0.0002 (2)	0.0017 (2)
O1	0.0393 (8)	0.0620 (10)	0.0582 (10)	0.0216 (7)	0.0088 (7)	0.0238 (8)
O2	0.0443 (8)	0.0601 (10)	0.0631 (10)	0.0182 (7)	0.0114 (7)	0.0283 (8)
O3	0.0363 (7)	0.0478 (8)	0.0540 (9)	-0.0098 (6)	0.0038 (6)	0.0115 (7)
O4	0.0442 (8)	0.0354 (7)	0.0462 (8)	0.0065 (6)	-0.0018 (6)	-0.0055 (6)
N1	0.0261 (8)	0.0457 (9)	0.0392 (9)	0.0047 (7)	-0.0004 (6)	0.0085 (7)
C1	0.0325 (9)	0.0349 (9)	0.0284 (9)	0.0056 (7)	-0.0025 (7)	-0.0021 (7)
C2	0.0264 (8)	0.0373 (9)	0.0295 (9)	0.0046 (7)	-0.0020 (7)	-0.0035 (8)
C3	0.0306 (9)	0.0342 (9)	0.0291 (9)	0.0030 (7)	-0.0014 (7)	-0.0019 (7)
C4	0.0353 (10)	0.0448 (11)	0.0460 (12)	0.0138 (9)	0.0047 (8)	0.0102 (9)
C5	0.0342 (10)	0.0539 (13)	0.0519 (13)	0.0159 (9)	0.0114 (9)	0.0118 (10)
C6	0.0356 (10)	0.0430 (11)	0.0372 (10)	0.0060 (8)	0.0059 (8)	0.0050 (9)
C7	0.0322 (9)	0.0382 (10)	0.0326 (10)	0.0053 (8)	-0.0027 (7)	-0.0012 (8)
C8	0.0333 (9)	0.0335 (9)	0.0297 (9)	-0.0038 (7)	0.0016 (7)	0.0022 (7)
C9	0.0438 (12)	0.0503 (12)	0.0512 (13)	0.0002 (10)	0.0056 (10)	-0.0115 (10)
C10	0.0680 (17)	0.0646 (16)	0.0611 (16)	-0.0072 (13)	0.0067 (13)	-0.0284 (13)
C11	0.0665 (17)	0.0720 (18)	0.0577 (15)	-0.0258 (14)	-0.0038 (13)	-0.0170 (13)
C12	0.0362 (11)	0.0722 (16)	0.0587 (14)	-0.0131 (11)	-0.0050 (10)	-0.0017 (12)
C13	0.0351 (10)	0.0513 (12)	0.0433 (12)	-0.0005 (9)	0.0032 (8)	-0.0024 (10)

### Geometric parameters ( $\text{\AA}$ , $^\circ$ )

C11—C6	1.730 (2)	C5—C6	1.378 (3)
S1—O3	1.4321 (15)	C8—C13	1.378 (3)
S1—O4	1.4257 (16)	C8—C9	1.378 (3)
S1—N1	1.6255 (17)	C9—C10	1.375 (3)
S1—C8	1.7572 (18)	C10—C11	1.377 (3)
O1—C7	1.305 (2)	C11—C12	1.372 (4)
O2—C7	1.211 (3)	C12—C13	1.385 (3)
O1—H10	0.8200	C2—H2	0.9300
N1—C3	1.411 (2)	C4—H4	0.9300
N1—H1N	0.8600	C5—H5	0.9300
C1—C6	1.394 (3)	C9—H9	0.9300
C1—C7	1.490 (3)	C10—H10	0.9300
C1—C2	1.391 (3)	C11—H11	0.9300
C2—C3	1.387 (2)	C12—H12	0.9300
C3—C4	1.387 (3)	C13—H13	0.9300
C4—C5	1.375 (3)		
O3—S1—O4	119.97 (9)	S1—C8—C9	118.95 (15)
O3—S1—N1	103.69 (9)	S1—C8—C13	119.76 (14)
O3—S1—C8	108.23 (9)	C9—C8—C13	121.29 (18)
O4—S1—N1	109.31 (9)	C8—C9—C10	119.3 (2)
O4—S1—C8	107.63 (9)	C9—C10—C11	119.8 (2)

N1—S1—C8	107.41 (8)	C10—C11—C12	120.9 (2)
C7—O1—H1O	109.00	C11—C12—C13	119.7 (2)
S1—N1—C3	126.74 (13)	C8—C13—C12	119.03 (19)
C3—N1—H1N	117.00	C1—C2—H2	119.00
S1—N1—H1N	117.00	C3—C2—H2	119.00
C2—C1—C6	118.08 (16)	C3—C4—H4	120.00
C2—C1—C7	118.53 (16)	C5—C4—H4	120.00
C6—C1—C7	123.38 (17)	C4—C5—H5	119.00
C1—C2—C3	121.90 (16)	C6—C5—H5	119.00
N1—C3—C2	117.57 (16)	C8—C9—H9	120.00
N1—C3—C4	123.56 (16)	C10—C9—H9	120.00
C2—C3—C4	118.87 (17)	C9—C10—H10	120.00
C3—C4—C5	119.72 (19)	C11—C10—H10	120.00
C4—C5—C6	121.4 (2)	C10—C11—H11	120.00
C1—C6—C5	120.03 (19)	C12—C11—H11	120.00
C11—C6—C5	116.31 (16)	C11—C12—H12	120.00
C11—C6—C1	123.64 (16)	C13—C12—H12	120.00
O1—C7—O2	122.32 (19)	C8—C13—H13	120.00
O1—C7—C1	113.71 (17)	C12—C13—H13	120.00
O2—C7—C1	123.97 (17)		
O3—S1—N1—C3	179.16 (16)	C2—C1—C7—O2	-177.51 (19)
O4—S1—N1—C3	-51.79 (18)	C6—C1—C7—O1	-176.84 (18)
C8—S1—N1—C3	64.71 (17)	C6—C1—C7—O2	3.7 (3)
O3—S1—C8—C9	-47.55 (18)	C1—C2—C3—N1	179.62 (16)
O3—S1—C8—C13	131.51 (16)	C1—C2—C3—C4	-1.0 (3)
O4—S1—C8—C9	-178.57 (16)	N1—C3—C4—C5	-179.45 (19)
O4—S1—C8—C13	0.48 (18)	C2—C3—C4—C5	1.2 (3)
N1—S1—C8—C9	63.83 (18)	C3—C4—C5—C6	-0.6 (3)
N1—S1—C8—C13	-117.12 (16)	C4—C5—C6—C11	178.09 (17)
S1—N1—C3—C2	-163.64 (14)	C4—C5—C6—C1	-0.4 (3)
S1—N1—C3—C4	17.0 (3)	S1—C8—C9—C10	179.51 (18)
C6—C1—C2—C3	0.1 (3)	C13—C8—C9—C10	0.5 (3)
C7—C1—C2—C3	-178.76 (17)	S1—C8—C13—C12	-179.00 (17)
C2—C1—C6—C11	-177.75 (15)	C9—C8—C13—C12	0.0 (3)
C2—C1—C6—C5	0.6 (3)	C8—C9—C10—C11	-0.9 (4)
C7—C1—C6—C11	1.1 (3)	C9—C10—C11—C12	0.8 (4)
C7—C1—C6—C5	179.40 (19)	C10—C11—C12—C13	-0.3 (4)
C2—C1—C7—O1	2.0 (3)	C11—C12—C13—C8	-0.1 (3)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H $\cdots$ <i>A</i>	<i>D</i> —H	H $\cdots$ <i>A</i>	<i>D</i> $\cdots$ <i>A</i>	<i>D</i> —H $\cdots$ <i>A</i>
O1—H1O $\cdots$ O2 <sup>i</sup>	0.82	1.83	2.648 (2)	178
N1—H1N $\cdots$ O3 <sup>ii</sup>	0.86	2.16	2.898 (2)	144
C4—H4 $\cdots$ O4	0.93	2.41	3.052 (3)	126
C6—C11 $\cdots$ CgB <sup>iii</sup>	1.730 (2)	3.8131 (12)	4.605 (2)	106.16 (8)
S1—O4 $\cdots$ CgA <sup>iv</sup>	1.4257 (16)	3.1405 (17)	4.2532 (9)	133.77 (8)

Symmetry codes: (i)  $-x, -y+3, -z$ ; (ii)  $-x, y+1/2, -z+1/2$ ; (iii)  $x, -y+3/2, z-1/2$ ; (iv)  $x, y-1, z$ .

Fig. 1






Fig. 2

