

N'-(5-Chloro-2-hydroxybenzylidene)-3,4,5-trihydroxybenzohydrazide dihydrate

Abeer A. Abdul Alhadi, Hapipah Mohd. Ali and Seik Weng Ng*

Department of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia
Correspondence e-mail: seikweng@um.edu.my


Received 23 March 2009; accepted 24 March 2009

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.007$ Å; R factor = 0.071; wR factor = 0.181; data-to-parameter ratio = 11.9.

The benzohydrazide molecule in the title dihydrate, $\text{C}_{14}\text{H}_{11}\text{ClN}_2\text{O}_5 \cdot 2\text{H}_2\text{O}$, is non-planar, with the two aromatic rings at either side of the $-\text{C}(=\text{O})-\text{NH}-\text{N}=\text{CH}-$ unit forming a dihedral angle of $29.7(2)^\circ$. The benzohydrazide molecule is linked to the water molecules by $\text{O}-\text{H}\cdots\text{O}$ and $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds, with other $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds leading to a layer structure.

Related literature

For the the parent *N'*-(2-hydroxybenzylidene)benzohydrazide, see: Lyubchova *et al.* (1995). For other *N'*-(2-hydroxy-5-nitrobenzylidene)benzohydrazides, see: Ali *et al.* (2005); Lyubchova *et al.* (1995); Xu & Liu (2006).



Experimental

Crystal data

$\text{C}_{14}\text{H}_{11}\text{ClN}_2\text{O}_5 \cdot 2\text{H}_2\text{O}$
 $M_r = 358.73$
Monoclinic, $P2_1/c$

$a = 30.5627(12)$ Å
 $b = 3.7539(2)$ Å
 $c = 12.8882(5)$ Å

$\beta = 90.450(3)^\circ$
 $V = 1478.61(11)$ Å³
 $Z = 4$
Mo $K\alpha$ radiation

$\mu = 0.30$ mm⁻¹
 $T = 100$ K
 $0.36 \times 0.04 \times 0.04$ mm

Data collection

Bruker SMART APEX diffractometer
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)
 $T_{\min} = 0.899$, $T_{\max} = 0.988$

10104 measured reflections
2623 independent reflections
1801 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.097$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.071$
 $wR(F^2) = 0.181$
 $S = 1.07$
2623 reflections

221 parameters
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.38$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.49$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{O1}-\text{H1}\cdots\text{N1}$	0.84	1.89	2.631 (5)	146
$\text{O3}-\text{H3}\cdots\text{O1w}^i$	0.84	1.96	2.737 (6)	153
$\text{O4}-\text{H4}\cdots\text{O1w}^{ii}$	0.84	1.80	2.599 (7)	158
$\text{O5}-\text{H5}\cdots\text{O2}^{ii}$	0.84	1.93	2.765 (5)	171
$\text{O1w}-\text{H11}\cdots\text{O3}$	0.83	2.28	2.969 (6)	140
$\text{O1w}-\text{H12}\cdots\text{O4}^{iii}$	0.84	2.07	2.900 (7)	170
$\text{O2w}-\text{H21}\cdots\text{O1}$	0.84	2.15	2.946 (5)	157
$\text{O2w}-\text{H22}\cdots\text{O2}^{iv}$	0.84	1.97	2.808 (5)	172
$\text{N2}-\text{H2}\cdots\text{O2w}^{ii}$	0.88	2.03	2.882 (5)	162

Symmetry codes: (i) $-x + 2, y + \frac{1}{2}, -z + \frac{3}{2}$; (ii) $x, -y + \frac{3}{2}, z - \frac{1}{2}$; (iii) $x, -y + \frac{5}{2}, z + \frac{1}{2}$; (iv) $x, y - 1, z$.

Data collection: APEX2 (Bruker, 2008); cell refinement: SAINT (Bruker, 2008); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: X-SEED (Barbour, 2001); software used to prepare material for publication: publCIF (Westrip, 2009).

We thank the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: TK2403).

References

- Ali, H. M., Puvaneswary, S. & Ng, S. W. (2005). *Acta Cryst.* **E61**, o3464–o3465.
Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
Bruker (2008). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
Lyubchova, A., Cossé-Barbi, A., Doucet, J. P., Robert, F., Souron, J.-P. & Quarton, M. (1995). *Acta Cryst.* **C51**, 1893–1895.
Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Westrip, S. P. (2009). publCIF. In preparation.
Xu, H.-M. & Liu, S.-X. (2006). *Acta Cryst.* **E62**, o3026–o3027.

supplementary materials

Acta Cryst. (2009). E65, o910 [doi:10.1107/S1600536809010812]

***N'*-(5-Chloro-2-hydroxybenzylidene)-3,4,5-trihydroxybenzohydrazide dihydrate**

A. A. Abdul Alhadi, H. M. Ali and S. W. Ng

Comment

(type here to add)

Experimental

5-Chloro-2-hydroxybenzaldehyde (0.31 g, 2 mmol) and 3,4,5-trihydroxybenzoylhydrazide (0.36 g, 2 mmol) were heated in ethanol (50 ml) for several hours. The solvent was removed and the product recrystallized from DMSO.

Refinement

Carbon-bound H-atoms were placed in calculated positions [C—H 0.95 Å, $U(H) = 1.2U(C)$], and were included in the refinement in the riding model approximation. The amino (0.88 Å) and hydroxy H-atoms (0.84 Å) were similarly generated with $U_{iso} = 1.2U_{eq}(\text{carrier atom})$ for N-H and $U_{iso} = 1.5U_{eq}(\text{carrier atom})$ for O-H. The water H-atoms were placed in chemically sensible positions on the basis of possible hydrogen bonds, but were not refined; $U_{iso} = 1.5U_{eq}(O)$.

Figures




Fig. 1. Thermal ellipsoid plot (Barbour, 2001) of $C_{14}H_{11}ClN_2O_5 \cdot 2H_2O$ at the 70% probability level; hydrogen atoms are drawn as spheres of arbitrary radius.

***N'*-(5-Chloro-2-hydroxybenzylidene)-3,4,5-trihydroxybenzohydrazide dihydrate**

Crystal data

$C_{14}H_{11}ClN_2O_5 \cdot 2H_2O$

$M_r = 358.73$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

$a = 30.5627(12) \text{ \AA}$

$b = 3.7539(2) \text{ \AA}$

$c = 12.8882(5) \text{ \AA}$

$\beta = 90.450(3)^\circ$

$V = 1478.61(11) \text{ \AA}^3$

$Z = 4$

$F_{000} = 744$

$D_x = 1.611 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation

$\lambda = 0.71073 \text{ \AA}$

Cell parameters from 1505 reflections

$\theta = 2.7\text{--}24.3^\circ$

$\mu = 0.30 \text{ mm}^{-1}$

$T = 100 \text{ K}$

Prism, yellow

$0.36 \times 0.04 \times 0.04 \text{ mm}$

supplementary materials

Data collection

Bruker SMART APEX diffractometer	2623 independent reflections
Radiation source: fine-focus sealed tube	1801 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.097$
$T = 100$ K	$\theta_{\text{max}} = 25.0^\circ$
ω scans	$\theta_{\text{min}} = 0.7^\circ$
Absorption correction: Multi-scan (SADABS; Sheldrick, 1996)	$h = -36 \rightarrow 36$
$T_{\text{min}} = 0.899$, $T_{\text{max}} = 0.988$	$k = -4 \rightarrow 4$
10104 measured reflections	$l = -15 \rightarrow 14$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.071$	H-atom parameters constrained
$wR(F^2) = 0.181$	$w = 1/[\sigma^2(F_o^2) + (0.0537P)^2 + 5.5995P]$
$S = 1.07$	where $P = (F_o^2 + 2F_c^2)/3$
2623 reflections	$(\Delta/\sigma)_{\text{max}} = 0.001$
221 parameters	$\Delta\rho_{\text{max}} = 0.38 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	$\Delta\rho_{\text{min}} = -0.49 \text{ e } \text{\AA}^{-3}$
	Extinction correction: none

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C11	0.53161 (4)	0.3168 (4)	0.59750 (10)	0.0244 (3)
O1	0.67086 (11)	0.6471 (10)	0.8994 (2)	0.0243 (9)
H1	0.6933	0.7236	0.8694	0.036*
O2	0.79657 (11)	0.9733 (10)	0.8420 (2)	0.0232 (9)
O3	0.94472 (12)	1.3696 (12)	0.7043 (3)	0.0349 (10)
H3	0.9663	1.3279	0.6661	0.052*
O4	0.94729 (12)	1.0933 (13)	0.5106 (3)	0.0395 (12)
H4	0.9481	0.9332	0.4651	0.059*
O5	0.87480 (11)	0.7706 (11)	0.4244 (2)	0.0245 (9)
H5	0.8498	0.7191	0.4013	0.037*
O1w	0.97110 (13)	0.8480 (13)	0.8663 (4)	0.0542 (14)
H11	0.9676	0.9037	0.8040	0.081*
H12	0.9666	1.0253	0.9042	0.081*
O2w	0.73755 (11)	0.1847 (10)	0.9961 (2)	0.0239 (8)
H21	0.7141	0.2633	0.9700	0.036*
H22	0.7554	0.1438	0.9482	0.036*
N1	0.71793 (13)	0.8677 (12)	0.7419 (3)	0.0194 (10)

N2	0.75565 (13)	0.9864 (12)	0.6939 (3)	0.0187 (9)
H2	0.7545	1.0529	0.6284	0.022*
C1	0.63896 (16)	0.5781 (13)	0.8275 (4)	0.0179 (11)
C2	0.60017 (17)	0.4266 (14)	0.8603 (4)	0.0209 (12)
H2A	0.5965	0.3740	0.9318	0.025*
C3	0.56699 (17)	0.3513 (14)	0.7916 (4)	0.0233 (12)
H3A	0.5403	0.2508	0.8151	0.028*
C4	0.57295 (16)	0.4243 (13)	0.6865 (4)	0.0189 (11)
C5	0.61125 (16)	0.5725 (13)	0.6514 (4)	0.0193 (11)
H5A	0.6148	0.6213	0.5797	0.023*
C6	0.64514 (16)	0.6516 (14)	0.7219 (4)	0.0179 (11)
C7	0.68550 (15)	0.7976 (13)	0.6815 (4)	0.0172 (11)
H7	0.6880	0.8421	0.6092	0.021*
C9	0.79416 (16)	1.0018 (14)	0.7457 (4)	0.0181 (11)
C10	0.83308 (16)	1.0434 (14)	0.6800 (4)	0.0183 (11)
C11	0.87050 (16)	1.1993 (15)	0.7207 (4)	0.0214 (12)
H11A	0.8704	1.2939	0.7890	0.026*
C12	0.90791 (16)	1.2173 (16)	0.6620 (4)	0.0266 (13)
C13	0.90870 (16)	1.0734 (15)	0.5623 (4)	0.0241 (13)
C14	0.87108 (16)	0.9157 (15)	0.5212 (4)	0.0219 (12)
C15	0.83293 (16)	0.9046 (14)	0.5784 (4)	0.0189 (11)
H15	0.8070	0.8046	0.5496	0.023*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C11	0.0204 (7)	0.0272 (7)	0.0257 (7)	-0.0018 (6)	-0.0025 (5)	-0.0039 (6)
O1	0.026 (2)	0.031 (2)	0.0151 (17)	-0.0024 (18)	-0.0023 (15)	-0.0001 (17)
O2	0.0211 (19)	0.036 (2)	0.0122 (18)	0.0034 (17)	-0.0014 (14)	-0.0031 (16)
O3	0.022 (2)	0.049 (3)	0.034 (2)	-0.014 (2)	-0.0055 (17)	0.006 (2)
O4	0.021 (2)	0.070 (4)	0.028 (2)	-0.009 (2)	0.0042 (17)	0.011 (2)
O5	0.0199 (18)	0.042 (2)	0.0118 (17)	-0.0017 (18)	0.0023 (14)	-0.0004 (17)
O1w	0.027 (2)	0.056 (3)	0.079 (3)	0.013 (2)	0.012 (2)	0.017 (3)
O2w	0.0223 (18)	0.035 (2)	0.0139 (17)	0.0042 (17)	-0.0002 (14)	-0.0012 (17)
N1	0.015 (2)	0.024 (2)	0.019 (2)	0.0020 (18)	-0.0001 (17)	0.0006 (19)
N2	0.018 (2)	0.026 (2)	0.0121 (19)	-0.0012 (19)	0.0017 (17)	-0.0016 (19)
C1	0.023 (3)	0.016 (3)	0.014 (2)	0.002 (2)	-0.001 (2)	0.002 (2)
C2	0.027 (3)	0.022 (3)	0.013 (2)	0.002 (2)	0.005 (2)	0.000 (2)
C3	0.022 (3)	0.023 (3)	0.026 (3)	0.000 (2)	0.009 (2)	0.000 (2)
C4	0.017 (3)	0.017 (3)	0.023 (3)	0.001 (2)	0.000 (2)	-0.003 (2)
C5	0.025 (3)	0.020 (3)	0.013 (2)	0.004 (2)	0.001 (2)	0.000 (2)
C6	0.022 (3)	0.018 (3)	0.014 (2)	0.002 (2)	0.002 (2)	-0.001 (2)
C7	0.024 (3)	0.016 (3)	0.011 (2)	0.005 (2)	0.003 (2)	0.001 (2)
C9	0.021 (3)	0.018 (3)	0.015 (3)	0.001 (2)	-0.001 (2)	0.002 (2)
C10	0.019 (3)	0.023 (3)	0.013 (2)	0.003 (2)	-0.003 (2)	0.006 (2)
C11	0.022 (3)	0.024 (3)	0.018 (3)	0.001 (2)	-0.004 (2)	0.001 (2)
C12	0.019 (3)	0.036 (3)	0.025 (3)	-0.007 (2)	-0.006 (2)	0.013 (3)
C13	0.017 (3)	0.035 (3)	0.020 (3)	-0.001 (2)	0.000 (2)	0.009 (2)

supplementary materials

C14	0.024 (3)	0.029 (3)	0.013 (2)	0.000 (2)	-0.002 (2)	0.007 (2)
C15	0.018 (3)	0.022 (3)	0.017 (2)	-0.001 (2)	-0.002 (2)	0.002 (2)

Geometric parameters (Å, °)

C11—C4	1.747 (5)	C1—C6	1.403 (6)
O1—C1	1.365 (6)	C2—C3	1.371 (7)
O1—H1	0.8400	C2—H2A	0.9500
O2—C9	1.247 (6)	C3—C4	1.395 (7)
O3—C12	1.371 (6)	C3—H3A	0.9500
O3—H3	0.8400	C4—C5	1.376 (7)
O4—C13	1.361 (6)	C5—C6	1.403 (7)
O4—H4	0.8400	C5—H5A	0.9500
O5—C14	1.366 (6)	C6—C7	1.450 (7)
O5—H5	0.8400	C7—H7	0.9500
O1w—H11	0.8347	C9—C10	1.474 (7)
O1w—H12	0.8378	C10—C11	1.384 (7)
O2w—H21	0.8421	C10—C15	1.408 (7)
O2w—H22	0.8400	C11—C12	1.377 (7)
N1—C7	1.283 (6)	C11—H11A	0.9500
N1—N2	1.386 (6)	C12—C13	1.395 (8)
N2—C9	1.350 (6)	C13—C14	1.394 (7)
N2—H2	0.8800	C14—C15	1.385 (7)
C1—C2	1.384 (7)	C15—H15	0.9500
C1—O1—H1	109.5	C1—C6—C7	122.9 (4)
C12—O3—H3	109.5	C5—C6—C7	118.3 (4)
C13—O4—H4	109.5	N1—C7—C6	121.0 (4)
C14—O5—H5	109.5	N1—C7—H7	119.5
H11—O1w—H12	110.0	C6—C7—H7	119.5
H21—O2w—H22	109.0	O2—C9—N2	122.2 (4)
C7—N1—N2	115.9 (4)	O2—C9—C10	122.6 (4)
C9—N2—N1	121.2 (4)	N2—C9—C10	115.1 (4)
C9—N2—H2	119.4	C11—C10—C15	120.3 (5)
N1—N2—H2	119.4	C11—C10—C9	119.7 (4)
O1—C1—C2	118.6 (4)	C15—C10—C9	119.9 (4)
O1—C1—C6	121.4 (4)	C12—C11—C10	120.0 (5)
C2—C1—C6	120.0 (4)	C12—C11—H11A	120.0
C3—C2—C1	121.2 (5)	C10—C11—H11A	120.0
C3—C2—H2A	119.4	O3—C12—C11	119.0 (5)
C1—C2—H2A	119.4	O3—C12—C13	120.5 (5)
C2—C3—C4	119.0 (5)	C11—C12—C13	120.5 (5)
C2—C3—H3A	120.5	O4—C13—C14	123.5 (5)
C4—C3—H3A	120.5	O4—C13—C12	116.8 (5)
C5—C4—C3	121.1 (5)	C14—C13—C12	119.7 (5)
C5—C4—C11	119.4 (4)	O5—C14—C15	123.4 (4)
C3—C4—C11	119.5 (4)	O5—C14—C13	116.2 (4)
C4—C5—C6	119.9 (4)	C15—C14—C13	120.3 (5)
C4—C5—H5A	120.0	C14—C15—C10	119.2 (4)
C6—C5—H5A	120.0	C14—C15—H15	120.4

C1—C6—C5	118.8 (5)	C10—C15—H15	120.4
C7—N1—N2—C9	-166.9 (5)	N2—C9—C10—C11	154.5 (5)
O1—C1—C2—C3	-180.0 (5)	O2—C9—C10—C15	148.7 (5)
C6—C1—C2—C3	-1.1 (8)	N2—C9—C10—C15	-29.5 (7)
C1—C2—C3—C4	1.0 (8)	C15—C10—C11—C12	-0.4 (8)
C2—C3—C4—C5	-0.6 (8)	C9—C10—C11—C12	175.6 (5)
C2—C3—C4—C11	178.0 (4)	C10—C11—C12—O3	-179.3 (5)
C3—C4—C5—C6	0.2 (8)	C10—C11—C12—C13	-1.2 (8)
C11—C4—C5—C6	-178.4 (4)	O3—C12—C13—O4	-0.1 (8)
O1—C1—C6—C5	179.6 (5)	C11—C12—C13—O4	-178.2 (5)
C2—C1—C6—C5	0.7 (7)	O3—C12—C13—C14	179.2 (5)
O1—C1—C6—C7	1.2 (8)	C11—C12—C13—C14	1.1 (8)
C2—C1—C6—C7	-177.7 (5)	O4—C13—C14—O5	1.3 (8)
C4—C5—C6—C1	-0.3 (7)	C12—C13—C14—O5	-178.0 (5)
C4—C5—C6—C7	178.2 (5)	O4—C13—C14—C15	179.9 (5)
N2—N1—C7—C6	176.3 (4)	C12—C13—C14—C15	0.6 (8)
C1—C6—C7—N1	-0.1 (8)	O5—C14—C15—C10	176.3 (5)
C5—C6—C7—N1	-178.5 (5)	C13—C14—C15—C10	-2.2 (8)
N1—N2—C9—O2	-13.1 (8)	C11—C10—C15—C14	2.1 (7)
N1—N2—C9—C10	165.1 (4)	C9—C10—C15—C14	-173.9 (5)
O2—C9—C10—C11	-27.4 (8)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
O1—H1...N1	0.84	1.89	2.631 (5)	146
O3—H3...O1w ⁱ	0.84	1.96	2.737 (6)	153
O4—H4...O1w ⁱⁱ	0.84	1.80	2.599 (7)	158
O5—H5...O2 ⁱⁱ	0.84	1.93	2.765 (5)	171
O1w—H11...O3	0.83	2.28	2.969 (6)	140
O1w—H12...O4 ⁱⁱⁱ	0.84	2.07	2.900 (7)	170
O2w—H21...O1	0.84	2.15	2.946 (5)	157
O2w—H22...O2 ^{iv}	0.84	1.97	2.808 (5)	172
N2—H2...O2w ⁱⁱ	0.88	2.03	2.882 (5)	162

Symmetry codes: (i) $-x+2, y+1/2, -z+3/2$; (ii) $x, -y+3/2, z-1/2$; (iii) $x, -y+5/2, z+1/2$; (iv) $x, y-1, z$.

Fig. 1

