

Bis(4-aminopyridinium) tetrachlorido-cobaltate(II)

Samuel Robinson Jebas,^{a*} A. Sinthiya,^b B. Ravindran Durai Nayagam,^c Dieter Schollmeyer^d and S. Alfred Cecil Raj^e

^aDepartment of Physics, Karunya University, Karunya Nagar, Coimbatore 641 114, India, ^bDepartment of Electronics, St. Josephs College, Tiruchirappalli 620 002, India, ^cDepartment of Chemistry, Popes College, Sawyerupuram 628 251, Tamilnadu, India, ^dInstitut für Organische Chemie, Universität Mainz, Duesbergweg 10-14, 55099 Mainz, Germany, and ^eDepartment of Physics, St. Josephs College, Tiruchirappalli 620 002, India

Correspondence e-mail: jebas2@gmail.com


Received 5 April 2009; accepted 7 April 2009

Key indicators: single-crystal X-ray study; $T = 173\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$; R factor = 0.021; wR factor = 0.052; data-to-parameter ratio = 22.5.

In the title compound, $(\text{C}_5\text{H}_7\text{N}_2)_2[\text{CoCl}_4]$, the cobalt(II) ion is coordinated by four chloride ions in a slightly distorted tetrahedral geometry. The crystal packing is stabilized by intermolecular N—H···Cl hydrogen bonding, forming a three-dimensional network. The crystal was a non-merohedral twin emulating tetragonal symmetry, but being in fact orthorhombic.

Related literature

For the biological activity of 4-aminopyridine, see: Judge & Bever (2006); Schwid *et al.* (1997); Strupp *et al.* (2004). For related structures, see: Anderson *et al.* (2005); Chao & Schempp (1977); Jebas *et al.* (2006); Zhang *et al.* (2005). For bond-length data, see: Anderson *et al.* (2005).



Experimental

Crystal data

$(\text{C}_5\text{H}_7\text{N}_2)_2[\text{CoCl}_4]$
 $M_r = 390.98$

Orthorhombic, $P_{\bar{2}}12_12_1$
 $a = 15.0051 (12)\text{ \AA}$

$b = 14.9751 (12)\text{ \AA}$
 $c = 7.1723 (6)\text{ \AA}$
 $V = 1611.6 (2)\text{ \AA}^3$
 $Z = 4$

Mo $K\alpha$ radiation
 $\mu = 1.72\text{ mm}^{-1}$
 $T = 173\text{ K}$
 $0.25 \times 0.22 \times 0.17\text{ mm}$

Data collection

Bruker APEXII SMART CCD diffractometer
Absorption correction: multi-scan (SADABS; Bruker, 2008)
 $T_{\min} = 0.650$, $T_{\max} = 0.746$

45299 measured reflections
3884 independent reflections
3802 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.035$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.021$
 $wR(F^2) = 0.052$
 $S = 1.02$
3884 reflections
173 parameters
H-atom parameters constrained

$\Delta\rho_{\text{max}} = 0.25\text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.13\text{ e \AA}^{-3}$
Absolute structure: Flack (1983),
1654 Friedel pairs
Flack parameter: -0.12 (2)

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N4—H4···Cl2 ⁱ	0.88	2.94	3.563 (3)	130
N4—H4···Cl3 ⁱ	0.88	2.67	3.335 (3)	134
N7—H7A···Cl2 ⁱⁱ	0.84	2.50	3.338 (2)	175
N7—H7B···Cl1	0.90	2.53	3.387 (2)	158
N11—H11···Cl1 ⁱⁱⁱ	0.87	2.52	3.272 (3)	144
N14—H14B···Cl2 ^{iv}	0.84	2.64	3.394 (3)	149
N14—H14A···Cl4	0.90	2.42	3.303 (2)	169

Symmetry codes: (i) $-x + \frac{1}{2}, -y, z - \frac{1}{2}$; (ii) $x - \frac{1}{2}, -y + \frac{1}{2}, -z + 1$; (iii) $x - \frac{1}{2}, -y + \frac{1}{2}, -z$; (iv) $-x + 1, y + \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: APEX2 (Bruker, 2008); cell refinement: SAINT (Bruker, 2008); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL and PLATON (Spek, 2009).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT2925).

References

- Anderson, F. P., Gallagher, J. F., Kenny, P. T. M. & Lough, A. J. (2005). *Acta Cryst. E61*, o1350–o1353.
- Bruker (2008). APEX2, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
- Chao, M. & Schempp, E. (1977). *Acta Cryst. B33*, 1557–1564.
- Flack, H. D. (1983). *Acta Cryst. A39*, 876–881.
- Jebas, S. R., Balasubramanian, T. & Light, M. E. (2006). *Acta Cryst. E62*, m1818–m1819.
- Judge, S. & Bever, C. (2006). *Pharmacol. Ther.* **111**, 224–259.
- Schwid, S. B., Petrie, M. D., McDermott, M. P., Tierney, D. S., Mason, D. H. & Goodman, A. D. (1997). *Neurology*, **48**, 817–821.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D65*, 148–155.
- Strupp, M., Kalla, R., Dichgans, M., Fraitinger, T., Glasauer, S. & Brandt, T. (2004). *Neurology*, **62**, 1623–1625.
- Zhang, H., Fang, L. & Yuan, R. (2005). *Acta Cryst. E61*, m677–m678.

supplementary materials

Acta Cryst. (2009). E65, m521 [doi:10.1107/S1600536809013270]

Bis(4-aminopyridinium) tetrachloridocobaltate(II)

S. R. Jebas, A. Sinthiya, B. Ravindran Durai Nayagam, D. Schollmeyer and S. A. C. Raj

Comment

4-aminopyridine (Fampridine) is used clinically in Lambert-Eaton myasthenic syndrome and multiple sclerosis because by blocking potassium channels it prolongs action potentials thereby increasing transmitter release at the neuromuscular junction (Judge & Bever, 2006; Schwid *et al.*, 1997; Strupp *et al.*, 2004). The structure of 4-aminopyridine has been reported (Chao & Schempp, 1977). Redetermination of the structure of 4-aminopyridine has been reported (Anderson *et al.*, 2005). As a part of our investigation of the reactions of the 4-aminopyridine with metals, we report here the crystal structure of the title compound (I).

The asymmetric unit of (I), consists of two molecules of 4-aminopyridinium cation and a $[\text{CoCl}_4]^{2-}$ anion. The bond lengths and bond angles of the 4-aminopyridinium are comparable with the values reported earlier for the 4-aminopyridine in its uncomplexed form (Anderson *et al.*, 2005; Chao & Schempp, 1977). Protonation of the atoms N4 and N11 of the 4-aminopyridine leads to the widening of C3—N4—C5 and C10—N11—C12 angles in the pyridine ring to 121.3 (7) $^{\circ}$ and 120.4 (7) $^{\circ}$, compared to 115.25 (13) $^{\circ}$ in 4-aminopyridine (Anderson *et al.*, 2005). The 4-aminopyridine ring is essentially planar with the maximum deviation from planarity of 0.014 (3) Å for the atoms C8 and N11 respectively.

The anion exhibits distorted tetrahedral geometry, with the Co^{II} ion is surrounded by four Cl atoms, with Cl—Co—Cl angles ranging from 106.19 (11)—115.63 (8) Å. The mean Co—Cl bond length, 2.2707 (2) Å, is close to those observed in similar complex (Jebas *et al.*, 2006; Zhang *et al.*, 2005).

The crystal packing (Fig. 2) is consolidated by intermolecular N—H···Cl hydrogen bonding to form a three dimensional network.

Experimental

4-aminopyridine (0.094 g, 1 mmol) and CoCl_2 (0.169 g, 1 mmol) in ethanol (10 ml each) and the solution was stirred well for 3 h. Blue crystals of (I) were obtained by slow evaporation of the solution over a period of one month.

Refinement

The crystals of (I) crystallized with nearly tetragonal lattice parameters. It was not possible to solve and refine the structure in any tetragonal space group but it could be easily solved and refined in orthorhombic space group P2₁2₁2₁. PLATON and the intensity statistic indicate twinning. Applying the twin instruction TWIN 0 1 0 1 0 0 0 0 -1 with a BASF of 0.340 (1) the R1 value drops to 0.021 (0.095 without TWIN instruction). The nonstandard setting for the orthorhombic cell was kept to simplify the twin matrix.

All the hydrogen atoms were fixed on the calculated positions and allowed to ride on their parent atoms with the C—H = 0.95 Å (aromatic); N—H = 0.84–0.89 Å with $U_{\text{iso}}(\text{C})$ in the range of 1.2 $U_{\text{eq}}(\text{C})$ –1.5 $U_{\text{eq}}(\text{N})$.

supplementary materials

Figures




Fig. 1. The molecular structure of the title compound, showing 50% probability displacement ellipsoids and the atom numbering scheme. Intramolecular hydrogen bondings are shown as dashed lines.




Fig. 2. The crystal packing of the title compound, viewed down the α axis, showing the three dimensional network.

Bis(4-aminopyridinium) tetrachloridocobalt(II)

Crystal data

(C ₅ H ₇ N ₂) ₂ [CoCl ₄]	$F_{000} = 788$
$M_r = 390.98$	$D_x = 1.611 \text{ Mg m}^{-3}$
Orthorhombic, $P2_12_12_1$	Mo $K\alpha$ radiation
Hall symbol: P 2ac 2ab	$\lambda = 0.71073 \text{ \AA}$
$a = 15.0051 (12) \text{ \AA}$	Cell parameters from 9192 reflections
$b = 14.9751 (12) \text{ \AA}$	$\theta = 2.7\text{--}27.8^\circ$
$c = 7.1723 (6) \text{ \AA}$	$\mu = 1.72 \text{ mm}^{-1}$
$V = 1611.6 (2) \text{ \AA}^3$	$T = 173 \text{ K}$
$Z = 4$	Block, blue
	$0.25 \times 0.22 \times 0.17 \text{ mm}$

Data collection

Bruker APEXII SMART CCD diffractometer	3802 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.035$
$T = 173 \text{ K}$	$\theta_{\max} = 28.0^\circ$
CCD scan	$\theta_{\min} = 1.4^\circ$
Absorption correction: multi-scan (SADABS; Bruker, 2008)	$h = -19 \rightarrow 19$
$T_{\min} = 0.650, T_{\max} = 0.746$	$k = -19 \rightarrow 19$

45299 measured reflections

$l = -9 \rightarrow 9$

3884 independent reflections

Refinement

Refinement on F^2

Hydrogen site location: inferred from neighbouring sites

Least-squares matrix: full

H-atom parameters constrained

$R[F^2 > 2\sigma(F^2)] = 0.021$

$$w = 1/[\sigma^2(F_o^2) + (0.0281P)^2 + 0.361P]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$wR(F^2) = 0.052$

$$(\Delta/\sigma)_{\max} < 0.001$$

$S = 1.02$

$$\Delta\rho_{\max} = 0.25 \text{ e \AA}^{-3}$$

3884 reflections

$$\Delta\rho_{\min} = -0.12 \text{ e \AA}^{-3}$$

173 parameters

Extinction correction: none

Primary atom site location: structure-invariant direct methods

Absolute structure: Flack (1983), 1654 Friedel pairs

Secondary atom site location: difference Fourier map Flack parameter: -0.12 (2)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger. Structure was refined as tetragonal twin with basf=0.34063

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Co1	0.49922 (2)	0.24481 (2)	0.37955 (6)	0.02580 (7)
Cl1	0.45208 (4)	0.12712 (5)	0.20160 (11)	0.03485 (16)
Cl2	0.59477 (5)	0.18722 (5)	0.59594 (11)	0.03872 (17)
Cl3	0.38406 (6)	0.30281 (5)	0.54058 (11)	0.04194 (19)
Cl4	0.56931 (5)	0.33990 (5)	0.18373 (14)	0.04232 (17)
C1	0.19394 (17)	0.06484 (18)	0.2826 (4)	0.0289 (5)
C2	0.2459 (2)	-0.0132 (2)	0.2552 (4)	0.0337 (6)
H2	0.3091	-0.0095	0.2540	0.040*
C3	0.2050 (2)	-0.0935 (2)	0.2307 (4)	0.0379 (7)
H3	0.2400	-0.1455	0.2116	0.045*
N4	0.11624 (18)	-0.10010 (17)	0.2331 (3)	0.0413 (6)
H4	0.0873	-0.1508	0.2248	0.050*
C5	0.0644 (2)	-0.0280 (2)	0.2604 (4)	0.0403 (7)
H5	0.0015	-0.0348	0.2631	0.048*
C6	0.10048 (18)	0.0545 (2)	0.2843 (4)	0.0342 (6)

supplementary materials

H6	0.0630	0.1049	0.3021	0.041*
N7	0.23233 (15)	0.14446 (15)	0.3030 (4)	0.0382 (5)
H7A	0.2002	0.1883	0.3330	0.057*
H7B	0.2909	0.1553	0.2895	0.057*
C8	0.32310 (18)	0.44022 (18)	0.0283 (4)	0.0289 (5)
C9	0.3084 (2)	0.34727 (18)	0.0205 (4)	0.0326 (6)
H9	0.3562	0.3066	0.0389	0.039*
C10	0.2245 (2)	0.3171 (2)	-0.0139 (4)	0.0398 (7)
H10	0.2142	0.2546	-0.0204	0.048*
N11	0.15584 (17)	0.37337 (19)	-0.0389 (3)	0.0418 (6)
H11	0.1000	0.3568	-0.0377	0.050*
C12	0.1679 (2)	0.4632 (2)	-0.0254 (4)	0.0382 (7)
H12	0.1183	0.5021	-0.0385	0.046*
C13	0.24960 (17)	0.4974 (3)	0.0064 (4)	0.0331 (6)
H13	0.2575	0.5602	0.0140	0.040*
N14	0.40431 (15)	0.47315 (16)	0.0593 (3)	0.0375 (5)
H14A	0.4522	0.4383	0.0767	0.056*
H14B	0.4130	0.5288	0.0656	0.056*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Co1	0.02267 (19)	0.0223 (2)	0.03247 (13)	0.00081 (16)	-0.00149 (13)	-0.00034 (12)
Cl1	0.0261 (3)	0.0370 (4)	0.0415 (4)	-0.0058 (2)	0.0005 (3)	-0.0129 (3)
Cl2	0.0389 (4)	0.0279 (3)	0.0493 (4)	0.0002 (2)	-0.0190 (3)	0.0004 (3)
Cl3	0.0407 (4)	0.0315 (4)	0.0536 (4)	0.0119 (3)	0.0134 (3)	0.0012 (3)
Cl4	0.0296 (3)	0.0346 (4)	0.0628 (4)	-0.0003 (3)	0.0073 (3)	0.0151 (3)
C1	0.0278 (13)	0.0327 (14)	0.0263 (13)	0.0030 (10)	-0.0015 (11)	0.0023 (11)
C2	0.0296 (15)	0.0333 (15)	0.0383 (15)	0.0032 (11)	0.0010 (11)	0.0018 (13)
C3	0.0486 (17)	0.0302 (14)	0.0349 (15)	0.0031 (12)	-0.0027 (13)	-0.0019 (11)
N4	0.0536 (15)	0.0364 (13)	0.0338 (12)	-0.0155 (11)	-0.0055 (11)	0.0029 (10)
C5	0.0334 (16)	0.0515 (17)	0.0359 (15)	-0.0080 (13)	-0.0017 (12)	0.0069 (12)
C6	0.0289 (14)	0.0408 (16)	0.0329 (14)	0.0033 (11)	-0.0021 (12)	0.0026 (12)
N7	0.0303 (11)	0.0273 (12)	0.0570 (15)	0.0018 (8)	0.0002 (11)	0.0017 (11)
C8	0.0279 (13)	0.0328 (13)	0.0261 (12)	0.0014 (10)	0.0004 (11)	0.0015 (11)
C9	0.0403 (15)	0.0260 (13)	0.0317 (13)	0.0018 (11)	0.0001 (12)	0.0001 (11)
C10	0.0500 (17)	0.0372 (17)	0.0321 (14)	-0.0146 (14)	0.0024 (13)	-0.0031 (12)
N11	0.0305 (12)	0.0627 (17)	0.0322 (11)	-0.0136 (12)	-0.0010 (10)	-0.0001 (12)
C12	0.0342 (15)	0.0499 (18)	0.0304 (14)	0.0011 (12)	0.0018 (12)	0.0038 (13)
C13	0.0307 (14)	0.0327 (16)	0.0357 (14)	0.0041 (10)	0.0040 (13)	0.0027 (11)
N14	0.0279 (11)	0.0295 (11)	0.0550 (15)	-0.0007 (8)	-0.0018 (11)	-0.0008 (10)

Geometric parameters (\AA , $^\circ$)

Co1—Cl3	2.2527 (8)	N7—H7A	0.8422
Co1—Cl4	2.2597 (8)	N7—H7B	0.8984
Co1—Cl2	2.2822 (8)	C8—N14	1.333 (3)
Co1—Cl1	2.2880 (8)	C8—C13	1.405 (4)
C1—N7	1.332 (3)	C8—C9	1.410 (4)

C1—C6	1.411 (4)	C9—C10	1.360 (4)
C1—C2	1.419 (4)	C9—H9	0.9500
C2—C3	1.361 (4)	C10—N11	1.344 (4)
C2—H2	0.9500	C10—H10	0.9500
C3—N4	1.335 (4)	N11—C12	1.360 (4)
C3—H3	0.9500	N11—H11	0.8737
N4—C5	1.345 (4)	C12—C13	1.348 (4)
N4—H4	0.8772	C12—H12	0.9500
C5—C6	1.359 (4)	C13—H13	0.9500
C5—H5	0.9500	N14—H14A	0.8967
C6—H6	0.9500	N14—H14B	0.8440
Cl3—Co1—Cl4	115.64 (3)	C1—N7—H7A	118.5
Cl3—Co1—Cl2	106.20 (4)	C1—N7—H7B	124.9
Cl4—Co1—Cl2	111.62 (3)	H7A—N7—H7B	116.6
Cl3—Co1—Cl1	110.24 (3)	N14—C8—C13	120.7 (3)
Cl4—Co1—Cl1	106.41 (4)	N14—C8—C9	121.0 (2)
Cl2—Co1—Cl1	106.41 (3)	C13—C8—C9	118.3 (3)
N7—C1—C6	121.8 (2)	C10—C9—C8	118.7 (3)
N7—C1—C2	121.0 (3)	C10—C9—H9	120.7
C6—C1—C2	117.2 (3)	C8—C9—H9	120.7
C3—C2—C1	119.8 (3)	N11—C10—C9	121.7 (3)
C3—C2—H2	120.1	N11—C10—H10	119.1
C1—C2—H2	120.1	C9—C10—H10	119.1
N4—C3—C2	121.0 (3)	C10—N11—C12	120.6 (3)
N4—C3—H3	119.5	C10—N11—H11	123.8
C2—C3—H3	119.5	C12—N11—H11	114.0
C3—N4—C5	121.2 (3)	C13—C12—N11	120.6 (3)
C3—N4—H4	123.8	C13—C12—H12	119.7
C5—N4—H4	114.8	N11—C12—H12	119.7
N4—C5—C6	121.2 (3)	C12—C13—C8	120.1 (3)
N4—C5—H5	119.4	C12—C13—H13	120.0
C6—C5—H5	119.4	C8—C13—H13	120.0
C5—C6—C1	119.6 (3)	C8—N14—H14A	122.7
C5—C6—H6	120.2	C8—N14—H14B	121.0
C1—C6—H6	120.2	H14A—N14—H14B	116.3
N7—C1—C2—C3	−178.6 (3)	N14—C8—C9—C10	−179.2 (3)
C6—C1—C2—C3	0.5 (4)	C13—C8—C9—C10	2.2 (4)
C1—C2—C3—N4	−0.4 (4)	C8—C9—C10—N11	−0.5 (4)
C2—C3—N4—C5	−0.2 (5)	C9—C10—N11—C12	−1.8 (4)
C3—N4—C5—C6	0.8 (4)	C10—N11—C12—C13	2.5 (5)
N4—C5—C6—C1	−0.7 (4)	N11—C12—C13—C8	−0.8 (4)
N7—C1—C6—C5	179.1 (3)	N14—C8—C13—C12	179.8 (3)
C2—C1—C6—C5	0.0 (4)	C9—C8—C13—C12	−1.5 (4)

Hydrogen-bond geometry (Å, °)


D—H···A	D—H	H···A	D···A	D—H···A
N4—H4···Cl2 ⁱ	0.88	2.94	3.563 (3)	130

supplementary materials

N4—H4···Cl3 ⁱ	0.88	2.67	3.335 (3)	134
N7—H7A···Cl2 ⁱⁱ	0.84	2.50	3.338 (2)	175
N7—H7B···Cl1	0.90	2.53	3.387 (2)	158
N11—H11···Cl1 ⁱⁱⁱ	0.87	2.52	3.272 (3)	144
N14—H14B···Cl2 ^{iv}	0.84	2.64	3.394 (3)	149
N14—H14A···Cl4	0.90	2.42	3.303 (2)	169

Symmetry codes: (i) $-x+1/2, -y, z-1/2$; (ii) $x-1/2, -y+1/2, -z+1$; (iii) $x-1/2, -y+1/2, -z$; (iv) $-x+1, y+1/2, -z+1/2$.

Fig. 1



supplementary materials

Fig. 2

