

Bromido{dicyclohexyl[2'-(dimethylamino)biphenyl-2-yl]phosphine- κP }-[2-(4,6-dimethylpyrimidin-2-yl)ferrocenyl- $\kappa^2 C^1, N$]palladium(II) dichloromethane solvate

Xin-Qi Hao,^a Hong-Mei Li,^b Hui Jiang,^a Yong-Tao Ma^a and Mao-Ping Song^{a*}

^aDepartment of Chemistry, Henan Key Laboratory of Chemical Biology and Organic Chemistry, Zhengzhou University, Zhengzhou 450052, People's Republic of China, and ^bLuoyang Normal University Library, Luoyang 471022, People's Republic of China

Correspondence e-mail: maopingsong@zzu.edu.cn


Received 25 March 2009; accepted 5 April 2009

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.006\text{ \AA}$; R factor = 0.038; wR factor = 0.099; data-to-parameter ratio = 16.7.

In the title compound, $[\text{FePdBr}(\text{C}_5\text{H}_5)(\text{C}_{11}\text{H}_{10}\text{N}_2)-(\text{C}_{26}\text{H}_{36}\text{NP})]\cdot\text{CH}_2\text{Cl}_2$, the Pd atom displays a distorted square-planar coordination environment. The five-membered metallacycle adopts an envelope conformation with the coordinated cyclopentadienyl C atom 0.4222 (4) \AA out of plane. The dihedral angle between the pyrimidinyl ring and substituted cyclopentadienyl ring is 21.47 (2) $^\circ$. In the crystal structure, the dimeric unit is generated through the C—H $\cdots\pi$ contact *via* a crystallographic inversion centre, while the C—H $\cdots\text{Cl}$ contacts in the dimeric centre link the dichlormethane molecules with the Pd complex molecules.

Related literature

For historical background of cyclopalladated compounds, see: Cope & Sickman (1965). For the properties of cyclopalladated compounds, see: Dupont *et al.* (2005); Gong *et al.* (2007); Xu *et al.* (2007). For related structures, see: Xu *et al.* (2008, 2009).



Experimental

Crystal data

$[\text{FePdBr}(\text{C}_5\text{H}_5)(\text{C}_{11}\text{H}_{10}\text{N}_2)-(\text{C}_{26}\text{H}_{36}\text{NP})]\cdot\text{CH}_2\text{Cl}_2$	$\beta = 76.590 (1)^\circ$
$M_r = 955.91$	$\gamma = 72.647 (1)^\circ$
Triclinic, $P\bar{1}$	$V = 2141.9 (3)\text{ \AA}^3$
$a = 10.2059 (9)\text{ \AA}$	$Z = 2$
$b = 10.3637 (9)\text{ \AA}$	Mo $K\alpha$ radiation
$c = 21.900 (2)\text{ \AA}$	$\mu = 1.89\text{ mm}^{-1}$
$\alpha = 81.195 (1)^\circ$	$T = 293\text{ K}$
	$0.37 \times 0.24 \times 0.21\text{ mm}$

Data collection

Bruker SMART APEX CCD area-detector diffractometer	16458 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Sheldrick, 1996)	7919 independent reflections
$T_{\min} = 0.540$, $T_{\max} = 0.697$	6716 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.017$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.038$	473 parameters
$wR(F^2) = 0.099$	H-atom parameters constrained
$S = 1.08$	$\Delta\rho_{\max} = 1.03\text{ e \AA}^{-3}$
7919 reflections	$\Delta\rho_{\min} = -0.69\text{ e \AA}^{-3}$

Table 1
Selected geometric parameters (\AA , $^\circ$).

Pd1—C1	1.997 (3)	Pd1—P1	2.2654 (9)
Pd1—N1	2.192 (3)	Pd1—Br1	2.6585 (6)
N1—Pd1—P1	173.00 (9)	C1—Pd1—Br1	160.40 (10)

Table 2
Hydrogen-bond geometry (\AA , $^\circ$).

D—H \cdots A	D—H	H \cdots A	D \cdots A	D—H \cdots A
C3—H3 \cdots Cl2 ⁱ	0.98	2.78	3.671 (5)	151
C16—H16C \cdots Cg1 ⁱⁱ	0.96	2.68	3.544 (6)	151

Symmetry codes: (i) $-x + 1, -y + 1, -z$; (ii) $-x + 1, -y + 2, -z$. Cg1 is the centroid of the C6—C10 Cp ring.

Data collection: *APEX2* (Bruker, 2004); cell refinement: *SAINT* (Bruker, 2004); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

This work was supported by the National Science Foundation of China (No. 20872133) and the Innovation Fund for Outstanding Scholars of Henan Province (No. 074200510005).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SI2165).

References

- Bruker (2004). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Cope, A. C. & Sickman, R. M. (1965). *J. Am. Chem. Soc.* **87**, 3272–3273.
- Dupont, J., Consorti, C. S. & Spencer, J. (2005). *Chem. Rev.* **105**, 2527–2571.
- Gong, J. F., Zhang, Y. H., Song, M. P. & Xu, C. (2007). *Organometallics*, **26**, 6487–6492.

metal-organic compounds

- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Xu, C., Gong, J. F., Guo, T., Zhang, Y. H. & Wu, Y. J. (2008). *J. Mol. Catal. A Chem.* **279**, 69–76.
- Xu, C., Gong, J. F. & Wu, Y. J. (2007). *Tetrahedron Lett.* **48**, 1619–1623.
- Xu, C., Wang, Z. Q., Fu, W. J., Lou, X. H., Li, Y. F., Cen, F. F., Ma, H. J. & Ji, B. M. (2009). *Organometallics*, **28**, 1909–1916.

supplementary materials

Acta Cryst. (2009). E65, m529-m530 [doi:10.1107/S1600536809012884]

Bromido{dicyclohexyl[2'-(dimethylamino)biphenyl-2-yl]phosphine- $\kappa P\}$ [2-(4,6-dimethylpyrimidin-2-yl)ferrocenyl- $\kappa^2 C^1,N$]palladium(II) dichloromethane solvate

X.-Q. Hao, H.-M. Li, H. Jiang, Y.-T. Ma and M.-P. Song

Comment

Cyclopalladated compounds containing a Pd—C bond intramolecularly stabilized by one donor atom were first reported in the middle 1960's (Cope & Sickman, 1965). Forty years later, the chemistry of these compound has developed into one of the most fruitful fields in organometallic chemistry (Dupont *et al.*, 2005). In the palladium-catalyzed coupling reactions, palladacycles have shown enormous superiority in many respects (Gong *et al.*, 2007; Xu *et al.*, 2007).

The Pd atom in the title complex is in a distorted square-planar environment bonded to the atoms shown in Fig. 1: the Br atom, the pyrimidinyl nitrogen atom and the carbon atom of the ferrocenyl moiety. The Pd—C1 (N1,P1,Br1) bond lengths and the P1—Pd1—N1 and C1—Pd1—Br1 bond angles (Table 1) are within comparable ranges of 162.08 (7) ° - 174.76 (7) ° and 162.27 (9) ° - 166.04 (8) ° (C—Pd—Cl), respectively (Xu *et al.*, 2009). The five-membered metallacycle (containing atoms C1/C2/C11/N1/Pd1) adopts an envelope conformation, with the C1 atom 0.4222 (4) Å out of plane. The dihedral angle between the pyrimidinyl ring and substituted cyclopentadienyl ring is 21.47 (2)°. In the crystal structure, the dimeric unit (Fig.2) is generated through the C—H···π contacts *via* a crystallographic inversion centre, and the C—H···Cl contact (Table 2) connects a symmetry-related dichlormethane molecule with the Pd complex (Table 2). This hydrogen bonding motif is different from those of related cyclopalladated ferrocene derivatives (Xu *et al.*, 2008; Xu *et al.*, 2009), in which C—H···Cl hydrogen bonds construct the one-dimensional chain structures.

Experimental

The title compound was prepared as described in literature (Xu *et al.*, 2009), using Li_2PdBr_4 instead of Li_2PdCl_4 and recrystallized from dichloromethane-petroleum ether solution at room temperature to give the desired product as red crystals suitable for single-crystal X-ray diffraction.

Refinement

H atoms attached to C atoms of the title compound were placed in geometrically idealized positions and treated as riding with C—H distances constrained to 0.93–0.96 Å, and with $U_{\text{iso}}(\text{H})=1.2U_{\text{eq}}(\text{C})$ ($1.5U_{\text{eq}}$ for methyl H).

Figures




Fig. 1. The molecular structure of the title compound with displacement ellipsoids at the 30% probability level. The dichloromethane molecule is omitted for clarity.

supplementary materials




Fig. 2. The dimeric unit of the title compound formed by the intermolecular C—H···Cl hydrogen bonds and C—H···π interactions. $Cg1$ is the centroid of the Cp ring C6—C10, symmetry code ii: 1 - x , 2 - y , - z .

Bromido{dicyclohexyl[2'-(dimethylamino)biphenyl-2-yl]phosphine-κP}[2- (4,6-dimethylpyrimidin-2-yl)ferrocenyl-κ²C¹,N]palladium(II) dichloromethane solvate

Crystal data

[FePdBr(C ₅ H ₅)(C ₁₁ H ₁₀ N ₂)(C ₂₆ H ₃₆ NP)]·CH ₂ Cl ₂	$Z = 2$
$M_r = 955.91$	$F_{000} = 976$
Triclinic, $P\bar{1}$	$D_x = 1.482 \text{ Mg m}^{-3}$
$a = 10.2059 (9) \text{ \AA}$	Mo $K\alpha$ radiation
$b = 10.3637 (9) \text{ \AA}$	$\lambda = 0.71073 \text{ \AA}$
$c = 21.900 (2) \text{ \AA}$	Cell parameters from 7107 reflections
$\alpha = 81.1950 (10)^\circ$	$\theta = 2.5\text{--}27.1^\circ$
$\beta = 76.5900 (10)^\circ$	$\mu = 1.89 \text{ mm}^{-1}$
$\gamma = 72.6470 (10)^\circ$	$T = 293 \text{ K}$
$V = 2141.9 (3) \text{ \AA}^3$	Block, red
	$0.37 \times 0.24 \times 0.21 \text{ mm}$

Data collection

Bruker SMART APEX CCD area-detector diffractometer	7919 independent reflections
Radiation source: fine-focus sealed tube	6716 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.017$
$T = 293 \text{ K}$	$\theta_{\text{max}} = 25.5^\circ$
φ and ω scans	$\theta_{\text{min}} = 2.5^\circ$
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	$h = -12\text{--}10$
$T_{\text{min}} = 0.540$, $T_{\text{max}} = 0.697$	$k = -12\text{--}12$
16458 measured reflections	$l = -26\text{--}26$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.038$	H-atom parameters constrained
$wR(F^2) = 0.099$	$w = 1/[σ^2(F_o^2) + (0.0421P)^2 + 2.3428P]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.08$	$(Δ/σ)_{\text{max}} = 0.001$
7919 reflections	$Δρ_{\text{max}} = 1.03 \text{ e \AA}^{-3}$
473 parameters	$Δρ_{\text{min}} = -0.69 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes)

are estimated using the full covariance matrix. The cell e.s.d.'s are taken

into account individually in the estimation of e.s.d.'s in distances, angles

and torsion angles; correlations between e.s.d.'s in cell parameters are only

used when they are defined by crystal symmetry. An approximate (isotropic)

treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Pd1	0.29216 (3)	0.66381 (3)	0.217306 (12)	0.03942 (9)
Fe1	0.63058 (6)	0.70760 (5)	0.12158 (2)	0.04439 (14)
P1	0.33910 (9)	0.56740 (8)	0.31270 (4)	0.03503 (19)
Br1	0.02020 (5)	0.68171 (6)	0.25105 (2)	0.07101 (15)
Cl2	0.1066 (2)	0.5000 (3)	0.07189 (12)	0.1586 (9)
Cl3	0.2456 (4)	0.2376 (3)	0.12552 (12)	0.1980 (14)
N1	0.2744 (3)	0.7531 (3)	0.12117 (14)	0.0477 (7)
N2	0.3968 (4)	0.7000 (3)	0.01637 (15)	0.0592 (9)
N3	0.7663 (3)	0.2896 (3)	0.29096 (16)	0.0565 (8)
C1	0.4926 (4)	0.6039 (3)	0.17412 (16)	0.0395 (7)
C2	0.5018 (4)	0.6044 (3)	0.10732 (16)	0.0443 (8)
C3	0.6430 (4)	0.5430 (4)	0.07875 (18)	0.0537 (10)
H3	0.6788	0.5297	0.0339	0.064*
C4	0.7212 (4)	0.5024 (4)	0.1271 (2)	0.0544 (10)
H4	0.8219	0.4579	0.1214	0.065*
C5	0.6299 (4)	0.5404 (3)	0.18529 (18)	0.0449 (8)
H5	0.6578	0.5268	0.2262	0.054*
C6	0.6949 (6)	0.8426 (5)	0.1587 (2)	0.0754 (14)
H6	0.7205	0.8302	0.2001	0.091*
C7	0.5625 (6)	0.9016 (4)	0.1460 (3)	0.0746 (14)
H7	0.4782	0.9384	0.1771	0.090*
C8	0.5682 (7)	0.9016 (5)	0.0818 (3)	0.0812 (16)
H8	0.4892	0.9383	0.0601	0.097*
C9	0.7057 (8)	0.8393 (6)	0.0538 (3)	0.096 (2)
H9	0.7411	0.8256	0.0090	0.115*
C10	0.7859 (5)	0.8021 (5)	0.1023 (3)	0.0872 (17)
H10	0.8866	0.7577	0.0970	0.105*

supplementary materials

C11	0.3860 (4)	0.6874 (4)	0.07887 (17)	0.0477 (9)
C12	0.2921 (6)	0.7939 (5)	-0.0063 (2)	0.0674 (13)
C13	0.1860 (6)	0.8765 (5)	0.0328 (2)	0.0724 (13)
H13	0.1202	0.9473	0.0159	0.087*
C14	0.1768 (5)	0.8542 (4)	0.0977 (2)	0.0618 (11)
C15	0.0679 (5)	0.9476 (5)	0.1408 (3)	0.0829 (15)
H15A	-0.0102	0.9103	0.1570	0.124*
H15B	0.0371	1.0344	0.1180	0.124*
H15C	0.1064	0.9585	0.1751	0.124*
C16	0.3020 (7)	0.8073 (5)	-0.0767 (2)	0.0927 (19)
H16A	0.2361	0.7675	-0.0862	0.139*
H16B	0.3951	0.7613	-0.0967	0.139*
H16C	0.2812	0.9016	-0.0921	0.139*
C17	0.3460 (4)	0.3850 (3)	0.32533 (16)	0.0397 (8)
H17	0.3978	0.3440	0.3592	0.048*
C18	0.2011 (4)	0.3582 (4)	0.3453 (2)	0.0540 (10)
H18A	0.1514	0.3996	0.3839	0.065*
H18B	0.1467	0.3982	0.3128	0.065*
C19	0.2181 (5)	0.2047 (4)	0.3555 (2)	0.0662 (12)
H19A	0.2673	0.1663	0.3899	0.079*
H19B	0.1264	0.1885	0.3674	0.079*
C20	0.2977 (5)	0.1351 (4)	0.2974 (2)	0.0716 (13)
H20A	0.2446	0.1673	0.2640	0.086*
H20B	0.3099	0.0380	0.3062	0.086*
C21	0.4392 (5)	0.1628 (4)	0.2760 (2)	0.0682 (12)
H21A	0.4859	0.1223	0.2368	0.082*
H21B	0.4962	0.1206	0.3073	0.082*
C22	0.4266 (4)	0.3154 (4)	0.26618 (19)	0.0518 (9)
H22A	0.5194	0.3291	0.2551	0.062*
H22B	0.3789	0.3559	0.2315	0.062*
C23	0.1984 (4)	0.6508 (3)	0.37623 (16)	0.0409 (8)
H23	0.1134	0.6308	0.3716	0.049*
C24	0.2157 (4)	0.5979 (4)	0.44399 (17)	0.0508 (9)
H24A	0.2947	0.6205	0.4527	0.061*
H24B	0.2339	0.4999	0.4491	0.061*
C25	0.0821 (5)	0.6618 (5)	0.4903 (2)	0.0681 (13)
H25A	0.0948	0.6303	0.5331	0.082*
H25B	0.0047	0.6332	0.4835	0.082*
C26	0.0467 (5)	0.8173 (5)	0.4817 (2)	0.0737 (13)
H26A	-0.0408	0.8554	0.5098	0.088*
H26B	0.1196	0.8466	0.4926	0.088*
C27	0.0340 (5)	0.8691 (4)	0.4137 (2)	0.0623 (11)
H27A	-0.0448	0.8474	0.4042	0.075*
H27B	0.0166	0.9670	0.4085	0.075*
C28	0.1661 (4)	0.8057 (4)	0.36824 (19)	0.0503 (9)
H28A	0.2440	0.8324	0.3758	0.060*
H28B	0.1547	0.8385	0.3254	0.060*
C29	0.4941 (3)	0.5949 (3)	0.33234 (15)	0.0372 (7)
C30	0.5028 (4)	0.7283 (3)	0.31298 (17)	0.0443 (8)

H30	0.4448	0.7823	0.2864	0.053*
C31	0.5932 (4)	0.7823 (4)	0.3316 (2)	0.0554 (10)
H31	0.5953	0.8713	0.3178	0.066*
C32	0.6802 (5)	0.7048 (5)	0.3703 (2)	0.0632 (11)
H32	0.7396	0.7413	0.3845	0.076*
C33	0.6785 (5)	0.5717 (4)	0.3881 (2)	0.0609 (11)
H33	0.7388	0.5194	0.4140	0.073*
C34	0.5909 (4)	0.5118 (4)	0.36905 (17)	0.0438 (8)
C35	0.6087 (4)	0.3642 (4)	0.39042 (18)	0.0460 (8)
C36	0.6959 (4)	0.2591 (4)	0.3527 (2)	0.0526 (9)
C37	0.7073 (5)	0.1256 (4)	0.3780 (2)	0.0662 (12)
H37	0.7620	0.0554	0.3536	0.079*
C38	0.6395 (6)	0.0938 (5)	0.4387 (3)	0.0771 (14)
H38	0.6486	0.0035	0.4542	0.092*
C39	0.5598 (5)	0.1945 (5)	0.4756 (2)	0.0699 (12)
H39	0.5159	0.1734	0.5165	0.084*
C40	0.5445 (4)	0.3289 (4)	0.45172 (19)	0.0553 (10)
H40	0.4900	0.3973	0.4772	0.066*
C42	0.8764 (5)	0.3547 (6)	0.2856 (3)	0.0867 (16)
H42A	0.8416	0.4319	0.3099	0.130*
H42B	0.9064	0.3839	0.2422	0.130*
H42C	0.9541	0.2915	0.3012	0.130*
C43	0.8130 (6)	0.1821 (5)	0.2488 (3)	0.0862 (16)
H43A	0.8887	0.1117	0.2625	0.129*
H43B	0.8445	0.2183	0.2068	0.129*
H43C	0.7368	0.1453	0.2495	0.129*
C44	0.2410 (10)	0.4080 (8)	0.1087 (4)	0.138 (3)
H44A	0.2313	0.4467	0.1477	0.166*
H44B	0.3292	0.4154	0.0820	0.166*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Pd1	0.03597 (15)	0.04066 (15)	0.04152 (15)	-0.01016 (11)	-0.01180 (11)	0.00234 (11)
Fe1	0.0491 (3)	0.0377 (3)	0.0443 (3)	-0.0144 (2)	-0.0046 (2)	0.0003 (2)
P1	0.0351 (4)	0.0315 (4)	0.0382 (4)	-0.0091 (3)	-0.0079 (3)	-0.0011 (3)
Br1	0.0486 (2)	0.0991 (4)	0.0643 (3)	-0.0267 (2)	-0.0149 (2)	0.0144 (2)
Cl2	0.0879 (13)	0.193 (2)	0.161 (2)	-0.0153 (13)	0.0182 (13)	-0.0262 (17)
Cl3	0.345 (5)	0.142 (2)	0.1273 (18)	-0.095 (2)	-0.063 (2)	0.0086 (15)
N1	0.0528 (19)	0.0448 (17)	0.0487 (18)	-0.0164 (15)	-0.0190 (15)	0.0061 (14)
N2	0.096 (3)	0.0469 (19)	0.0463 (19)	-0.0302 (19)	-0.0288 (18)	0.0041 (15)
N3	0.0462 (18)	0.0532 (19)	0.062 (2)	-0.0014 (15)	-0.0091 (16)	-0.0063 (16)
C1	0.0429 (19)	0.0334 (17)	0.0402 (18)	-0.0112 (15)	-0.0069 (15)	0.0024 (14)
C2	0.059 (2)	0.0326 (17)	0.0422 (19)	-0.0151 (16)	-0.0081 (17)	-0.0029 (15)
C3	0.070 (3)	0.039 (2)	0.045 (2)	-0.0119 (18)	0.0007 (19)	-0.0070 (16)
C4	0.047 (2)	0.041 (2)	0.065 (3)	-0.0057 (17)	0.0009 (19)	-0.0053 (18)
C5	0.043 (2)	0.0377 (19)	0.047 (2)	-0.0068 (15)	-0.0078 (16)	0.0053 (15)
C6	0.107 (4)	0.071 (3)	0.069 (3)	-0.053 (3)	-0.028 (3)	0.007 (2)

supplementary materials

C7	0.094 (4)	0.040 (2)	0.089 (4)	-0.027 (2)	-0.003 (3)	-0.011 (2)
C8	0.117 (5)	0.046 (3)	0.093 (4)	-0.034 (3)	-0.050 (4)	0.025 (3)
C9	0.159 (6)	0.076 (4)	0.058 (3)	-0.075 (4)	0.020 (4)	-0.004 (3)
C10	0.057 (3)	0.066 (3)	0.136 (5)	-0.030 (2)	-0.002 (3)	-0.001 (3)
C11	0.069 (3)	0.0349 (18)	0.047 (2)	-0.0231 (18)	-0.0209 (19)	0.0027 (16)
C12	0.112 (4)	0.050 (2)	0.060 (3)	-0.042 (3)	-0.045 (3)	0.016 (2)
C13	0.092 (4)	0.058 (3)	0.078 (3)	-0.023 (3)	-0.052 (3)	0.019 (2)
C14	0.063 (3)	0.054 (2)	0.074 (3)	-0.018 (2)	-0.034 (2)	0.011 (2)
C15	0.067 (3)	0.067 (3)	0.100 (4)	0.005 (2)	-0.025 (3)	0.003 (3)
C16	0.177 (6)	0.064 (3)	0.063 (3)	-0.052 (4)	-0.066 (4)	0.018 (2)
C17	0.0454 (19)	0.0312 (17)	0.0453 (19)	-0.0144 (15)	-0.0129 (15)	0.0022 (14)
C18	0.056 (2)	0.049 (2)	0.062 (2)	-0.0258 (19)	-0.0076 (19)	-0.0030 (18)
C19	0.073 (3)	0.053 (2)	0.080 (3)	-0.036 (2)	-0.014 (2)	0.006 (2)
C20	0.090 (3)	0.042 (2)	0.093 (4)	-0.030 (2)	-0.022 (3)	-0.008 (2)
C21	0.076 (3)	0.038 (2)	0.089 (3)	-0.015 (2)	-0.008 (3)	-0.013 (2)
C22	0.057 (2)	0.040 (2)	0.057 (2)	-0.0152 (17)	-0.0027 (18)	-0.0091 (17)
C23	0.0379 (18)	0.0372 (18)	0.0446 (19)	-0.0077 (14)	-0.0058 (15)	-0.0026 (15)
C24	0.055 (2)	0.047 (2)	0.043 (2)	-0.0045 (18)	-0.0075 (17)	-0.0050 (16)
C25	0.071 (3)	0.067 (3)	0.046 (2)	0.003 (2)	-0.002 (2)	-0.003 (2)
C26	0.068 (3)	0.069 (3)	0.069 (3)	0.001 (2)	0.004 (2)	-0.025 (2)
C27	0.062 (3)	0.037 (2)	0.074 (3)	0.0025 (18)	-0.002 (2)	-0.0109 (19)
C28	0.053 (2)	0.0358 (19)	0.057 (2)	-0.0084 (17)	-0.0060 (18)	-0.0027 (16)
C29	0.0369 (17)	0.0375 (17)	0.0375 (17)	-0.0094 (14)	-0.0087 (14)	-0.0040 (14)
C30	0.045 (2)	0.0369 (18)	0.052 (2)	-0.0125 (16)	-0.0096 (16)	-0.0033 (16)
C31	0.061 (2)	0.046 (2)	0.065 (3)	-0.0234 (19)	-0.012 (2)	-0.0048 (19)
C32	0.063 (3)	0.063 (3)	0.079 (3)	-0.030 (2)	-0.024 (2)	-0.013 (2)
C33	0.059 (3)	0.063 (3)	0.069 (3)	-0.018 (2)	-0.032 (2)	0.001 (2)
C34	0.045 (2)	0.0404 (19)	0.045 (2)	-0.0093 (16)	-0.0131 (16)	-0.0013 (15)
C35	0.045 (2)	0.042 (2)	0.052 (2)	-0.0058 (16)	-0.0220 (17)	0.0005 (16)
C36	0.045 (2)	0.047 (2)	0.062 (2)	-0.0005 (17)	-0.0219 (18)	0.0005 (18)
C37	0.068 (3)	0.045 (2)	0.076 (3)	0.001 (2)	-0.021 (2)	0.000 (2)
C38	0.088 (4)	0.047 (3)	0.088 (4)	-0.009 (2)	-0.026 (3)	0.012 (2)
C39	0.081 (3)	0.065 (3)	0.062 (3)	-0.022 (3)	-0.019 (2)	0.011 (2)
C40	0.062 (3)	0.054 (2)	0.050 (2)	-0.013 (2)	-0.0198 (19)	0.0008 (18)
C42	0.051 (3)	0.109 (4)	0.098 (4)	-0.023 (3)	-0.007 (3)	-0.011 (3)
C43	0.085 (4)	0.076 (3)	0.082 (4)	0.000 (3)	-0.001 (3)	-0.025 (3)
C44	0.198 (9)	0.139 (7)	0.125 (6)	-0.101 (6)	-0.080 (6)	0.033 (5)

Geometric parameters (\AA , $^\circ$)

Pd1—C1	1.997 (3)	C18—H18A	0.9700
Pd1—N1	2.192 (3)	C18—H18B	0.9700
Pd1—P1	2.2654 (9)	C19—C20	1.501 (6)
Pd1—Br1	2.6585 (6)	C19—H19A	0.9700
Fe1—C2	2.025 (4)	C19—H19B	0.9700
Fe1—C3	2.028 (4)	C20—C21	1.510 (6)
Fe1—C9	2.032 (5)	C20—H20A	0.9700
Fe1—C7	2.034 (4)	C20—H20B	0.9700
Fe1—C10	2.035 (5)	C21—C22	1.533 (5)

Fe1—C8	2.041 (4)	C21—H21A	0.9700
Fe1—C4	2.047 (4)	C21—H21B	0.9700
Fe1—C6	2.047 (5)	C22—H22A	0.9700
Fe1—C5	2.053 (3)	C22—H22B	0.9700
Fe1—C1	2.058 (3)	C23—C28	1.530 (5)
P1—C29	1.841 (3)	C23—C24	1.532 (5)
P1—C17	1.851 (3)	C23—H23	0.9800
P1—C23	1.857 (3)	C24—C25	1.533 (5)
Cl2—C44	1.715 (8)	C24—H24A	0.9700
Cl3—C44	1.737 (7)	C24—H24B	0.9700
N1—C14	1.343 (5)	C25—C26	1.535 (6)
N1—C11	1.364 (5)	C25—H25A	0.9700
N2—C11	1.337 (5)	C25—H25B	0.9700
N2—C12	1.346 (6)	C26—C27	1.525 (6)
N3—C36	1.414 (5)	C26—H26A	0.9700
N3—C42	1.449 (6)	C26—H26B	0.9700
N3—C43	1.454 (6)	C27—C28	1.513 (5)
C1—C5	1.420 (5)	C27—H27A	0.9700
C1—C2	1.443 (5)	C27—H27B	0.9700
C2—C3	1.425 (5)	C28—H28A	0.9700
C2—C11	1.448 (5)	C28—H28B	0.9700
C3—C4	1.411 (6)	C29—C30	1.405 (5)
C3—H3	0.9800	C29—C34	1.419 (5)
C4—C5	1.425 (5)	C30—C31	1.372 (5)
C4—H4	0.9800	C30—H30	0.9300
C5—H5	0.9800	C31—C32	1.368 (6)
C6—C7	1.381 (7)	C31—H31	0.9300
C6—C10	1.398 (8)	C32—C33	1.379 (6)
C6—H6	0.9800	C32—H32	0.9300
C7—C8	1.394 (7)	C33—C34	1.393 (5)
C7—H7	0.9800	C33—H33	0.9300
C8—C9	1.393 (8)	C34—C35	1.499 (5)
C8—H8	0.9800	C35—C40	1.396 (5)
C9—C10	1.427 (8)	C35—C36	1.420 (5)
C9—H9	0.9800	C36—C37	1.391 (6)
C10—H10	0.9800	C37—C38	1.389 (7)
C12—C13	1.371 (7)	C37—H37	0.9300
C12—C16	1.510 (6)	C38—C39	1.360 (7)
C13—C14	1.391 (6)	C38—H38	0.9300
C13—H13	0.9300	C39—C40	1.386 (6)
C14—C15	1.488 (7)	C39—H39	0.9300
C15—H15A	0.9600	C40—H40	0.9300
C15—H15B	0.9600	C42—H42A	0.9600
C15—H15C	0.9600	C42—H42B	0.9600
C16—H16A	0.9600	C42—H42C	0.9600
C16—H16B	0.9600	C43—H43A	0.9600
C16—H16C	0.9600	C43—H43B	0.9600
C17—C22	1.521 (5)	C43—H43C	0.9600
C17—C18	1.537 (5)	C44—H44A	0.9700

supplementary materials

C17—H17	0.9800	C44—H44B	0.9700
C18—C19	1.534 (5)		
C1—Pd1—N1	80.12 (13)	C14—C15—H15B	109.5
C1—Pd1—P1	93.07 (10)	H15A—C15—H15B	109.5
N1—Pd1—P1	173.00 (9)	C14—C15—H15C	109.5
C1—Pd1—Br1	160.40 (10)	H15A—C15—H15C	109.5
N1—Pd1—Br1	92.84 (8)	H15B—C15—H15C	109.5
P1—Pd1—Br1	94.13 (3)	C12—C16—H16A	109.5
C2—Fe1—C3	41.18 (15)	C12—C16—H16B	109.5
C2—Fe1—C9	122.5 (2)	H16A—C16—H16B	109.5
C3—Fe1—C9	106.90 (19)	C12—C16—H16C	109.5
C2—Fe1—C7	123.4 (2)	H16A—C16—H16C	109.5
C3—Fe1—C7	159.1 (2)	H16B—C16—H16C	109.5
C9—Fe1—C7	67.4 (2)	C22—C17—C18	109.9 (3)
C2—Fe1—C10	159.3 (2)	C22—C17—P1	109.9 (2)
C3—Fe1—C10	122.8 (2)	C18—C17—P1	113.7 (2)
C9—Fe1—C10	41.1 (2)	C22—C17—H17	107.7
C7—Fe1—C10	67.2 (2)	C18—C17—H17	107.7
C2—Fe1—C8	107.86 (19)	P1—C17—H17	107.7
C3—Fe1—C8	122.8 (2)	C19—C18—C17	109.7 (3)
C9—Fe1—C8	40.0 (2)	C19—C18—H18A	109.7
C7—Fe1—C8	40.0 (2)	C17—C18—H18A	109.7
C10—Fe1—C8	67.7 (2)	C19—C18—H18B	109.7
C2—Fe1—C4	68.24 (16)	C17—C18—H18B	109.7
C3—Fe1—C4	40.50 (17)	H18A—C18—H18B	108.2
C9—Fe1—C4	123.1 (2)	C20—C19—C18	111.8 (4)
C7—Fe1—C4	159.5 (2)	C20—C19—H19A	109.3
C10—Fe1—C4	108.1 (2)	C18—C19—H19A	109.3
C8—Fe1—C4	158.8 (2)	C20—C19—H19B	109.3
C2—Fe1—C6	158.8 (2)	C18—C19—H19B	109.3
C3—Fe1—C6	159.1 (2)	H19A—C19—H19B	107.9
C9—Fe1—C6	67.9 (2)	C19—C20—C21	110.7 (4)
C7—Fe1—C6	39.6 (2)	C19—C20—H20A	109.5
C10—Fe1—C6	40.1 (2)	C21—C20—H20A	109.5
C8—Fe1—C6	67.1 (2)	C19—C20—H20B	109.5
C4—Fe1—C6	124.0 (2)	C21—C20—H20B	109.5
C2—Fe1—C5	68.29 (15)	H20A—C20—H20B	108.1
C3—Fe1—C5	68.69 (16)	C20—C21—C22	111.7 (4)
C9—Fe1—C5	159.4 (3)	C20—C21—H21A	109.3
C7—Fe1—C5	123.73 (19)	C22—C21—H21A	109.3
C10—Fe1—C5	123.2 (2)	C20—C21—H21B	109.3
C8—Fe1—C5	159.2 (2)	C22—C21—H21B	109.3
C4—Fe1—C5	40.68 (15)	H21A—C21—H21B	107.9
C6—Fe1—C5	108.64 (18)	C17—C22—C21	111.2 (3)
C2—Fe1—C1	41.39 (14)	C17—C22—H22A	109.4
C3—Fe1—C1	69.73 (15)	C21—C22—H22A	109.4
C9—Fe1—C1	158.8 (3)	C17—C22—H22B	109.4
C7—Fe1—C1	107.86 (17)	C21—C22—H22B	109.4
C10—Fe1—C1	158.2 (2)	H22A—C22—H22B	108.0

C8—Fe1—C1	123.1 (2)	C28—C23—C24	110.0 (3)
C4—Fe1—C1	68.68 (14)	C28—C23—P1	113.1 (2)
C6—Fe1—C1	122.59 (18)	C24—C23—P1	116.3 (2)
C5—Fe1—C1	40.43 (14)	C28—C23—H23	105.5
C29—P1—C17	110.59 (16)	C24—C23—H23	105.5
C29—P1—C23	101.08 (16)	P1—C23—H23	105.5
C17—P1—C23	106.19 (16)	C23—C24—C25	109.8 (3)
C29—P1—Pd1	115.27 (11)	C23—C24—H24A	109.7
C17—P1—Pd1	112.84 (12)	C25—C24—H24A	109.7
C23—P1—Pd1	109.86 (11)	C23—C24—H24B	109.7
C14—N1—C11	117.0 (3)	C25—C24—H24B	109.7
C14—N1—Pd1	132.7 (3)	H24A—C24—H24B	108.2
C11—N1—Pd1	110.3 (2)	C24—C25—C26	111.1 (4)
C11—N2—C12	116.1 (4)	C24—C25—H25A	109.4
C36—N3—C42	116.4 (4)	C26—C25—H25A	109.4
C36—N3—C43	116.8 (4)	C24—C25—H25B	109.4
C42—N3—C43	109.8 (4)	C26—C25—H25B	109.4
C5—C1—C2	106.1 (3)	H25A—C25—H25B	108.0
C5—C1—Pd1	142.9 (3)	C27—C26—C25	110.4 (4)
C2—C1—Pd1	109.9 (3)	C27—C26—H26A	109.6
C5—C1—Fe1	69.6 (2)	C25—C26—H26A	109.6
C2—C1—Fe1	68.06 (19)	C27—C26—H26B	109.6
Pd1—C1—Fe1	132.86 (17)	C25—C26—H26B	109.6
C3—C2—C1	109.1 (3)	H26A—C26—H26B	108.1
C3—C2—C11	130.0 (3)	C28—C27—C26	111.1 (3)
C1—C2—C11	119.3 (3)	C28—C27—H27A	109.4
C3—C2—Fe1	69.5 (2)	C26—C27—H27A	109.4
C1—C2—Fe1	70.6 (2)	C28—C27—H27B	109.4
C11—C2—Fe1	114.3 (2)	C26—C27—H27B	109.4
C4—C3—C2	107.3 (3)	H27A—C27—H27B	108.0
C4—C3—Fe1	70.5 (2)	C27—C28—C23	110.7 (3)
C2—C3—Fe1	69.3 (2)	C27—C28—H28A	109.5
C4—C3—H3	126.3	C23—C28—H28A	109.5
C2—C3—H3	126.3	C27—C28—H28B	109.5
Fe1—C3—H3	126.3	C23—C28—H28B	109.5
C3—C4—C5	108.6 (3)	H28A—C28—H28B	108.1
C3—C4—Fe1	69.0 (2)	C30—C29—C34	117.3 (3)
C5—C4—Fe1	69.9 (2)	C30—C29—P1	111.3 (3)
C3—C4—H4	125.7	C34—C29—P1	131.0 (3)
C5—C4—H4	125.7	C31—C30—C29	122.9 (4)
Fe1—C4—H4	125.7	C31—C30—H30	118.6
C1—C5—C4	109.0 (3)	C29—C30—H30	118.6
C1—C5—Fe1	69.99 (19)	C32—C31—C30	119.7 (4)
C4—C5—Fe1	69.5 (2)	C32—C31—H31	120.1
C1—C5—H5	125.5	C30—C31—H31	120.1
C4—C5—H5	125.5	C31—C32—C33	118.9 (4)
Fe1—C5—H5	125.5	C31—C32—H32	120.6
C7—C6—C10	108.1 (5)	C33—C32—H32	120.6
C7—C6—Fe1	69.7 (3)	C32—C33—C34	123.2 (4)

supplementary materials

C10—C6—Fe1	69.5 (3)	C32—C33—H33	118.4
C7—C6—H6	125.9	C34—C33—H33	118.4
C10—C6—H6	125.9	C33—C34—C29	117.8 (3)
Fe1—C6—H6	125.9	C33—C34—C35	115.7 (3)
C6—C7—C8	109.0 (5)	C29—C34—C35	126.5 (3)
C6—C7—Fe1	70.8 (3)	C40—C35—C36	118.7 (4)
C8—C7—Fe1	70.3 (3)	C40—C35—C34	118.3 (3)
C6—C7—H7	125.5	C36—C35—C34	122.8 (3)
C8—C7—H7	125.5	C37—C36—N3	121.2 (4)
Fe1—C7—H7	125.5	C37—C36—C35	117.9 (4)
C9—C8—C7	108.1 (5)	N3—C36—C35	120.9 (3)
C9—C8—Fe1	69.7 (3)	C38—C37—C36	122.0 (4)
C7—C8—Fe1	69.7 (3)	C38—C37—H37	119.0
C9—C8—H8	125.9	C36—C37—H37	119.0
C7—C8—H8	125.9	C39—C38—C37	120.1 (4)
Fe1—C8—H8	125.9	C39—C38—H38	119.9
C8—C9—C10	107.2 (5)	C37—C38—H38	119.9
C8—C9—Fe1	70.3 (3)	C38—C39—C40	119.6 (4)
C10—C9—Fe1	69.6 (3)	C38—C39—H39	120.2
C8—C9—H9	126.4	C40—C39—H39	120.2
C10—C9—H9	126.4	C39—C40—C35	121.7 (4)
Fe1—C9—H9	126.4	C39—C40—H40	119.1
C6—C10—C9	107.5 (5)	C35—C40—H40	119.1
C6—C10—Fe1	70.4 (3)	N3—C42—H42A	109.5
C9—C10—Fe1	69.3 (3)	N3—C42—H42B	109.5
C6—C10—H10	126.3	H42A—C42—H42B	109.5
C9—C10—H10	126.3	N3—C42—H42C	109.5
Fe1—C10—H10	126.3	H42A—C42—H42C	109.5
N2—C11—N1	125.8 (4)	H42B—C42—H42C	109.5
N2—C11—C2	120.5 (4)	N3—C43—H43A	109.5
N1—C11—C2	113.6 (3)	N3—C43—H43B	109.5
N2—C12—C13	121.1 (4)	H43A—C43—H43B	109.5
N2—C12—C16	116.5 (5)	N3—C43—H43C	109.5
C13—C12—C16	122.3 (5)	H43A—C43—H43C	109.5
C12—C13—C14	120.0 (4)	H43B—C43—H43C	109.5
C12—C13—H13	120.0	C12—C44—Cl3	112.9 (4)
C14—C13—H13	120.0	C12—C44—H44A	109.0
N1—C14—C13	119.2 (4)	Cl3—C44—H44A	109.0
N1—C14—C15	119.8 (4)	Cl2—C44—H44B	109.0
C13—C14—C15	120.8 (4)	Cl3—C44—H44B	109.0
C14—C15—H15A	109.5	H44A—C44—H44B	107.8
C1—Pd1—P1—C29	43.93 (15)	C10—C6—C7—Fe1	59.0 (3)
N1—Pd1—P1—C29	31.0 (7)	C2—Fe1—C7—C6	162.9 (3)
Br1—Pd1—P1—C29	-154.31 (12)	C3—Fe1—C7—C6	-160.3 (4)
C1—Pd1—P1—C17	-84.43 (15)	C9—Fe1—C7—C6	-82.1 (4)
N1—Pd1—P1—C17	-97.3 (7)	C10—Fe1—C7—C6	-37.4 (3)
Br1—Pd1—P1—C17	77.32 (12)	C8—Fe1—C7—C6	-119.5 (5)
C1—Pd1—P1—C23	157.29 (15)	C4—Fe1—C7—C6	43.1 (7)
N1—Pd1—P1—C23	144.4 (7)	C5—Fe1—C7—C6	78.2 (3)

Br1—Pd1—P1—C23	-40.96 (12)	C1—Fe1—C7—C6	119.9 (3)
C1—Pd1—N1—C14	-159.6 (4)	C2—Fe1—C7—C8	-77.6 (4)
P1—Pd1—N1—C14	-146.5 (6)	C3—Fe1—C7—C8	-40.8 (7)
Br1—Pd1—N1—C14	38.9 (4)	C9—Fe1—C7—C8	37.3 (4)
C1—Pd1—N1—C11	21.8 (2)	C10—Fe1—C7—C8	82.1 (4)
P1—Pd1—N1—C11	34.9 (8)	C4—Fe1—C7—C8	162.6 (5)
Br1—Pd1—N1—C11	-139.8 (2)	C6—Fe1—C7—C8	119.5 (5)
N1—Pd1—C1—C5	172.1 (4)	C5—Fe1—C7—C8	-162.3 (3)
P1—Pd1—C1—C5	-6.3 (4)	C1—Fe1—C7—C8	-120.6 (3)
Br1—Pd1—C1—C5	-117.7 (4)	C6—C7—C8—C9	1.1 (5)
N1—Pd1—C1—C2	-22.0 (2)	Fe1—C7—C8—C9	-59.3 (3)
P1—Pd1—C1—C2	159.5 (2)	C6—C7—C8—Fe1	60.4 (3)
Br1—Pd1—C1—C2	48.1 (4)	C2—Fe1—C8—C9	-119.6 (4)
N1—Pd1—C1—Fe1	56.1 (2)	C3—Fe1—C8—C9	-76.7 (4)
P1—Pd1—C1—Fe1	-122.3 (2)	C7—Fe1—C8—C9	119.4 (5)
Br1—Pd1—C1—Fe1	126.2 (2)	C10—Fe1—C8—C9	38.9 (3)
C2—Fe1—C1—C5	118.0 (3)	C4—Fe1—C8—C9	-43.8 (7)
C3—Fe1—C1—C5	80.6 (2)	C6—Fe1—C8—C9	82.4 (4)
C9—Fe1—C1—C5	165.1 (5)	C5—Fe1—C8—C9	164.7 (5)
C7—Fe1—C1—C5	-121.4 (3)	C1—Fe1—C8—C9	-162.6 (3)
C10—Fe1—C1—C5	-48.4 (6)	C2—Fe1—C8—C7	121.0 (3)
C8—Fe1—C1—C5	-162.8 (3)	C3—Fe1—C8—C7	163.9 (3)
C4—Fe1—C1—C5	37.1 (2)	C9—Fe1—C8—C7	-119.4 (5)
C6—Fe1—C1—C5	-80.5 (3)	C10—Fe1—C8—C7	-80.5 (4)
C3—Fe1—C1—C2	-37.4 (2)	C4—Fe1—C8—C7	-163.2 (5)
C9—Fe1—C1—C2	47.1 (6)	C6—Fe1—C8—C7	-37.0 (3)
C7—Fe1—C1—C2	120.6 (3)	C5—Fe1—C8—C7	45.3 (7)
C10—Fe1—C1—C2	-166.4 (5)	C1—Fe1—C8—C7	78.0 (4)
C8—Fe1—C1—C2	79.3 (3)	C7—C8—C9—C10	-0.7 (5)
C4—Fe1—C1—C2	-80.8 (2)	Fe1—C8—C9—C10	-60.1 (3)
C6—Fe1—C1—C2	161.5 (3)	C7—C8—C9—Fe1	59.3 (3)
C5—Fe1—C1—C2	-118.0 (3)	C2—Fe1—C9—C8	78.9 (4)
C2—Fe1—C1—Pd1	-97.3 (3)	C3—Fe1—C9—C8	121.2 (3)
C3—Fe1—C1—Pd1	-134.7 (3)	C7—Fe1—C9—C8	-37.3 (3)
C9—Fe1—C1—Pd1	-50.2 (6)	C10—Fe1—C9—C8	-118.0 (5)
C7—Fe1—C1—Pd1	23.3 (3)	C4—Fe1—C9—C8	162.6 (3)
C10—Fe1—C1—Pd1	96.3 (5)	C6—Fe1—C9—C8	-80.3 (4)
C8—Fe1—C1—Pd1	-18.1 (3)	C5—Fe1—C9—C8	-164.5 (5)
C4—Fe1—C1—Pd1	-178.2 (3)	C1—Fe1—C9—C8	43.9 (7)
C6—Fe1—C1—Pd1	64.2 (3)	C2—Fe1—C9—C10	-163.1 (3)
C5—Fe1—C1—Pd1	144.7 (4)	C3—Fe1—C9—C10	-120.8 (3)
C5—C1—C2—C3	-0.4 (4)	C7—Fe1—C9—C10	80.6 (4)
Pd1—C1—C2—C3	-171.6 (2)	C8—Fe1—C9—C10	118.0 (5)
Fe1—C1—C2—C3	59.1 (3)	C4—Fe1—C9—C10	-79.4 (4)
C5—C1—C2—C11	-167.1 (3)	C6—Fe1—C9—C10	37.7 (3)
Pd1—C1—C2—C11	21.7 (4)	C5—Fe1—C9—C10	-46.5 (7)
Fe1—C1—C2—C11	-107.6 (3)	C1—Fe1—C9—C10	161.8 (4)
C5—C1—C2—Fe1	-59.5 (2)	C7—C6—C10—C9	0.6 (5)
Pd1—C1—C2—Fe1	129.34 (18)	Fe1—C6—C10—C9	59.7 (3)

supplementary materials

C9—Fe1—C2—C3	78.2 (3)	C7—C6—C10—Fe1	−59.2 (3)
C7—Fe1—C2—C3	161.0 (3)	C8—C9—C10—C6	0.1 (5)
C10—Fe1—C2—C3	45.6 (6)	Fe1—C9—C10—C6	−60.4 (3)
C8—Fe1—C2—C3	119.7 (3)	C8—C9—C10—Fe1	60.5 (3)
C4—Fe1—C2—C3	−38.1 (2)	C2—Fe1—C10—C6	162.0 (5)
C6—Fe1—C2—C3	−167.7 (5)	C3—Fe1—C10—C6	−163.9 (3)
C5—Fe1—C2—C3	−82.0 (2)	C9—Fe1—C10—C6	118.3 (5)
C1—Fe1—C2—C3	−120.1 (3)	C7—Fe1—C10—C6	37.0 (3)
C3—Fe1—C2—C1	120.1 (3)	C8—Fe1—C10—C6	80.4 (4)
C9—Fe1—C2—C1	−161.7 (3)	C4—Fe1—C10—C6	−121.7 (3)
C7—Fe1—C2—C1	−78.9 (3)	C5—Fe1—C10—C6	−79.4 (4)
C10—Fe1—C2—C1	165.7 (5)	C1—Fe1—C10—C6	−44.0 (6)
C8—Fe1—C2—C1	−120.2 (3)	C2—Fe1—C10—C9	43.7 (7)
C4—Fe1—C2—C1	82.0 (2)	C3—Fe1—C10—C9	77.8 (4)
C6—Fe1—C2—C1	−47.6 (6)	C7—Fe1—C10—C9	−81.3 (4)
C5—Fe1—C2—C1	38.06 (19)	C8—Fe1—C10—C9	−37.9 (3)
C3—Fe1—C2—C11	−125.7 (4)	C4—Fe1—C10—C9	120.0 (3)
C9—Fe1—C2—C11	−47.5 (4)	C6—Fe1—C10—C9	−118.3 (5)
C7—Fe1—C2—C11	35.3 (4)	C5—Fe1—C10—C9	162.3 (3)
C10—Fe1—C2—C11	−80.1 (6)	C1—Fe1—C10—C9	−162.3 (4)
C8—Fe1—C2—C11	−6.0 (3)	C12—N2—C11—N1	5.3 (6)
C4—Fe1—C2—C11	−163.8 (3)	C12—N2—C11—C2	−170.8 (3)
C6—Fe1—C2—C11	66.6 (6)	C14—N1—C11—N2	−10.9 (5)
C5—Fe1—C2—C11	152.2 (3)	Pd1—N1—C11—N2	168.0 (3)
C1—Fe1—C2—C11	114.2 (3)	C14—N1—C11—C2	165.5 (3)
C1—C2—C3—C4	0.9 (4)	Pd1—N1—C11—C2	−15.6 (4)
C11—C2—C3—C4	165.6 (4)	C3—C2—C11—N2	10.2 (6)
Fe1—C2—C3—C4	60.5 (3)	C1—C2—C11—N2	173.6 (3)
C1—C2—C3—Fe1	−59.7 (2)	Fe1—C2—C11—N2	93.2 (4)
C11—C2—C3—Fe1	105.1 (4)	C3—C2—C11—N1	−166.4 (4)
C2—Fe1—C3—C4	−118.1 (3)	C1—C2—C11—N1	−3.0 (5)
C9—Fe1—C3—C4	121.5 (3)	Fe1—C2—C11—N1	−83.4 (3)
C7—Fe1—C3—C4	−167.6 (5)	C11—N2—C12—C13	3.9 (6)
C10—Fe1—C3—C4	79.4 (3)	C11—N2—C12—C16	−179.0 (4)
C8—Fe1—C3—C4	162.4 (3)	N2—C12—C13—C14	−7.1 (7)
C6—Fe1—C3—C4	49.4 (6)	C16—C12—C13—C14	175.9 (4)
C5—Fe1—C3—C4	−37.2 (2)	C11—N1—C14—C13	7.0 (6)
C1—Fe1—C3—C4	−80.5 (2)	Pd1—N1—C14—C13	−171.6 (3)
C9—Fe1—C3—C2	−120.3 (3)	C11—N1—C14—C15	−168.0 (4)
C7—Fe1—C3—C2	−49.5 (6)	Pd1—N1—C14—C15	13.5 (6)
C10—Fe1—C3—C2	−162.5 (3)	C12—C13—C14—N1	1.3 (7)
C8—Fe1—C3—C2	−79.5 (3)	C12—C13—C14—C15	176.2 (5)
C4—Fe1—C3—C2	118.1 (3)	C29—P1—C17—C22	−87.5 (3)
C6—Fe1—C3—C2	167.5 (5)	C23—P1—C17—C22	163.6 (3)
C5—Fe1—C3—C2	81.0 (2)	Pd1—P1—C17—C22	43.2 (3)
C1—Fe1—C3—C2	37.6 (2)	C29—P1—C17—C18	148.7 (3)
C2—C3—C4—C5	−1.0 (4)	C23—P1—C17—C18	39.9 (3)
Fe1—C3—C4—C5	58.8 (3)	Pd1—P1—C17—C18	−80.5 (3)
C2—C3—C4—Fe1	−59.8 (2)	C22—C17—C18—C19	57.1 (4)

C2—Fe1—C4—C3	38.7 (2)	P1—C17—C18—C19	−179.1 (3)
C9—Fe1—C4—C3	−76.8 (3)	C17—C18—C19—C20	−58.0 (5)
C7—Fe1—C4—C3	167.4 (5)	C18—C19—C20—C21	56.7 (5)
C10—Fe1—C4—C3	−119.6 (3)	C19—C20—C21—C22	−54.9 (5)
C8—Fe1—C4—C3	−44.7 (6)	C18—C17—C22—C21	−56.3 (4)
C6—Fe1—C4—C3	−160.9 (3)	P1—C17—C22—C21	177.7 (3)
C5—Fe1—C4—C3	120.3 (3)	C20—C21—C22—C17	55.5 (5)
C1—Fe1—C4—C3	83.4 (2)	C29—P1—C23—C28	71.5 (3)
C2—Fe1—C4—C5	−81.6 (2)	C17—P1—C23—C28	−173.0 (3)
C3—Fe1—C4—C5	−120.3 (3)	Pd1—P1—C23—C28	−50.7 (3)
C9—Fe1—C4—C5	163.0 (3)	C29—P1—C23—C24	−57.1 (3)
C7—Fe1—C4—C5	47.1 (6)	C17—P1—C23—C24	58.3 (3)
C10—Fe1—C4—C5	120.2 (3)	Pd1—P1—C23—C24	−179.3 (2)
C8—Fe1—C4—C5	−164.9 (5)	C28—C23—C24—C25	58.1 (4)
C6—Fe1—C4—C5	78.8 (3)	P1—C23—C24—C25	−171.8 (3)
C1—Fe1—C4—C5	−36.9 (2)	C23—C24—C25—C26	−57.2 (5)
C2—C1—C5—C4	−0.2 (4)	C24—C25—C26—C27	56.0 (5)
Pd1—C1—C5—C4	166.0 (3)	C25—C26—C27—C28	−56.1 (5)
Fe1—C1—C5—C4	−58.7 (3)	C26—C27—C28—C23	57.9 (5)
C2—C1—C5—Fe1	58.5 (2)	C24—C23—C28—C27	−58.8 (4)
Pd1—C1—C5—Fe1	−135.3 (4)	P1—C23—C28—C27	169.4 (3)
C3—C4—C5—C1	0.7 (4)	C17—P1—C29—C30	169.9 (2)
Fe1—C4—C5—C1	59.0 (2)	C23—P1—C29—C30	−78.0 (3)
C3—C4—C5—Fe1	−58.3 (3)	Pd1—P1—C29—C30	40.4 (3)
C2—Fe1—C5—C1	−38.9 (2)	C17—P1—C29—C34	−18.8 (4)
C3—Fe1—C5—C1	−83.4 (2)	C23—P1—C29—C34	93.4 (3)
C9—Fe1—C5—C1	−164.6 (5)	Pd1—P1—C29—C34	−148.2 (3)
C7—Fe1—C5—C1	77.6 (3)	C34—C29—C30—C31	−4.5 (5)
C10—Fe1—C5—C1	160.6 (3)	P1—C29—C30—C31	168.2 (3)
C8—Fe1—C5—C1	44.3 (6)	C29—C30—C31—C32	0.4 (6)
C4—Fe1—C5—C1	−120.4 (3)	C30—C31—C32—C33	2.3 (7)
C6—Fe1—C5—C1	118.7 (3)	C31—C32—C33—C34	−0.8 (7)
C2—Fe1—C5—C4	81.4 (3)	C32—C33—C34—C29	−3.4 (6)
C3—Fe1—C5—C4	37.0 (2)	C32—C33—C34—C35	177.1 (4)
C9—Fe1—C5—C4	−44.3 (6)	C30—C29—C34—C33	5.8 (5)
C7—Fe1—C5—C4	−162.0 (3)	P1—C29—C34—C33	−165.1 (3)
C10—Fe1—C5—C4	−79.0 (3)	C30—C29—C34—C35	−174.8 (3)
C8—Fe1—C5—C4	164.6 (5)	P1—C29—C34—C35	14.3 (6)
C6—Fe1—C5—C4	−120.9 (3)	C33—C34—C35—C40	80.8 (5)
C1—Fe1—C5—C4	120.4 (3)	C29—C34—C35—C40	−98.7 (5)
C2—Fe1—C6—C7	−42.9 (6)	C33—C34—C35—C36	−94.5 (5)
C3—Fe1—C6—C7	160.3 (4)	C29—C34—C35—C36	86.0 (5)
C9—Fe1—C6—C7	80.9 (4)	C42—N3—C36—C37	−113.7 (5)
C10—Fe1—C6—C7	119.5 (5)	C43—N3—C36—C37	18.8 (6)
C8—Fe1—C6—C7	37.4 (3)	C42—N3—C36—C35	67.7 (5)
C4—Fe1—C6—C7	−163.2 (3)	C43—N3—C36—C35	−159.8 (4)
C5—Fe1—C6—C7	−120.8 (3)	C40—C35—C36—C37	3.1 (6)
C1—Fe1—C6—C7	−78.3 (3)	C34—C35—C36—C37	178.4 (4)
C2—Fe1—C6—C10	−162.4 (5)	C40—C35—C36—N3	−178.3 (4)

supplementary materials


C3—Fe1—C6—C10	40.7 (7)	C34—C35—C36—N3	-3.0 (6)
C9—Fe1—C6—C10	-38.7 (4)	N3—C36—C37—C38	179.6 (4)
C7—Fe1—C6—C10	-119.5 (5)	C35—C36—C37—C38	-1.7 (7)
C8—Fe1—C6—C10	-82.1 (4)	C36—C37—C38—C39	-0.5 (8)
C4—Fe1—C6—C10	77.3 (4)	C37—C38—C39—C40	1.3 (8)
C5—Fe1—C6—C10	119.7 (3)	C38—C39—C40—C35	0.1 (7)
C1—Fe1—C6—C10	162.2 (3)	C36—C35—C40—C39	-2.3 (6)
C10—C6—C7—C8	-1.0 (5)	C34—C35—C40—C39	-177.9 (4)
Fe1—C6—C7—C8	-60.1 (3)		

Hydrogen-bond geometry (\AA , $^\circ$)

$D\text{—H}\cdots A$	$D\text{—H}$	$H\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
C3—H3 ⁱ —Cl2 ⁱ	0.98	2.78	3.671 (5)	151
C16—H16C ⁱⁱ —Cg1 ⁱⁱ	0.96	2.68	3.544 (6)	151

Symmetry codes: (i) $-x+1, -y+1, -z$; (ii) $-x+1, -y+2, -z$.

Fig. 1



supplementary materials

Fig. 2

