

Poly[tetraqua- μ_3 -benzene-1,2-di-carboxylato- μ_3 -bromido-penta- μ_2 -bromido-octa- μ_3 -isonicotinato-hepta-copper(I)trilanthanum(III)]

Guo-Ming Wang,* Zeng-Xin Li, Shu-Yun Xue and Hui-Luan Liu

Department of Chemistry, Teachers' College of Qingdao University, Shandong 266071, People's Republic of China

Correspondence e-mail: gmwang_pub@163.com


Received 7 April 2009; accepted 15 April 2009

Key indicators: single-crystal X-ray study; $T = 295\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.012\text{ \AA}$; disorder in main residue; R factor = 0.052; wR factor = 0.126; data-to-parameter ratio = 14.0.

A new lanthanum(III)-copper(I) heterometallic coordination polymer, $[\text{Cu}_7\text{La}_3\text{Br}_6(\text{C}_6\text{H}_4\text{NO}_2)_8(\text{C}_8\text{H}_4\text{O}_4)(\text{H}_2\text{O})_4]_n$, has been prepared by a hydrothermal method. Of the three La atoms in the asymmetric unit, two are eight-coordinate with bicapped trigonal-prismatic configurations; the third is nine-coordinated and has a tricapped trigonal-prismatic coordination geometry. Of the seven Cu atoms, two are two-coordinate with CuBrN and CuN_2 ligand sets, three have trigonal configurations, *viz.* CuBrN_2 , CuBr_2N and CuBr_3 , while the remaining two adopt distorted tetrahedral CuBr_3N geometries. In the crystal structure, adjacent La centers are linked by isonicotinate (IN^-) and benzene-1,2-dicarboxylate ligands to form a two-dimensional La-carboxylate layer in the *ab* plane. These layers are further interconnected with each other by bridging $[\text{Cu}(\text{IN})_2]$ motifs, leading to an unusual three-dimensional heterometallic Cu-halide-lanthanide-organic framework, with the inorganic $[\text{Cu}_6\text{Br}_6]_n$ chains located in the resulting channels. Two Cu atoms are disordered over two positions, both with site occupancy factors of 0.80 and 0.20. O-H-O hydrogen bonding between water molecules and carboxylate O atoms helps to consolidate the crystal packing.


Related literature

For background on the structures and applications of heterometallic lanthanide-transition metal (Ln-TM) coordination polymers, see: Benelli & Gatteschi (2002); Shibasaki & Yoshikawa (2002); Zhao, Cheng *et al.* (2004); Zhao, Chen *et al.* (2004); Guillou *et al.* (2006); Wang *et al.* (2006). For some examples of extended heterometallic Ln-TM architectures, see: Ren *et al.* (2003); Prasad *et al.* (2007); Cheng *et al.* (2008); Deng *et al.* (2008); Wang, Li *et al.* (2008). For the coordination modes of isonicotinate and benzene-1,2-dicarboxylate ligands, see: Gu & Xue (2007); Wang, Duan *et al.* (2008).



Experimental

Crystal data



$M_r = 2553.96$

Monoclinic, $P2_1/c$

$a = 10.1071(5)\text{ \AA}$

$b = 19.6311(3)\text{ \AA}$

$c = 34.4015(2)\text{ \AA}$

$\beta = 92.480(2)^\circ$

$V = 6819.3(4)\text{ \AA}^3$

$Z = 4$

Mo $K\alpha$ radiation

$\mu = 7.57\text{ mm}^{-1}$

$T = 295\text{ K}$

$0.20 \times 0.10 \times 0.09\text{ mm}$

Data collection

Bruker APEXII area-detector diffractometer

Absorption correction: multi-scan (*SADABS*; Sheldrick, 1996)

$T_{\min} = 0.313$, $T_{\max} = 0.549$

(expected range = 0.288–0.506)

52743 measured reflections

13363 independent reflections

9870 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.071$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.052$

$wR(F^2) = 0.126$

$S = 1.18$

13363 reflections

955 parameters

H-atom parameters constrained

$\Delta\rho_{\max} = 1.73\text{ e \AA}^{-3}$

$\Delta\rho_{\min} = -2.22\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O16-H16D \cdots O21 ⁱ	0.86	2.09	2.901 (7)	157
O23-H23D \cdots O22	0.93	2.00	2.861 (8)	153
O24-H24D \cdots O20	0.85	2.22	2.844 (11)	130

Symmetry code: (i) $x + 1, y, z$.

Data collection: *APEX2* (Bruker, 2002); cell refinement: *SAINT* (Bruker, 2002); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXL97*.

This work was supported by the Qingdao University Research Fund (grant No. 063-06300522).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SJ2614).

References

- Benelli, C. & Gatteschi, D. (2002). *Chem. Rev.* **102**, 2369–2388.
- Bruker (2002). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Cheng, J. W., Zhang, J., Zheng, S. T. & Yang, G. Y. (2008). *Chem. Eur. J.* **14**, 88–97.
- Deng, H., Li, Y. H., Qiu, C. Y., Liu, Z. H. & Zeller, M. (2008). *Inorg. Chem. Commun.* **11**, 1151–1154.
- Gu, X. J. & Xue, D. F. (2007). *Inorg. Chem.* **46**, 3212–3216.
- Guillou, O., Daiguebonne, C., Camara, M. & Kerbellec, N. (2006). *Inorg. Chem.* **45**, 8468–8470.
- Prasad, T. K., Rajasekharan, M. V. & Costes, J. P. (2007). *Angew. Chem. Int. Ed.* **46**, 2851–2854.
- Ren, Y., Long, L., Mao, B., Yuan, Y., Huang, R. & Zheng, L. (2003). *Angew. Chem. Int. Ed.* **42**, 532–535.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Shibasaki, M. & Yoshikawa, N. (2002). *Chem. Rev.* **102**, 2187–2210.
- Wang, G.-M., Duan, C.-S., Liu, H.-L. & Li, H. (2008). *Acta Cryst. E* **64**, m468–m469.
- Wang, G.-M., Li, Z.-X., Zheng, Q.-H. & Liu, H.-L. (2008). *Acta Cryst. E* **64**, m1260–m1261.
- Wang, Z., Shen, X., Wang, J., Zhang, P., Li, Y., Nfor, E., Song, Y., Ohkoshi, S., Hashimoto, K. & You, X. (2006). *Angew. Chem. Int. Ed.* **45**, 3287–3291.
- Zhao, B., Chen, X., Cheng, P., Liao, D., Yan, S. & Jiang, Z. (2004). *J. Am. Chem. Soc.* **126**, 15394–15395.
- Zhao, B., Cheng, P., Chen, X., Cheng, C., Shi, W., Liao, D., Yan, S. & Jiang, Z. (2004). *J. Am. Chem. Soc.* **126**, 3012–3013.

supplementary materials

Acta Cryst. (2009). E65, m550-m551 [doi:10.1107/S1600536809014081]

Poly[tetraaqua- μ_3 -benzene-1,2-dicarboxylato- μ_3 -bromido-penta- μ_2 -bromido-octa- μ_3 -isonicotinato-heptacopper(I)trilanthanum(III)]

G.-M. Wang, Z.-X. Li, S.-Y. Xue and H.-L. Liu

Comment

In recent years, the design and construction of heterometallic lanthanide(Ln)-transition metal(TM) coordination frameworks have attracted considerable attention because of their intriguing topological architectures and potential applications in for example magnetism, luminescence, and heterogeneous catalysis (Benelli & Gatteschi, 2002; Shibasaki & Yoshikawa, 2002; Zhao, Cheng *et al.*, 2004; Zhao, Chen *et al.*, 2004; Guillou *et al.*, 2006; Wang *et al.*, 2006). Compared with the assembly of homometallic Ln and TM compounds, the analogous chemistry and synthetic strategy of heterometallic Ln—TM coordination frameworks is still underdeveloped. (Ren *et al.*, 2003; Prasad *et al.*, 2007; Cheng *et al.*, 2008; Deng *et al.* (2008); Wang, Li *et al.* (2008)). This may be attributed to the variable and versatile coordination numbers of the lanthanide ions, their low stereochemical preference, as well as the competitive reactions of Ln and TM metals for the same organic ligands. Fortunately, according to the hard-soft acid base theory, the Ln and TM ions have different affinities for O and N donors, which makes it possible to construct unusual heterometallic Ln—TM frameworks by choosing multifunctional ligands with both oxygen and nitrogen donors. Therefore, isonicotinic acid (HIN) has been chosen here as the bifunctional ligand. Meanwhile, we also introduced another multifunctional ligand, the deprotonated 1,2-benzenedicarboxylic acid, (BDC^{2-}), into the reaction system simultaneously, exploring the construction of heterometallic Ln—TM compounds with high dimensionality. The title compound $[La_3Cu_7Br_6(IN)_8(BDC)(H_2O)_4]_n$ (1) is reported here and displays novel three-dimensional coordination features.

As shown in Fig. 1, the asymmetric unit contains three unique lanthanum(III) atoms, seven copper(I) ions, six bromide ions, one BDC^{2-} ligand and eight IN^- ligands, as well as four aqua ligands. The La1 and La2 atoms are both eight-coordinate with bicapped trigonal prismatic geometries. Atom La1 is surrounded by six carboxylate oxygen atoms from six IN^- ligands, one carboxylate oxygen atom from a BDC^{2-} ligand and an aqua ligand. Atom La2, on the other hand, is coordinated by five carboxylate oxygen atoms from five IN^- ligands, two carboxylate oxygen atoms from a BDC^{2-} ligand and an aqua ligand. The La3 atom is nine-coordinated and has a tricapped trigonal-prismatic coordination environment comprising two coordinated water molecules, five carboxylate oxygen atoms from five IN^- ligands and two carboxylate oxygen atoms from one BDC^{2-} ligand. The La—O bond lengths range from 2.387 (6) to 2.797 (6) Å.

The identical La(III) ions are linked by mixed IN^- and BDC^{2-} ligands to form a two-dimensional La-carboxylate layer in the *ab* plane (Fig. 2). Interestingly, compared to the abundant and versatile coordination modes that found in IN^- and BDC^{2-} ligands (Gu & Xue, 2007; Wang, Duan *et al.*, 2008), only a single bidentate (for IN^-) and a unique pentadentate (for BDC^{2-}) modes are adopted in the La-carboxylate layer (Fig. 3). These La-carboxylate layers are further interconnected by $[Cu(1)(IN)_2]$ linear bridging to give rise to an unusual Cu-halide-lanthanide-organic framework with one-dimensional channels (Fig. 4), in which the inorganic $[Cu_6Br_6]_n$ chains are located. As shown in Fig. 5, the inorganic $[Cu_6Br_6]_n$ motif contains six unique Cu(I) atoms with three different types of coordination modes and six Br^- ions. The Cu2 atom is two-co-

supplementary materials

ordinated with a nerally linear geometry: one $\mu_2\text{-Br}1$ ion and one N atom from one bridging IN⁻ ligand. The Cu3, Cu4 and Cu5 atoms are three-coordinate with trigonal coordination environments: two isonicotinate N atoms and one $\mu_2\text{-Br}2$ ion are bonded to Cu3; one $\mu_2\text{-Br}2$ ion, one $\mu_3\text{-Br}3$ ion and one $\mu_3\text{-Br}4$ ion to Cu4; one $\mu_2\text{-Br}5$ ion, one $\mu_3\text{-Br}4$ ion and one N atom to Cu5. The remaining Cu6 and Cu7 atoms, however, are coordinated to one N atom and three $\mu_2\text{-Br}$ (Br3, Br5 and Br6 for Cu6; Br1, Br4 and Br6 for Cu7) ions respectively, defining distorted tetrahedral geometries. The Cu—N and Cu—Br distances are in the range 1.921 (7)–2.031 (8) Å and 2.228 (2)–2.670 (2) Å, respectively. Therefore, the overall structure of 1 can also be viewed as one-dimensional [Cu₆Br₆]_n chains inserted into the channels of a three-dimensional heterometallic Cu-halide-lanthanide-organic framework (Fig. 6).

Experimental

The title compound was synthesized under mild hydrothermal conditions. Typically, a mixture of La₂O₃ (0.5 mmol; 0.163 g), CuBr₂ (0.067 g, 0.30 mmol), HIN (2.00 mmol, 0247 g), H₂BDC (1.00 mmol, 0.167 g) and H₂O (8 ml) was sealed in a 25 ml Teflon-lined steel autoclave and heated under autogenous pressure at 443 K for 9 days. The brown prism-like crystals obtained were recovered by filtration, washed with distilled water and dried in air. Although copper(II) salts were used as starting materials, the Cu centers in the product are in the +1 oxidation state. This is attributed to a reduction reaction occurring under the hydrothermal conditions used.

Refinement

H atoms bound to C atoms were positioned geometrically, with C—H distances of 0.93 Å, and constrained to ride on their parent atoms [$U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$]. H atoms bound to O atoms were located in a difference Fourier map and treated as riding, with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{O})$. Atoms Cu2 and Cu3 were refined as disordered over two positions, with site occupancy factors of fixed at 0.80 and 0.20 respectively in the final refinement.

Figures




Fig. 1. The molecular structure of 1, showing the atom-numbering scheme. Displacement ellipsoids are drawn at the 50% probability level. [Symmetry codes: (i) -1 + x , y , z ; (ii) - x , 1 - y , - z ; (iii) 1 - x , 1 - y , - z ; (iv) - x , - y , - z ; (v) 1 + x , 1/2 - y , -1/2 + z ; (vi) 1 + x , y , z ; (vii) 1 - x , 1/2 + y , -1/2 - z ; (viii) 1 - x , -1/2 + y , -1/2 - z .]




Fig. 2. Two-dimensional La-carboxylate layer in the *ab* plane.




Fig. 3. The coordination modes of IN^- and BDC^{2-} found in the La-carboxylate layer.




Fig. 4. A view of the three-dimensional Cu-halide-lanthanide-organic framework with one-dimensional channels.




Fig. 5. One-dimensional infinite $[\text{Cu}_6\text{Br}_6]_n$ chains along the a axis.




Fig. 6. Framework of 1 viewed along the a axis.

Poly[tetraqua- μ_3 -benzene-1,2-dicarboxylato- μ_3 -bromido-penta- μ_2 -bromido-octa- μ_3 -isonicotinato-heptacopper(I)trilanthanum(III)]

Crystal data

$[\text{Cu}_7\text{La}_3\text{Br}_6(\text{C}_6\text{H}_4\text{NO}_2)_8(\text{C}_8\text{H}_4\text{O}_4)(\text{H}_2\text{O})_4]$	$F_{000} = 4848$
$M_r = 2553.96$	$D_x = 2.488 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
Hall symbol: -P 2ybc	$\lambda = 0.71073 \text{ \AA}$
$a = 10.1071 (5) \text{ \AA}$	Cell parameters from 52743 reflections
$b = 19.6311 (3) \text{ \AA}$	$\theta = 1.2\text{--}26.0^\circ$
$c = 34.4015 (2) \text{ \AA}$	$\mu = 7.57 \text{ mm}^{-1}$
$\beta = 92.480 (2)^\circ$	$T = 295 \text{ K}$
$V = 6819.3 (4) \text{ \AA}^3$	Prism, brown
$Z = 4$	$0.20 \times 0.10 \times 0.09 \text{ mm}$

Data collection

Bruker APEXII area-detector diffractometer	13363 independent reflections
Radiation source: fine-focus sealed tube	9870 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.071$
$T = 295 \text{ K}$	$\theta_{\text{max}} = 26.0^\circ$
ϕ and ω scans	$\theta_{\text{min}} = 1.2^\circ$
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	$h = -12 \rightarrow 12$
$T_{\text{min}} = 0.313$, $T_{\text{max}} = 0.549$	$k = -23 \rightarrow 24$

supplementary materials

52743 measured reflections

$l = -42 \rightarrow 42$

Refinement

Refinement on F^2

Secondary atom site location: difference Fourier map

Least-squares matrix: full

Hydrogen site location: inferred from neighbouring sites

$R[F^2 > 2\sigma(F^2)] = 0.052$

H-atom parameters constrained

$wR(F^2) = 0.126$

$$w = 1/[\sigma^2(F_o^2) + (0.0498P)^2 + 4.7192P]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$S = 1.18$

$$(\Delta/\sigma)_{\max} = 0.001$$

13363 reflections

$$\Delta\rho_{\max} = 1.73 \text{ e \AA}^{-3}$$

955 parameters

$$\Delta\rho_{\min} = -2.22 \text{ e \AA}^{-3}$$

Primary atom site location: structure-invariant direct methods

Extinction correction: none

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\text{sigma}(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R- factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
La1	0.22600 (4)	0.48720 (2)	0.006803 (12)	0.01347 (11)	
La2	0.29679 (4)	0.14367 (2)	-0.026957 (12)	0.01318 (11)	
La3	-0.24624 (4)	0.16813 (2)	-0.017184 (13)	0.01663 (12)	
Cu1	0.38381 (16)	0.52109 (8)	-0.24321 (3)	0.0605 (4)	
Cu2	0.1352 (4)	0.2896 (2)	-0.21115 (11)	0.0565 (9)	0.80
Cu2'	0.0878 (17)	0.3052 (9)	-0.2218 (4)	0.074 (5)	0.20
Cu3	0.8350 (3)	0.1565 (2)	-0.26477 (12)	0.0440 (7)	0.80
Cu3'	0.8908 (14)	0.1708 (11)	-0.2644 (5)	0.064 (4)	0.20
Cu4	0.63663 (16)	0.28675 (8)	-0.26968 (5)	0.0653 (4)	
Cu5	0.61139 (14)	0.41101 (7)	-0.23317 (4)	0.0530 (4)	
Cu6	0.88795 (15)	0.43063 (7)	-0.27424 (3)	0.0514 (4)	
Cu7	1.21736 (16)	0.34075 (8)	-0.28116 (4)	0.0576 (4)	
Br1	0.21910 (10)	0.22413 (5)	-0.25622 (3)	0.0408 (3)	
Br2	0.57870 (12)	0.17237 (6)	-0.26488 (4)	0.0548 (3)	
Br3	0.85917 (10)	0.30796 (5)	-0.25766 (3)	0.0443 (3)	
Br4	0.47465 (9)	0.36921 (5)	-0.28606 (3)	0.0359 (2)	
Br5	0.76335 (9)	0.50127 (5)	-0.23036 (3)	0.0380 (2)	

Br6	1.13084 (9)	0.44473 (5)	-0.25274 (3)	0.0362 (2)
O1	0.1380 (6)	0.5739 (3)	-0.04055 (17)	0.0325 (15)
O2	-0.0776 (5)	0.5648 (3)	-0.05687 (16)	0.0269 (14)
O3	0.3829 (5)	0.4716 (3)	-0.04459 (15)	0.0279 (14)
O4	0.5802 (6)	0.5205 (3)	-0.05111 (16)	0.0308 (15)
O5	-0.1574 (6)	0.4261 (3)	-0.05461 (18)	0.0383 (17)
O6	0.0482 (6)	0.4339 (3)	-0.03153 (17)	0.0392 (17)
O7	0.3772 (6)	0.6005 (3)	0.00383 (19)	0.0406 (17)
H7A	0.4177	0.5917	-0.0167	0.080*
H7B	0.4321	0.5951	0.0231	0.080*
O8	0.2809 (6)	0.3656 (3)	0.00780 (17)	0.0325 (15)
O9	0.2517 (7)	0.2573 (3)	-0.00553 (19)	0.0432 (18)
O10	0.6667 (6)	0.2366 (4)	0.0380 (2)	0.050 (2)
O11	0.4973 (6)	0.1853 (3)	0.00866 (16)	0.0275 (14)
O12	0.6320 (6)	0.0893 (4)	-0.06027 (17)	0.0433 (18)
O13	0.4174 (5)	0.0698 (3)	-0.07032 (17)	0.0306 (15)
O14	0.6171 (7)	0.2426 (4)	-0.0589 (2)	0.064 (3)
O15	0.4104 (6)	0.2149 (3)	-0.07631 (18)	0.0381 (16)
O16	0.4042 (5)	0.0575 (3)	0.02259 (16)	0.0295 (14)
H16C	0.3978	0.0251	0.0388	0.080*
H16D	0.4832	0.0734	0.0256	0.080*
O17	0.1231 (5)	0.1276 (3)	0.01773 (16)	0.0285 (14)
O18	-0.0733 (6)	0.1657 (4)	0.03231 (17)	0.0381 (17)
O19	0.1224 (6)	0.1580 (3)	-0.07778 (18)	0.0331 (15)
O20	-0.0813 (6)	0.1887 (4)	-0.06543 (17)	0.0381 (16)
O21	-0.3099 (5)	0.0712 (3)	0.02882 (15)	0.0217 (13)
O22	-0.1878 (5)	-0.0234 (3)	0.03169 (15)	0.0197 (12)
O23	-0.1039 (5)	0.0580 (3)	-0.03168 (17)	0.0321 (15)
H23C	-0.0254	0.0630	-0.0290	0.080*
H23D	-0.1202	0.0204	-0.0159	0.080*
O24	-0.1433 (10)	0.2896 (4)	-0.0098 (3)	0.084 (3)
H24C	-0.1207	0.3284	-0.0007	0.101*
H24D	-0.0999	0.2827	-0.0300	0.101*
C1	0.0388 (8)	0.5791 (4)	-0.0635 (2)	0.0238 (19)
C2	0.0667 (7)	0.6024 (4)	-0.1045 (2)	0.0205 (18)
C3	-0.0229 (8)	0.5899 (5)	-0.1351 (2)	0.030 (2)
H3A	-0.1032	0.5689	-0.1305	0.036*
C4	0.0066 (9)	0.6087 (5)	-0.1727 (3)	0.037 (2)
H4A	-0.0560	0.6005	-0.1928	0.044*
C5	0.2079 (10)	0.6501 (5)	-0.1517 (3)	0.040 (3)
H5A	0.2882	0.6708	-0.1567	0.049*
C6	0.1829 (9)	0.6333 (5)	-0.1140 (3)	0.034 (2)
H6A	0.2463	0.6431	-0.0944	0.040*
C7	0.4703 (8)	0.5001 (4)	-0.0643 (2)	0.0206 (18)
C8	0.4435 (8)	0.5066 (4)	-0.1068 (2)	0.0238 (19)
C9	0.3389 (9)	0.4730 (5)	-0.1256 (2)	0.032 (2)
H9A	0.2799	0.4476	-0.1115	0.038*
C10	0.3229 (10)	0.4776 (5)	-0.1654 (3)	0.044 (3)
H10A	0.2540	0.4534	-0.1777	0.052*

supplementary materials

C11	0.5003 (10)	0.5473 (5)	-0.1686 (3)	0.041 (3)
H11A	0.5567	0.5733	-0.1833	0.050*
C12	0.5248 (9)	0.5449 (5)	-0.1292 (2)	0.030 (2)
H12A	0.5955	0.5688	-0.1177	0.035*
C13	-0.0358 (8)	0.4176 (4)	-0.0570 (2)	0.0199 (18)
C14	0.0095 (8)	0.3870 (4)	-0.0943 (2)	0.0205 (18)
C15	-0.0494 (9)	0.4072 (5)	-0.1290 (3)	0.032 (2)
H15A	-0.1170	0.4394	-0.1295	0.038*
C16	-0.0078 (10)	0.3795 (5)	-0.1629 (3)	0.039 (2)
H16A	-0.0463	0.3949	-0.1864	0.047*
C17	0.1388 (9)	0.3108 (5)	-0.1299 (3)	0.039 (3)
H17A	0.2010	0.2759	-0.1301	0.047*
C18	0.1073 (8)	0.3381 (4)	-0.0946 (3)	0.030 (2)
H18A	0.1508	0.3240	-0.0716	0.036*
C19	0.2871 (8)	0.3047 (4)	0.0165 (2)	0.0218 (18)
C20	0.3417 (8)	0.2869 (4)	0.0570 (2)	0.0220 (19)
C21	0.2682 (8)	0.3108 (4)	0.0879 (3)	0.027 (2)
H21A	0.1893	0.3341	0.0827	0.032*
C22	0.3115 (9)	0.3002 (4)	0.1262 (3)	0.032 (2)
H22A	0.2605	0.3151	0.1464	0.038*
C23	0.4283 (9)	0.2680 (5)	0.1342 (3)	0.034 (2)
H23A	0.4574	0.2609	0.1599	0.040*
C24	0.5047 (9)	0.2454 (5)	0.1040 (3)	0.033 (2)
H24A	0.5864	0.2251	0.1098	0.039*
C25	0.4602 (7)	0.2528 (4)	0.0652 (2)	0.0211 (18)
C26	0.5453 (8)	0.2248 (4)	0.0353 (3)	0.0244 (19)
C27	0.5335 (8)	0.0750 (4)	-0.0816 (2)	0.0197 (18)
C28	0.5547 (8)	0.0625 (4)	-0.1237 (2)	0.0201 (18)
C29	0.6764 (8)	0.0441 (5)	-0.1370 (3)	0.032 (2)
H29A	0.7490	0.0395	-0.1196	0.038*
C30	0.6903 (9)	0.0325 (5)	-0.1760 (2)	0.033 (2)
H30A	0.7731	0.0192	-0.1841	0.039*
C31	0.4752 (10)	0.0582 (6)	-0.1901 (3)	0.045 (3)
H31A	0.4060	0.0645	-0.2085	0.054*
C32	0.4512 (9)	0.0690 (5)	-0.1518 (3)	0.038 (2)
H32A	0.3666	0.0805	-0.1444	0.045*
C33	0.5163 (9)	0.2446 (5)	-0.0812 (3)	0.031 (2)
C34	0.5301 (8)	0.2866 (4)	-0.1169 (2)	0.0242 (19)
C35	0.4653 (9)	0.2684 (5)	-0.1515 (3)	0.032 (2)
H35A	0.4074	0.2315	-0.1525	0.038*
C36	0.4877 (9)	0.3059 (5)	-0.1848 (3)	0.037 (2)
H36A	0.4469	0.2921	-0.2083	0.045*
C37	0.6226 (9)	0.3787 (5)	-0.1509 (3)	0.033 (2)
H37A	0.6742	0.4179	-0.1500	0.040*
C38	0.6105 (9)	0.3424 (4)	-0.1166 (3)	0.031 (2)
H38A	0.6563	0.3559	-0.0939	0.038*
C39	0.0442 (8)	0.1505 (4)	0.0412 (2)	0.0208 (18)
C40	0.0887 (7)	0.1565 (4)	0.0830 (2)	0.0195 (18)
C41	0.0109 (8)	0.1899 (4)	0.1090 (2)	0.027 (2)

H41A	-0.0680	0.2106	0.1006	0.032*
C42	0.0531 (9)	0.1919 (5)	0.1480 (3)	0.035 (2)
H42A	0.0019	0.2150	0.1656	0.042*
C43	0.2389 (9)	0.1316 (5)	0.1347 (3)	0.033 (2)
H43A	0.3188	0.1123	0.1434	0.040*
C44	0.2062 (8)	0.1270 (5)	0.0962 (3)	0.030 (2)
H44A	0.2609	0.1048	0.0792	0.036*
C45	0.0064 (8)	0.1715 (4)	-0.0876 (2)	0.0199 (18)
C46	-0.0323 (8)	0.1683 (4)	-0.1304 (2)	0.0214 (18)
C47	-0.1537 (8)	0.1923 (4)	-0.1445 (2)	0.027 (2)
H47A	-0.2133	0.2114	-0.1277	0.033*
C48	-0.1848 (9)	0.1875 (5)	-0.1837 (3)	0.033 (2)
H48A	-0.2639	0.2070	-0.1930	0.040*
C49	0.0058 (9)	0.1337 (5)	-0.1953 (2)	0.032 (2)
H49A	0.0608	0.1126	-0.2126	0.038*
C50	0.0497 (8)	0.1386 (4)	-0.1569 (3)	0.027 (2)
H50A	0.1330	0.1224	-0.1489	0.032*
C51	-0.2364 (7)	0.0287 (4)	0.0465 (2)	0.0179 (17)
C52	-0.2047 (7)	0.0423 (4)	0.0886 (2)	0.0179 (17)
C53	-0.2863 (8)	0.0798 (5)	0.1118 (3)	0.033 (2)
H53A	-0.3620	0.1001	0.1007	0.039*
C54	-0.2575 (10)	0.0875 (5)	0.1509 (3)	0.040 (3)
H54A	-0.3163	0.1113	0.1659	0.048*
C55	-0.0661 (9)	0.0266 (5)	0.1462 (3)	0.035 (2)
H55A	0.0116	0.0091	0.1577	0.042*
C56	-0.0915 (8)	0.0148 (5)	0.1070 (2)	0.029 (2)
H56A	-0.0334	-0.0114	0.0930	0.035*
N1	0.1216 (7)	0.6380 (4)	-0.1813 (2)	0.0334 (19)
N2	0.4013 (8)	0.5150 (4)	-0.1872 (2)	0.040 (2)
N3	0.0842 (9)	0.3320 (4)	-0.1638 (2)	0.043 (2)
N4	0.5940 (8)	0.0390 (4)	-0.2026 (2)	0.0356 (19)
N5	0.5652 (7)	0.3610 (4)	-0.1848 (2)	0.0315 (18)
N6	0.1655 (8)	0.1616 (4)	0.1610 (2)	0.0346 (19)
N7	-0.1092 (8)	0.1570 (4)	-0.2089 (2)	0.0346 (19)
N8	-0.1468 (8)	0.0616 (4)	0.1681 (2)	0.0337 (19)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
La1	0.0140 (2)	0.0170 (2)	0.0094 (2)	0.00099 (17)	0.00035 (17)	0.00074 (18)
La2	0.0137 (2)	0.0148 (2)	0.0109 (2)	0.00006 (18)	-0.00093 (17)	-0.00007 (18)
La3	0.0145 (2)	0.0225 (3)	0.0126 (2)	0.00030 (19)	-0.00268 (17)	0.00349 (19)
Cu1	0.0995 (12)	0.0690 (11)	0.0132 (6)	0.0036 (9)	0.0041 (7)	0.0086 (6)
Cu2	0.075 (2)	0.055 (2)	0.041 (2)	-0.0055 (15)	0.0266 (14)	-0.0164 (14)
Cu2'	0.114 (13)	0.076 (9)	0.037 (8)	0.019 (8)	0.046 (8)	-0.005 (6)
Cu3	0.058 (2)	0.0613 (18)	0.0126 (10)	0.0103 (15)	-0.0006 (14)	-0.0047 (10)
Cu3'	0.093 (12)	0.086 (12)	0.014 (4)	0.031 (9)	-0.001 (8)	-0.006 (6)
Cu4	0.0700 (10)	0.0659 (11)	0.0600 (10)	-0.0001 (8)	0.0015 (8)	0.0028 (8)

supplementary materials

Cu5	0.0697 (9)	0.0580 (9)	0.0309 (7)	-0.0293 (8)	-0.0008 (6)	0.0128 (6)
Cu6	0.0762 (10)	0.0571 (9)	0.0201 (7)	0.0036 (7)	-0.0047 (6)	0.0050 (6)
Cu7	0.0823 (11)	0.0660 (10)	0.0236 (7)	-0.0031 (8)	-0.0095 (7)	0.0041 (6)
Br1	0.0451 (6)	0.0387 (6)	0.0395 (6)	-0.0013 (5)	0.0128 (5)	-0.0057 (5)
Br2	0.0543 (7)	0.0558 (8)	0.0539 (7)	-0.0024 (6)	-0.0033 (6)	-0.0041 (6)
Br3	0.0484 (6)	0.0371 (6)	0.0475 (6)	0.0005 (5)	0.0032 (5)	0.0082 (5)
Br4	0.0291 (5)	0.0463 (6)	0.0317 (5)	-0.0051 (4)	-0.0062 (4)	0.0062 (5)
Br5	0.0354 (5)	0.0380 (6)	0.0405 (6)	-0.0063 (4)	0.0006 (4)	-0.0055 (5)
Br6	0.0335 (5)	0.0523 (7)	0.0230 (5)	-0.0105 (4)	0.0018 (4)	-0.0016 (4)
O1	0.029 (3)	0.042 (4)	0.026 (3)	-0.003 (3)	-0.006 (3)	0.020 (3)
O2	0.016 (3)	0.034 (4)	0.030 (3)	0.003 (3)	0.007 (3)	0.016 (3)
O3	0.028 (3)	0.042 (4)	0.014 (3)	-0.007 (3)	0.010 (2)	0.000 (3)
O4	0.028 (3)	0.041 (4)	0.022 (3)	-0.009 (3)	-0.009 (3)	0.003 (3)
O5	0.039 (4)	0.037 (4)	0.039 (4)	-0.002 (3)	0.005 (3)	-0.025 (3)
O6	0.054 (4)	0.039 (4)	0.024 (4)	-0.004 (3)	-0.014 (3)	-0.010 (3)
O7	0.032 (4)	0.044 (4)	0.045 (4)	-0.001 (3)	0.002 (3)	0.003 (3)
O8	0.048 (4)	0.020 (4)	0.028 (4)	0.001 (3)	-0.006 (3)	0.003 (3)
O9	0.063 (5)	0.017 (4)	0.047 (4)	0.015 (3)	-0.025 (4)	-0.012 (3)
O10	0.022 (4)	0.061 (5)	0.066 (5)	-0.005 (3)	0.003 (3)	-0.038 (4)
O11	0.031 (3)	0.027 (4)	0.025 (3)	-0.001 (3)	-0.004 (3)	-0.006 (3)
O12	0.031 (4)	0.079 (6)	0.020 (3)	-0.006 (3)	-0.004 (3)	-0.018 (3)
O13	0.028 (3)	0.033 (4)	0.032 (4)	-0.006 (3)	0.015 (3)	-0.011 (3)
O14	0.034 (4)	0.088 (6)	0.069 (5)	0.000 (4)	-0.009 (4)	0.066 (5)
O15	0.038 (4)	0.046 (4)	0.030 (4)	-0.009 (3)	0.001 (3)	0.017 (3)
O16	0.024 (3)	0.030 (4)	0.034 (4)	0.003 (3)	-0.006 (3)	0.010 (3)
O17	0.024 (3)	0.048 (4)	0.015 (3)	0.007 (3)	0.010 (2)	0.000 (3)
O18	0.021 (3)	0.073 (5)	0.019 (3)	0.016 (3)	-0.012 (3)	-0.012 (3)
O19	0.029 (4)	0.040 (4)	0.029 (4)	0.004 (3)	-0.007 (3)	0.006 (3)
O20	0.033 (4)	0.062 (5)	0.020 (3)	-0.005 (3)	0.004 (3)	0.005 (3)
O21	0.026 (3)	0.023 (3)	0.016 (3)	0.004 (2)	0.001 (2)	0.002 (2)
O22	0.027 (3)	0.013 (3)	0.019 (3)	0.005 (2)	-0.001 (2)	-0.007 (2)
O23	0.023 (3)	0.040 (4)	0.033 (4)	0.005 (3)	0.007 (3)	-0.001 (3)
O24	0.136 (8)	0.038 (5)	0.079 (6)	-0.036 (5)	0.025 (6)	-0.010 (5)
C1	0.030 (5)	0.019 (5)	0.022 (5)	0.009 (4)	0.006 (4)	-0.002 (4)
C2	0.019 (4)	0.017 (4)	0.026 (5)	0.003 (3)	0.004 (3)	0.005 (4)
C3	0.030 (5)	0.033 (6)	0.026 (5)	-0.008 (4)	0.000 (4)	0.004 (4)
C4	0.035 (5)	0.049 (7)	0.026 (5)	-0.009 (5)	-0.001 (4)	-0.001 (5)
C5	0.041 (6)	0.061 (7)	0.020 (5)	-0.024 (5)	0.001 (4)	0.000 (5)
C6	0.027 (5)	0.047 (6)	0.026 (5)	-0.008 (4)	-0.006 (4)	0.003 (4)
C7	0.026 (5)	0.019 (5)	0.017 (4)	0.001 (4)	0.002 (3)	-0.003 (3)
C8	0.033 (5)	0.026 (5)	0.012 (4)	-0.002 (4)	0.000 (3)	0.000 (4)
C9	0.034 (5)	0.048 (6)	0.014 (4)	-0.011 (4)	0.000 (4)	0.001 (4)
C10	0.048 (6)	0.056 (7)	0.027 (5)	-0.020 (5)	-0.007 (5)	0.005 (5)
C11	0.063 (7)	0.038 (6)	0.024 (5)	-0.008 (5)	0.010 (5)	0.005 (5)
C12	0.039 (5)	0.031 (5)	0.018 (5)	-0.011 (4)	0.003 (4)	0.009 (4)
C13	0.022 (4)	0.018 (5)	0.021 (4)	-0.005 (3)	0.005 (4)	-0.002 (3)
C14	0.021 (4)	0.016 (4)	0.025 (5)	-0.006 (3)	0.006 (3)	0.001 (4)
C15	0.036 (5)	0.033 (6)	0.026 (5)	0.008 (4)	-0.003 (4)	-0.005 (4)
C16	0.051 (6)	0.043 (7)	0.025 (5)	0.003 (5)	-0.001 (5)	-0.001 (5)

supplementary materials

C17	0.030 (5)	0.030 (6)	0.059 (7)	0.004 (4)	0.020 (5)	0.003 (5)
C18	0.023 (5)	0.029 (5)	0.037 (6)	0.004 (4)	0.005 (4)	0.004 (4)
C19	0.024 (4)	0.021 (5)	0.021 (4)	0.003 (4)	0.001 (3)	-0.004 (4)
C20	0.028 (5)	0.011 (4)	0.026 (5)	0.001 (3)	-0.003 (4)	-0.004 (4)
C21	0.024 (5)	0.020 (5)	0.037 (5)	0.002 (4)	0.003 (4)	0.000 (4)
C22	0.046 (6)	0.022 (5)	0.027 (5)	0.004 (4)	0.009 (4)	0.004 (4)
C23	0.050 (6)	0.028 (5)	0.022 (5)	-0.001 (5)	-0.005 (4)	-0.006 (4)
C24	0.035 (5)	0.031 (6)	0.031 (5)	0.004 (4)	-0.014 (4)	-0.002 (4)
C25	0.017 (4)	0.024 (5)	0.022 (4)	-0.005 (3)	0.000 (3)	-0.003 (4)
C26	0.016 (4)	0.023 (5)	0.035 (5)	0.005 (3)	0.004 (4)	0.005 (4)
C27	0.025 (5)	0.021 (5)	0.014 (4)	0.002 (3)	0.001 (3)	-0.006 (3)
C28	0.025 (4)	0.025 (5)	0.010 (4)	0.002 (4)	0.000 (3)	-0.002 (3)
C29	0.024 (5)	0.044 (6)	0.027 (5)	0.003 (4)	0.000 (4)	-0.001 (4)
C30	0.035 (5)	0.047 (6)	0.017 (5)	0.007 (5)	0.012 (4)	-0.008 (4)
C31	0.050 (7)	0.069 (8)	0.013 (5)	0.004 (6)	-0.018 (4)	-0.010 (5)
C32	0.031 (5)	0.062 (7)	0.021 (5)	0.002 (5)	0.002 (4)	-0.007 (5)
C33	0.030 (5)	0.038 (6)	0.024 (5)	0.011 (4)	-0.002 (4)	0.014 (4)
C34	0.015 (4)	0.027 (5)	0.031 (5)	0.002 (3)	0.003 (4)	0.009 (4)
C35	0.031 (5)	0.033 (6)	0.031 (5)	-0.004 (4)	0.005 (4)	0.004 (4)
C36	0.046 (6)	0.037 (6)	0.029 (5)	-0.015 (5)	-0.007 (4)	0.000 (5)
C37	0.043 (6)	0.025 (5)	0.031 (5)	-0.011 (4)	-0.010 (4)	0.005 (4)
C38	0.031 (5)	0.022 (5)	0.041 (6)	-0.004 (4)	-0.002 (4)	0.006 (4)
C39	0.021 (4)	0.025 (5)	0.017 (4)	-0.005 (4)	0.005 (3)	0.000 (4)
C40	0.015 (4)	0.024 (5)	0.020 (4)	0.000 (3)	-0.003 (3)	0.001 (4)
C41	0.030 (5)	0.028 (5)	0.022 (5)	0.000 (4)	0.002 (4)	0.001 (4)
C42	0.041 (6)	0.041 (6)	0.024 (5)	-0.003 (5)	0.007 (4)	-0.006 (4)
C43	0.035 (5)	0.031 (6)	0.034 (6)	0.004 (4)	-0.008 (4)	-0.001 (4)
C44	0.019 (4)	0.041 (6)	0.029 (5)	-0.006 (4)	-0.003 (4)	-0.005 (4)
C45	0.019 (4)	0.018 (4)	0.023 (5)	-0.007 (3)	-0.002 (4)	0.010 (4)
C46	0.025 (4)	0.018 (4)	0.022 (5)	0.000 (3)	0.000 (3)	0.007 (4)
C47	0.026 (5)	0.034 (5)	0.022 (5)	0.006 (4)	-0.002 (4)	-0.002 (4)
C48	0.031 (5)	0.045 (6)	0.023 (5)	0.013 (4)	-0.007 (4)	0.005 (4)
C49	0.041 (6)	0.036 (6)	0.020 (5)	0.011 (4)	0.010 (4)	0.003 (4)
C50	0.019 (4)	0.031 (5)	0.031 (5)	0.008 (4)	0.001 (4)	0.000 (4)
C51	0.013 (4)	0.026 (5)	0.015 (4)	-0.009 (3)	0.001 (3)	0.008 (4)
C52	0.020 (4)	0.016 (4)	0.018 (4)	-0.002 (3)	-0.001 (3)	-0.003 (3)
C53	0.024 (5)	0.045 (6)	0.029 (5)	0.009 (4)	-0.004 (4)	-0.003 (4)
C54	0.045 (6)	0.051 (7)	0.025 (5)	0.006 (5)	0.006 (5)	-0.017 (5)
C55	0.043 (6)	0.034 (6)	0.026 (5)	-0.007 (5)	-0.007 (4)	-0.007 (4)
C56	0.030 (5)	0.038 (6)	0.020 (5)	0.005 (4)	-0.003 (4)	-0.004 (4)
N1	0.035 (4)	0.043 (5)	0.022 (4)	-0.013 (4)	-0.001 (3)	0.000 (4)
N2	0.051 (5)	0.049 (5)	0.018 (4)	-0.016 (4)	-0.001 (4)	-0.005 (4)
N3	0.055 (6)	0.044 (6)	0.033 (5)	-0.012 (4)	0.020 (4)	-0.006 (4)
N4	0.057 (5)	0.038 (5)	0.012 (4)	0.007 (4)	0.001 (4)	-0.007 (3)
N5	0.028 (4)	0.036 (5)	0.030 (4)	-0.014 (3)	-0.002 (3)	0.006 (4)
N6	0.048 (5)	0.036 (5)	0.019 (4)	-0.005 (4)	-0.004 (4)	0.001 (3)
N7	0.042 (5)	0.045 (5)	0.017 (4)	0.011 (4)	-0.004 (3)	-0.002 (4)
N8	0.048 (5)	0.035 (5)	0.018 (4)	-0.007 (4)	0.002 (4)	-0.008 (4)

supplementary materials

Geometric parameters (\AA , $^\circ$)

La1—O6	2.420 (6)	C4—N1	1.342 (11)
La1—O4 ⁱ	2.434 (5)	C4—H4A	0.9300
La1—O3	2.445 (5)	C5—N1	1.334 (11)
La1—O8	2.451 (6)	C5—C6	1.370 (12)
La1—O5 ⁱⁱ	2.486 (6)	C5—H5A	0.9300
La1—O1	2.494 (5)	C6—H6A	0.9300
La1—O2 ⁱⁱ	2.545 (5)	C7—C8	1.480 (11)
La1—O7	2.703 (6)	C8—C12	1.374 (11)
La2—O9	2.399 (6)	C8—C9	1.383 (11)
La2—O17	2.404 (5)	C9—C10	1.373 (12)
La2—O13	2.443 (6)	C9—H9A	0.9300
La2—O19	2.444 (6)	C10—N2	1.336 (12)
La2—O11	2.462 (5)	C10—H10A	0.9300
La2—O15	2.515 (6)	C11—N2	1.326 (12)
La2—O16	2.606 (5)	C11—C12	1.369 (12)
La2—O22 ⁱⁱⁱ	2.607 (5)	C11—H11A	0.9300
La3—O18	2.387 (5)	C12—H12A	0.9300
La3—O14 ^{iv}	2.436 (6)	C13—C14	1.505 (11)
La3—O20	2.436 (6)	C14—C15	1.369 (11)
La3—O12 ^{iv}	2.439 (6)	C14—C18	1.378 (11)
La3—O10 ^{iv}	2.515 (6)	C15—C16	1.371 (12)
La3—O21	2.574 (5)	C15—H15A	0.9300
La3—O24	2.610 (7)	C16—N3	1.319 (12)
La3—O23	2.657 (6)	C16—H16A	0.9300
La3—O11 ^{iv}	2.797 (6)	C17—N3	1.338 (13)
La3—C26 ^{iv}	3.043 (8)	C17—C18	1.377 (13)
Cu1—N4 ^v	1.920 (7)	C17—H17A	0.9300
Cu1—N2	1.931 (7)	C18—H18A	0.9300
Cu2—Cu2'	0.666 (13)	C19—C20	1.516 (11)
Cu2—N3	1.917 (9)	C20—C25	1.390 (11)
Cu2—Br1	2.212 (4)	C20—C21	1.405 (11)
Cu2—Cu7 ^{iv}	2.770 (4)	C21—C22	1.384 (12)
Cu2'—N3	2.064 (18)	C21—H21A	0.9300
Cu2'—Br1	2.415 (16)	C22—C23	1.357 (12)
Cu2'—Cu7 ^{iv}	2.569 (17)	C22—H22A	0.9300
Cu2'—Br3 ^{iv}	2.572 (16)	C23—C24	1.393 (12)
Cu3—Cu3'	0.629 (14)	C23—H23A	0.9300
Cu3—N1 ^{vi}	1.959 (8)	C24—C25	1.400 (11)
Cu3—N7 ^{vii}	1.980 (8)	C24—H24A	0.9300
Cu3—Br2	2.609 (4)	C25—C26	1.476 (11)
Cu3'—N7 ^{vii}	1.927 (19)	C26—La3 ^{vii}	3.043 (8)
Cu3'—N1 ^{vi}	1.976 (19)	C27—C28	1.493 (10)
Cu3'—Br3	2.72 (2)	C28—C29	1.380 (11)

Cu4—Br3	2.3071 (19)	C28—C32	1.399 (11)
Cu4—Br2	2.328 (2)	C29—C30	1.376 (11)
Cu4—Br4	2.3533 (19)	C29—H29A	0.9300
Cu4—Cu5	2.761 (2)	C30—N4	1.311 (11)
Cu5—N5	2.005 (7)	C30—H30A	0.9300
Cu5—Br5	2.3441 (16)	C31—N4	1.347 (12)
Cu5—Br4	2.3827 (16)	C31—C32	1.369 (12)
Cu6—N8 ^{viii}	2.004 (7)	C31—H31A	0.9300
Cu6—Br5	2.4397 (17)	C32—H32A	0.9300
Cu6—Br3	2.4946 (18)	C33—C34	1.490 (11)
Cu6—Br6	2.5481 (17)	C34—C38	1.364 (11)
Cu7—N6 ^{viii}	2.037 (7)	C34—C35	1.381 (12)
Cu7—Br6	2.4415 (18)	C35—C36	1.389 (12)
Cu7—Br1 ^{vii}	2.4447 (18)	C35—H35A	0.9300
Cu7—Cu2 ^{vii}	2.569 (17)	C36—N5	1.335 (11)
Cu7—Br4 ^{vii}	2.6722 (19)	C36—H36A	0.9300
Cu7—Cu2 ^{vii}	2.770 (4)	C37—N5	1.326 (11)
Br1—Cu7 ^{iv}	2.4447 (18)	C37—C38	1.386 (12)
Br3—Cu2 ^{vii}	2.572 (17)	C37—H37A	0.9300
Br4—Cu7 ^{iv}	2.6722 (19)	C38—H38A	0.9300
O1—C1	1.253 (9)	C39—C40	1.491 (11)
O2—C1	1.240 (9)	C40—C44	1.380 (11)
O2—La1 ⁱⁱ	2.545 (5)	C40—C41	1.382 (11)
O3—C7	1.268 (9)	C41—C42	1.389 (12)
O4—C7	1.247 (9)	C41—H41A	0.9300
O4—La1 ⁱ	2.434 (5)	C42—N6	1.342 (12)
O5—C13	1.247 (9)	C42—H42A	0.9300
O5—La1 ⁱⁱ	2.486 (6)	C43—N6	1.332 (12)
O6—C13	1.236 (9)	C43—C44	1.354 (12)
O7—H7A	0.8505	C43—H43A	0.9300
O7—H7B	0.8512	C44—H44A	0.9300
O8—C19	1.232 (10)	C45—C46	1.507 (11)
O9—C19	1.244 (10)	C46—C47	1.383 (11)
O10—C26	1.247 (9)	C46—C50	1.388 (11)
O10—La3 ^{vii}	2.515 (6)	C47—C48	1.374 (11)
O11—C26	1.279 (10)	C47—H47A	0.9300
O11—La3 ^{vii}	2.797 (6)	C48—N7	1.324 (11)
O12—C27	1.243 (9)	C48—H48A	0.9300
O12—La3 ^{vii}	2.439 (6)	C49—N7	1.317 (11)
O13—C27	1.256 (9)	C49—C50	1.377 (12)
O14—C33	1.250 (10)	C49—H49A	0.9300
O14—La3 ^{vii}	2.436 (6)	C50—H50A	0.9300
O15—C33	1.237 (10)	C51—C52	1.493 (10)
O16—H16C	0.8495	C52—C53	1.384 (11)
O16—H16D	0.8589	C52—C56	1.393 (11)
O17—C39	1.243 (9)	C53—C54	1.371 (12)

supplementary materials

O18—C39	1.250 (9)	C53—H53A	0.9300
O19—C45	1.235 (9)	C54—N8	1.343 (12)
O20—C45	1.241 (10)	C54—H54A	0.9300
O21—C51	1.258 (9)	C55—N8	1.328 (11)
O22—C51	1.252 (9)	C55—C56	1.379 (12)
O22—La2 ⁱⁱⁱ	2.607 (5)	C55—H55A	0.9300
O23—H23C	0.8013	C56—H56A	0.9300
O23—H23D	0.9328	N1—Cu3 ^v	1.959 (8)
O24—H24C	0.8501	N1—Cu3 ^{iv}	1.976 (19)
O24—H24D	0.8498	N4—Cu1 ^{vi}	1.920 (7)
C1—C2	1.521 (11)	N6—Cu7 ^{ix}	2.037 (7)
C2—C6	1.375 (11)	N7—Cu3 ^{iv}	1.927 (19)
C2—C3	1.380 (11)	N7—Cu3 ^{iv}	1.980 (8)
C3—C4	1.390 (12)	N8—Cu6 ^{ix}	2.004 (7)
C3—H3A	0.9300		
O6—La1—O4 ⁱ	150.8 (2)	H23C—O23—H23D	103.4
O6—La1—O3	92.4 (2)	La3—O24—H24C	162.5
O4 ⁱ —La1—O3	85.15 (19)	La3—O24—H24D	89.5
O6—La1—O8	75.4 (2)	H24C—O24—H24D	107.7
O4 ⁱ —La1—O8	75.8 (2)	O2—C1—O1	127.2 (8)
O3—La1—O8	74.6 (2)	O2—C1—C2	117.0 (7)
O6—La1—O5 ⁱⁱ	115.8 (2)	O1—C1—C2	115.7 (7)
O4 ⁱ —La1—O5 ⁱⁱ	82.4 (2)	C6—C2—C3	115.9 (8)
O3—La1—O5 ⁱⁱ	140.6 (2)	C6—C2—C1	123.3 (8)
O8—La1—O5 ⁱⁱ	136.6 (2)	C3—C2—C1	120.8 (7)
O6—La1—O1	72.8 (2)	C2—C3—C4	120.1 (8)
O4 ⁱ —La1—O1	134.92 (19)	C2—C3—H3A	119.9
O3—La1—O1	80.7 (2)	C4—C3—H3A	119.9
O8—La1—O1	138.4 (2)	N1—C4—C3	122.8 (8)
O5 ⁱⁱ —La1—O1	82.1 (2)	N1—C4—H4A	118.6
O6—La1—O2 ⁱⁱ	75.5 (2)	C3—C4—H4A	118.6
O4 ⁱ —La1—O2 ⁱⁱ	91.89 (18)	N1—C5—C6	122.6 (9)
O3—La1—O2 ⁱⁱ	149.2 (2)	N1—C5—H5A	118.7
O8—La1—O2 ⁱⁱ	74.9 (2)	C6—C5—H5A	118.7
O5 ⁱⁱ —La1—O2 ⁱⁱ	68.7 (2)	C5—C6—C2	121.7 (8)
O1—La1—O2 ⁱⁱ	120.78 (18)	C5—C6—H6A	119.1
O6—La1—O7	138.0 (2)	C2—C6—H6A	119.1
O4 ⁱ —La1—O7	68.51 (19)	O4—C7—O3	125.2 (7)
O3—La1—O7	71.97 (19)	O4—C7—C8	117.0 (7)
O8—La1—O7	132.38 (19)	O3—C7—C8	117.7 (7)
O5 ⁱⁱ —La1—O7	68.7 (2)	C12—C8—C9	117.5 (8)
O1—La1—O7	66.41 (19)	C12—C8—C7	120.8 (8)
O2 ⁱⁱ —La1—O7	134.9 (2)	C9—C8—C7	121.7 (8)

O9—La2—O17	76.8 (2)	C10—C9—C8	119.3 (8)
O9—La2—O13	147.7 (2)	C10—C9—H9A	120.3
O17—La2—O13	135.3 (2)	C8—C9—H9A	120.3
O9—La2—O19	88.4 (2)	N2—C10—C9	123.3 (9)
O17—La2—O19	87.0 (2)	N2—C10—H10A	118.3
O13—La2—O19	89.7 (2)	C9—C10—H10A	118.3
O9—La2—O11	72.6 (2)	N2—C11—C12	124.1 (9)
O17—La2—O11	109.47 (19)	N2—C11—H11A	117.9
O13—La2—O11	94.55 (19)	C12—C11—H11A	117.9
O19—La2—O11	150.5 (2)	C11—C12—C8	119.2 (9)
O9—La2—O15	77.8 (2)	C11—C12—H12A	120.4
O17—La2—O15	150.7 (2)	C8—C12—H12A	120.4
O13—La2—O15	70.3 (2)	O6—C13—O5	124.6 (8)
O19—La2—O15	77.8 (2)	O6—C13—C14	118.9 (7)
O11—La2—O15	76.2 (2)	O5—C13—C14	116.6 (7)
O9—La2—O16	118.8 (2)	C15—C14—C18	118.7 (8)
O17—La2—O16	77.99 (19)	C15—C14—C13	119.3 (8)
O13—La2—O16	79.00 (19)	C18—C14—C13	122.0 (8)
O19—La2—O16	144.06 (19)	C14—C15—C16	119.3 (9)
O11—La2—O16	65.12 (18)	C14—C15—H15A	120.3
O15—La2—O16	127.83 (19)	C16—C15—H15A	120.3
O9—La2—O22 ⁱⁱⁱ	140.9 (2)	N3—C16—C15	122.9 (9)
O17—La2—O22 ⁱⁱⁱ	66.62 (19)	N3—C16—H16A	118.5
O13—La2—O22 ⁱⁱⁱ	69.22 (17)	C15—C16—H16A	118.5
O19—La2—O22 ⁱⁱⁱ	76.68 (18)	N3—C17—C18	123.1 (9)
O11—La2—O22 ⁱⁱⁱ	131.93 (18)	N3—C17—H17A	118.5
O15—La2—O22 ⁱⁱⁱ	131.72 (19)	C18—C17—H17A	118.5
O16—La2—O22 ⁱⁱⁱ	67.40 (17)	C17—C18—C14	118.2 (9)
O18—La3—O14 ^{iv}	144.2 (3)	C17—C18—H18A	120.9
O18—La3—O20	89.3 (2)	C14—C18—H18A	120.9
O14 ^{iv} —La3—O20	83.5 (2)	O8—C19—O9	124.5 (8)
O18—La3—O12 ^{iv}	139.1 (2)	O8—C19—C20	117.4 (7)
O14 ^{iv} —La3—O12 ^{iv}	76.4 (3)	O9—C19—C20	118.1 (8)
O20—La3—O12 ^{iv}	91.9 (2)	C25—C20—C21	119.1 (8)
O18—La3—O10 ^{iv}	75.0 (2)	C25—C20—C19	124.9 (7)
O14 ^{iv} —La3—O10 ^{iv}	85.1 (3)	C21—C20—C19	115.8 (7)
O20—La3—O10 ^{iv}	134.2 (2)	C22—C21—C20	120.9 (8)
O12 ^{iv} —La3—O10 ^{iv}	127.9 (2)	C22—C21—H21A	119.5
O18—La3—O21	75.0 (2)	C20—C21—H21A	119.5
O14 ^{iv} —La3—O21	131.0 (2)	C23—C22—C21	120.1 (9)
O20—La3—O21	137.6 (2)	C23—C22—H22A	120.0
O12 ^{iv} —La3—O21	76.8 (2)	C21—C22—H22A	120.0
O10 ^{iv} —La3—O21	80.0 (2)	C22—C23—C24	120.1 (9)
O18—La3—O24	70.9 (3)	C22—C23—H23A	120.0
O14 ^{iv} —La3—O24	73.9 (3)	C24—C23—H23A	120.0

supplementary materials

O20—La3—O24	68.5 (3)	C23—C24—C25	120.9 (8)
O12 ^{iv} —La3—O24	145.8 (3)	C23—C24—H24A	119.5
O10 ^{iv} —La3—O24	65.7 (3)	C25—C24—H24A	119.5
O21—La3—O24	136.4 (2)	C20—C25—C24	118.8 (8)
O18—La3—O23	74.5 (2)	C20—C25—C26	124.2 (7)
O14 ^{iv} —La3—O23	132.5 (3)	C24—C25—C26	117.0 (7)
O20—La3—O23	67.4 (2)	O10—C26—O11	120.5 (8)
O12 ^{iv} —La3—O23	68.4 (2)	O10—C26—C25	118.7 (8)
O10 ^{iv} —La3—O23	141.9 (2)	O11—C26—C25	120.6 (7)
O21—La3—O23	70.47 (17)	O10—C26—La3 ^{vii}	53.7 (4)
O24—La3—O23	123.1 (3)	O11—C26—La3 ^{vii}	66.8 (4)
O18—La3—O11 ^{iv}	115.82 (19)	C25—C26—La3 ^{vii}	171.5 (6)
O14 ^{iv} —La3—O11 ^{iv}	66.7 (2)	O12—C27—O13	124.9 (7)
O20—La3—O11 ^{iv}	150.15 (19)	O12—C27—C28	117.7 (7)
O12 ^{iv} —La3—O11 ^{iv}	79.61 (18)	O13—C27—C28	117.4 (7)
O10 ^{iv} —La3—O11 ^{iv}	48.42 (18)	C29—C28—C32	116.4 (8)
O21—La3—O11 ^{iv}	68.52 (17)	C29—C28—C27	122.2 (7)
O24—La3—O11 ^{iv}	103.3 (2)	C32—C28—C27	121.4 (7)
O23—La3—O11 ^{iv}	132.43 (17)	C30—C29—C28	119.9 (8)
O18—La3—C26 ^{iv}	95.1 (2)	C30—C29—H29A	120.1
O14 ^{iv} —La3—C26 ^{iv}	74.9 (3)	C28—C29—H29A	120.1
O20—La3—C26 ^{iv}	148.7 (2)	N4—C30—C29	124.0 (8)
O12 ^{iv} —La3—C26 ^{iv}	104.4 (2)	N4—C30—H30A	118.0
O10 ^{iv} —La3—C26 ^{iv}	23.6 (2)	C29—C30—H30A	118.0
O21—La3—C26 ^{iv}	73.01 (19)	N4—C31—C32	123.2 (8)
O24—La3—C26 ^{iv}	83.7 (3)	N4—C31—H31A	118.4
O23—La3—C26 ^{iv}	143.5 (2)	C32—C31—H31A	118.4
O11 ^{iv} —La3—C26 ^{iv}	24.84 (19)	C31—C32—C28	119.6 (9)
N4 ^v —Cu1—N2	166.2 (3)	C31—C32—H32A	120.2
Cu2'—Cu2—N3	93.2 (18)	C28—C32—H32A	120.2
Cu2'—Cu2—Br1	99.8 (18)	O15—C33—O14	126.0 (8)
N3—Cu2—Br1	166.5 (3)	O15—C33—C34	119.1 (7)
Cu2'—Cu2—Cu7 ^{iv}	65.7 (18)	O14—C33—C34	114.9 (8)
N3—Cu2—Cu7 ^{iv}	133.0 (3)	C38—C34—C35	118.3 (8)
Br1—Cu2—Cu7 ^{iv}	57.48 (10)	C38—C34—C33	121.0 (8)
Cu2—Cu2'—N3	68.0 (16)	C35—C34—C33	120.6 (8)
Cu2—Cu2'—Br1	64.5 (16)	C34—C35—C36	119.0 (8)
N3—Cu2'—Br1	132.4 (7)	C34—C35—H35A	120.5
Cu2—Cu2'—Cu7 ^{iv}	101 (2)	C36—C35—H35A	120.5
N3—Cu2'—Cu7 ^{iv}	136.8 (10)	N5—C36—C35	123.1 (9)
Br1—Cu2'—Cu7 ^{iv}	58.6 (4)	N5—C36—H36A	118.5
Cu2—Cu2'—Br3 ^{iv}	152 (3)	C35—C36—H36A	118.5

N3—Cu2'—Br3 ^{iv}	113.9 (7)	N5—C37—C38	124.2 (8)
Br1—Cu2'—Br3 ^{iv}	106.3 (7)	N5—C37—H37A	117.9
Cu7 ^{iv} —Cu2'—Br3 ^{iv}	95.3 (5)	C38—C37—H37A	117.9
Cu3'—Cu3—N1 ^{vi}	82.3 (19)	C34—C38—C37	118.7 (9)
Cu3'—Cu3—N7 ^{vii}	76.0 (19)	C34—C38—H38A	120.6
N1 ^{vi} —Cu3—N7 ^{vii}	148.7 (4)	C37—C38—H38A	120.6
Cu3'—Cu3—Br2	147 (2)	O17—C39—O18	123.8 (7)
N1 ^{vi} —Cu3—Br2	106.5 (3)	O17—C39—C40	118.7 (7)
N7 ^{vii} —Cu3—Br2	104.0 (3)	O18—C39—C40	117.5 (7)
Cu3—Cu3'—N7 ^{vii}	86 (2)	C44—C40—C41	119.4 (8)
Cu3—Cu3'—N1 ^{vi}	79.3 (19)	C44—C40—C39	120.2 (7)
N7 ^{vii} —Cu3'—N1 ^{vi}	152.7 (12)	C41—C40—C39	120.4 (7)
Cu3—Cu3'—Br3	109 (2)	C40—C41—C42	118.7 (8)
N7 ^{vii} —Cu3'—Br3	92.9 (7)	C40—C41—H41A	120.7
N1 ^{vi} —Cu3'—Br3	113.5 (9)	C42—C41—H41A	120.7
Br3—Cu4—Br2	114.05 (8)	N6—C42—C41	121.9 (9)
Br3—Cu4—Br4	125.56 (8)	N6—C42—H42A	119.0
Br2—Cu4—Br4	120.39 (8)	C41—C42—H42A	119.0
Br3—Cu4—Cu5	82.39 (6)	N6—C43—C44	125.1 (9)
Br2—Cu4—Cu5	142.32 (8)	N6—C43—H43A	117.5
Br4—Cu4—Cu5	54.84 (5)	C44—C43—H43A	117.5
N5—Cu5—Br5	120.8 (2)	C43—C44—C40	117.7 (9)
N5—Cu5—Br4	108.5 (2)	C43—C44—H44A	121.2
Br5—Cu5—Br4	130.48 (6)	C40—C44—H44A	121.2
N5—Cu5—Cu4	88.5 (2)	O19—C45—O20	125.8 (8)
Br5—Cu5—Cu4	127.83 (7)	O19—C45—C46	117.5 (8)
Br4—Cu5—Cu4	53.85 (5)	O20—C45—C46	116.7 (7)
N8 ^{viii} —Cu6—Br5	119.7 (2)	C47—C46—C50	117.3 (8)
N8 ^{viii} —Cu6—Br3	106.4 (2)	C47—C46—C45	121.1 (8)
Br5—Cu6—Br3	109.80 (6)	C50—C46—C45	121.5 (7)
N8 ^{viii} —Cu6—Br6	113.8 (2)	C48—C47—C46	118.9 (8)
Br5—Cu6—Br6	105.98 (6)	C48—C47—H47A	120.6
Br3—Cu6—Br6	99.15 (6)	C46—C47—H47A	120.6
N6 ^{viii} —Cu7—Br6	109.2 (2)	N7—C48—C47	124.0 (8)
N6 ^{viii} —Cu7—Br1 ^{vii}	108.6 (2)	N7—C48—H48A	118.0
Br6—Cu7—Br1 ^{vii}	129.77 (7)	C47—C48—H48A	118.0
N6 ^{viii} —Cu7—Cu2 ^{vii}	130.7 (5)	N7—C49—C50	124.0 (8)
Br6—Cu7—Cu2 ^{vii}	72.8 (4)	N7—C49—H49A	118.0
Br1 ^{vii} —Cu7—Cu2 ^{vii}	57.5 (4)	C50—C49—H49A	118.0
N6 ^{viii} —Cu7—Br4 ^{vii}	98.8 (2)	C49—C50—C46	118.9 (8)
Br6—Cu7—Br4 ^{vii}	102.53 (6)	C49—C50—H50A	120.5
Br1 ^{vii} —Cu7—Br4 ^{vii}	103.04 (6)	C46—C50—H50A	120.5
Cu2 ^{vii} —Cu7—Br4 ^{vii}	129.6 (4)	O22—C51—O21	125.4 (7)

supplementary materials

N6 ^{viii} —Cu7—Cu2 ^{vii}	140.1 (3)	O22—C51—C52	118.1 (7)
Br6—Cu7—Cu2 ^{vii}	80.14 (9)	O21—C51—C52	116.6 (7)
Br1 ^{vii} —Cu7—Cu2 ^{vii}	49.72 (9)	C53—C52—C56	116.2 (7)
Cu2 ^{vii} —Cu7—Cu2 ^{vii}	13.7 (3)	C53—C52—C51	123.1 (7)
Br4 ^{vii} —Cu7—Cu2 ^{vii}	117.39 (9)	C56—C52—C51	120.6 (7)
Cu2—Br1—Cu2'	15.8 (3)	C54—C53—C52	121.2 (8)
Cu2—Br1—Cu7 ^{iv}	72.81 (11)	C54—C53—H53A	119.4
Cu2'—Br1—Cu7 ^{iv}	63.8 (4)	C52—C53—H53A	119.4
Cu4—Br2—Cu3	82.10 (12)	N8—C54—C53	121.9 (9)
Cu4—Br3—Cu6	104.81 (7)	N8—C54—H54A	119.0
Cu4—Br3—Cu2 ^{vii}	158.6 (4)	C53—C54—H54A	119.0
Cu6—Br3—Cu2 ^{vii}	91.1 (4)	N8—C55—C56	123.3 (9)
Cu4—Br3—Cu3'	85.6 (3)	N8—C55—H55A	118.3
Cu6—Br3—Cu3'	156.9 (4)	C56—C55—H55A	118.3
Cu2 ^{vii} —Br3—Cu3'	85.1 (5)	C55—C56—C52	119.6 (8)
Cu4—Br4—Cu5	71.31 (6)	C55—C56—H56A	120.2
Cu4—Br4—Cu7 ^{iv}	120.66 (6)	C52—C56—H56A	120.2
Cu5—Br4—Cu7 ^{iv}	123.90 (6)	C5—N1—C4	116.9 (8)
Cu5—Br5—Cu6	84.24 (6)	C5—N1—Cu3 ^v	121.7 (6)
Cu7—Br6—Cu6	98.64 (6)	C4—N1—Cu3 ^v	121.3 (6)
C1—O1—La1	136.1 (6)	C5—N1—Cu3 ^w	132.7 (8)
C1—O2—La1 ⁱⁱ	143.8 (5)	C4—N1—Cu3 ^w	109.2 (8)
C7—O3—La1	145.1 (5)	Cu3 ^v —N1—Cu3 ^w	18.4 (4)
C7—O4—La1 ⁱ	152.6 (6)	C11—N2—C10	116.4 (8)
C13—O5—La1 ⁱⁱ	116.1 (5)	C11—N2—Cu1	119.1 (6)
C13—O6—La1	165.8 (6)	C10—N2—Cu1	124.4 (6)
La1—O7—H7A	99.1	C16—N3—C17	117.6 (8)
La1—O7—H7B	102.7	C16—N3—Cu2	122.9 (7)
H7A—O7—H7B	107.5	C17—N3—Cu2	119.4 (7)
C19—O8—La1	163.1 (6)	C16—N3—Cu2'	104.2 (8)
C19—O9—La2	146.1 (6)	C17—N3—Cu2'	137.9 (8)
C26—O10—La3 ^{vii}	102.7 (5)	Cu2—N3—Cu2'	18.8 (4)
C26—O11—La2	146.8 (5)	C30—N4—C31	116.8 (7)
C26—O11—La3 ^{vii}	88.4 (5)	C30—N4—Cu1 ^{vi}	123.0 (6)
La2—O11—La3 ^{vii}	123.8 (2)	C31—N4—Cu1 ^{vi}	120.2 (6)
C27—O12—La3 ^{vii}	150.7 (6)	C37—N5—C36	116.5 (8)
C27—O13—La2	129.7 (5)	C37—N5—Cu5	119.6 (6)
C33—O14—La3 ^{vii}	143.8 (7)	C36—N5—Cu5	123.6 (6)
C33—O15—La2	141.0 (6)	C43—N6—C42	117.2 (8)
La2—O16—H16C	150.9	C43—N6—Cu7 ^{ix}	121.8 (6)
La2—O16—H16D	101.5	C42—N6—Cu7 ^{ix}	121.0 (6)
H16C—O16—H16D	107.0	C49—N7—C48	116.8 (7)
C39—O17—La2	151.1 (6)	C49—N7—Cu3 ^{iv}	111.3 (8)

supplementary materials

C39—O18—La3	147.0 (5)	C48—N7—Cu3 ^{iv}	127.6 (9)
C45—O19—La2	150.1 (6)	C49—N7—Cu3 ^{iv}	123.6 (6)
C45—O20—La3	154.4 (6)	C48—N7—Cu3 ^{iv}	119.4 (6)
C51—O21—La3	129.1 (5)	Cu3 ^{iv} —N7—Cu3 ^{iv}	18.5 (4)
C51—O22—La2 ⁱⁱⁱ	126.3 (5)	C55—N8—C54	117.7 (8)
La3—O23—H23C	114.8	C55—N8—Cu6 ^{ix}	121.1 (6)
La3—O23—H23D	115.0	C54—N8—Cu6 ^{ix}	121.0 (6)

Symmetry codes: (i) $-x+1, -y+1, -z$; (ii) $-x, -y+1, -z$; (iii) $-x, -y, -z$; (iv) $x-1, y, z$; (v) $-x+1, y+1/2, -z-1/2$; (vi) $-x+1, y-1/2, -z-1/2$; (vii) $x+1, y, z$; (viii) $x+1, -y+1/2, z-1/2$; (ix) $x-1, -y+1/2, z+1/2$.

Hydrogen-bond geometry (\AA , $^\circ$)

$D\text{—H}\cdots A$	$D\text{—H}$	$H\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
O16—H16D \cdots O21 ^{vii}	0.86	2.09	2.901 (7)	157
O23—H23D \cdots O22	0.93	2.00	2.861 (8)	153
O24—H24D \cdots O20	0.85	2.22	2.844 (11)	130

Symmetry codes: (vii) $x+1, y, z$.

supplementary materials

Fig. 1





Fig. 2



supplementary materials

Fig. 3





Fig. 4



supplementary materials

Fig. 5




Fig. 6

