

2-[*(E*)-2-(1*H*-Indol-3-yl)ethenyl]-1-methylpyridinium 4-bromobenzene-sulfonate¹

Suchada Chantrapromma,^{a*} Thawanrat Kobkeatthawin^a and Hoong-Kun Fun^b[§]

^aCrystal Materials Research Unit, Department of Chemistry, Faculty of Science, Prince of Songkla University, Hat-Yai, Songkhla 90112, Thailand, and ^bX-ray Crystallography Unit, School of Physics, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia

Correspondence e-mail: suchada.c@psu.ac.th

Received 16 March 2009; accepted 27 March 2009

Key indicators: single-crystal X-ray study; $T = 100\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$; R factor = 0.033; wR factor = 0.085; data-to-parameter ratio = 39.6.


In the title compound, $\text{C}_{16}\text{H}_{15}\text{N}_2^+\cdot\text{C}_6\text{H}_4\text{BrO}_3\text{S}^-$, the cation exists in the *E* configuration and is essentially planar with a dihedral angle of $3.10(5)^\circ$ between the pyridinium ring and the indole ring system. The π -conjugated planes of the cation and the anion are inclined to each other at a dihedral angle of $64.32(4)^\circ$. In the crystal structure, the cations are stacked in an antiparallel manner along the a axis. The anions are linked into a chain along the a axis. The cations and the anions are linked into a three-dimensional network by $\text{N}-\text{H}\cdots\text{O}$ and weak $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds. The crystal structure is further stabilized by $\text{C}-\text{H}\cdots\pi$ interactions. A $\pi-\pi$ interaction between the five-membered heterocyclic ring of the indole system and the pyridinium ring is also observed with a centroid–centroid distance of $3.5855(7)\text{ \AA}$.

Related literature

For bond-length data, see: Allen *et al.* (1987). For background to non-linear optical materials research, see: Coe *et al.* (2003); Dittrich *et al.* (2003); Ogawa *et al.* (2008); Otero *et al.* (2002); Weir *et al.* (2003); Yang *et al.* (2007). For related structures, see, for example: Chanawanno *et al.* (2008); Chantrapromma *et al.* (2006, 2007, 2008, 2009); Jindawong *et al.* (2005). For the stability of the temperature controller used in the data collection, see: Cosier & Glazer (1986).

¹This paper is dedicated to the late Her Royal Highness Princess Galyani Vadhana Krom Luang Naradhiwas Rajanagarindra for her patronage of Science in Thailand.

[§]Additional correspondence author, e-mail: hkfun@usm.my.



Experimental

Crystal data

$\text{C}_{16}\text{H}_{15}\text{N}_2^+\cdot\text{C}_6\text{H}_4\text{BrO}_3\text{S}^-$	$V = 2012.49(5)\text{ \AA}^3$
$M_r = 471.36$	$Z = 4$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
$a = 7.5188(1)\text{ \AA}$	$\mu = 2.17\text{ mm}^{-1}$
$b = 13.3659(2)\text{ \AA}$	$T = 100\text{ K}$
$c = 20.2670(3)\text{ \AA}$	$0.31 \times 0.27 \times 0.16\text{ mm}$
$\beta = 98.850(1)^\circ$	

Data collection

Bruker APEXII CCD area-detector diffractometer	67716 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 2005)	10581 independent reflections
$T_{\min} = 0.548$, $T_{\max} = 0.706$	8073 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.033$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.033$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.085$	$\Delta\rho_{\text{max}} = 1.02\text{ e \AA}^{-3}$
$S = 1.03$	$\Delta\rho_{\text{min}} = -0.66\text{ e \AA}^{-3}$
10581 reflections	
267 parameters	

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N2—H1N2 \cdots O2 ⁱ	0.85 (2)	1.91 (2)	2.7593 (14)	175.2 (17)
C1—H1A \cdots O3 ⁱⁱ	0.93	2.53	3.2067 (16)	130
C7—H7A \cdots O1	0.93	2.58	3.3095 (16)	136
C9—H9A \cdots O1	0.93	2.58	3.2426 (16)	128
C14—H14A \cdots O1 ⁱⁱⁱ	0.93	2.56	3.2987 (16)	137
C16—H16C \cdots O1 ⁱⁱⁱ	0.96	2.36	3.2739 (17)	158
C19—H19A \cdots O3 ^{iv}	0.93	2.51	3.2052 (16)	131
C21—H21A \cdots O2 ^v	0.93	2.28	3.1310 (15)	152
C4—H4A \cdots Cg3	0.93	2.82	3.5579 (13)	137
C16—H16A \cdots Cg3 ^{vi}	0.96	2.69	3.5731 (13)	154
C16—H16B \cdots Cg1 ^{vii}	0.96	2.74	3.4836 (14)	135

Symmetry codes: (i) $-x + 2, y + \frac{1}{2}, -z + \frac{1}{2}$; (ii) $x, -y + \frac{3}{2}, z + \frac{1}{2}$; (iii) $-x + 2, -y + 2, -z + 1$; (iv) $-x + 2, y - \frac{1}{2}, -z + \frac{1}{2}$; (v) $x - 1, y, z$; (vi) $x, -y + \frac{1}{2}, z - \frac{1}{2}$; (vii) $-x + 1, -y + 2, -z + 1$. Cg1 and Cg3 are the centroids of the N2/C8-C10/Cl5 and C10-C15 rings, respectively.

Data collection: *APEX2* (Bruker, 2005); cell refinement: *SAINT* (Bruker, 2005); data reduction: *SAINT*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL* and *PLATON* (Spek, 2009).

The authors thank Prince of Songkla University for financial support through the Crystal Materials Research Unit. The authors also thank Universiti Sains Malaysia for the Research University Golden Goose grant No. 1001/PFIZIK/811012.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: IS2401).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Bruker (2005). *APEX2, SAINT and SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Chanawanno, K., Chantrapromma, S. & Fun, H.-K. (2008). *Acta Cryst. E* **64**, o1882–o1883.
- Chantrapromma, S., Chotika, L., Ruanwas, P. & Fun, H.-K. (2008). *Acta Cryst. E* **64**, o574–o575.
- Chantrapromma, S., Jansrisewangwong, P., Musor, R. & Fun, H.-K. (2009). *Acta Cryst. E* **65**, o217–o218.
- Chantrapromma, S., Ruanwas, P., Fun, H.-K. & Patil, P. S. (2006). *Acta Cryst. E* **62**, o5494–o5496.
- Chantrapromma, S., Suwanwong, T. & Fun, H.-K. (2007). *Acta Cryst. E* **63**, o821–o823.
- Coe, B. J., Harris, J. A., Asselberghs, I., Wostyn, K., Clays, K., Persoons, A., Brunschwig, B. S., Coles, S. J., Gelbrich, T., Light, M. E., Hursthous, M. B. & Nakatani, K. (2003). *Adv. Funct. Mater.* **13**, 347–357.
- Cosier, J. & Glazer, A. M. (1986). *J. Appl. Cryst.* **19**, 105–107.
- Dittrich, Ph., Bartlome, R., Montemezzani, G. & Günter, P. (2003). *Appl. Surf. Sci.* **220**, 88–95.
- Jindawong, B., Chantrapromma, S., Fun, H.-K., Yu, X.-L. & Karalai, C. (2005). *Acta Cryst. E* **61**, o1340–o1342.
- Ogawa, J., Okada, S., Glavcheva, Z. & Nakanishi, H. (2008). *J. Cryst. Growth*, **310**, 836–842.
- Otero, M., Herranz, M. A., Seoane, C., Martín, N., Garín, J., Orduna, J., Alcalá, R. & Villacampa, B. (2002). *Tetrahedron*, **58**, 7463–7475.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.
- Weir, C. A. M., Hadizad, T., Beaudin, A. M. R. & Wang, Z.-Y. (2003). *Tetrahedron Lett.* **44**, 4697–4700.
- Yang, Z., Wörle, M., Mutter, L., Jazbinsek, M. & Günter, P. (2007). *Cryst. Growth Des.* **7**, 83–86.

supplementary materials

Acta Cryst. (2009). E65, o950-o951 [doi:10.1107/S1600536809011386]

2-[(*E*)-2-(1*H*-Indol-3-yl)ethenyl]-1-methylpyridinium 4-bromobenzenesulfonate

S. Chantrapromma, T. Kobkeatthawin and H.-K. Fun

Comment

Organic crystals with extensive conjugated π systems are attractive candidates for nonlinear optic (NLO) studies because of their large hyperpolarizability (β) and ease of preparation (Coe *et al.*, 2003; Dittrich *et al.*, 2003; Ogawa *et al.*, 2008; Otero *et al.*, 2002; Weir *et al.*, 2003; Yang *et al.*, 2007). One strategy to enhance the hyperpolarizability of the cations is by elongation of its π -conjugation system. Based on these studies, we have previously synthesized and reported the crystal structure of the pyridinium salts (Chanawanno *et al.*, 2008; Chantrapromma *et al.*, 2006, 2007, 2008, 2009; Jindawong *et al.*, 2005) in order to study for their NLO properties. We herein report the crystal structure of the title compound, (I), which is another pyridinium salt.

Figure 1 shows the asymmetric unit of (I), which consists of a $C_{16}H_{15}N_2^+$ cation and a $C_6H_4BrO_3S^-$ anion. The cation exists in the *E* configuration with respect to the C6=C7 double bond [1.3568 (16) Å] and is essentially planar with a dihedral angle between the pyridinium and indole rings being 3.10 (5) $^\circ$, the torsion angles C4–C5–C6–C7 = -2.13 (19) $^\circ$ and C6–C7–C8–C15 = 3.9 (2) $^\circ$. The indole ring system is planar with the most deviation of -0.0137 (12) Å for atom C8. The π -conjugated planes of the cation and the anion are inclined to each other with the interplanar angle between them being 64.32 (4) $^\circ$. The methyl group is co-planar with the attached N1/C1–C5 ring. The bond lengths in (I) are in normal ranges (Allen *et al.*, 1987) and comparable with those in related structures (Chanawanno *et al.*, 2008; Chantrapromma *et al.*, 2006, 2007, 2008, 2009; Jindawong *et al.*, 2005).

In the crystal packing (Fig. 2), all O atoms of the sulfonate group are involved in weak C—H···O interactions (Table 1). The arrangement of the cations and anions is interesting (Fig. 2). The cations are stacked in an antiparallel manner along the *a* axis and the anions are linked together into chains along the same direction. The cations are linked to the anions into a three dimensional network by N—H···O hydrogen bonds and weak C—H···O interactions (Table 1). The crystal structure is further stabilized by C—H··· π interactions (Table 1). A π – π interaction with a distance $Cg1\cdots Cg2 = 3.5855$ (7) Å (symmetry code: 2-x, 2-y, 1-z) is observed; $Cg1$ and $Cg2$ are the centroids of the N2/C8–C10/C15 and N1/C1–C5 rings, respectively.

Experimental

A solution of indole-3-carboxaldehyde (2.47 g, 17.02 mmol) in methanol (25 ml) was added dropwise to a stirred solution of 1,2-dimethylpyridinium iodide (4.00 g, 17.02 mmol) in methanol (15 ml) in the presence of piperidine (1.68 ml, 17.02 mmol) over a period of 15 mins at room temperature. The mixture was then refluxed for 1 hr in the nitrogen atmosphere. The solid formed was filtered, washed with diethyl ether and recrystallized from methanol to give orange crystals of 2-[(*E*)-2-(1*H*-Indol-3-yl)ethenyl]-1-methylpyridinium iodide (compound A) (5.61 g, 91%; m.p. 537–539 K).

Silver(I) *p*-bromobenzenesulfonate (compound B) was synthesized according to our previously reported procedure (Chantrapromma *et al.*, 2006). The title compound was synthesized by dissolving compound B (0.20 g, 0.58 mmol) in 20 ml methanol which upon heating was added a solution of compound A (0.21 g, 0.58 mmol) in hot methanol (30 ml). The mixture turned yellow and cloudy immediately. After stirring for 0.5 hr, the precipitate of silver iodide was filtered and the

supplementary materials

filtrate was evaporated to give an orange gum. Orange block-shaped single crystals of the title compound suitable for *x*-ray structure determination were recrystallized from methanol by slow evaporation of the solvent at room temperature after a few weeks (m.p. 508–510 K).

Refinement

H atom attached to N was located from the difference map and refined isotropically. The remaining H atoms were placed in calculated positions with C—H = 0.93 Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ for aromatic and CH, and with C—H = 0.96 Å and $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$ for CH_3 atoms. A rotating group model was used for the methyl groups. The highest residual electron density peak is located at 0.59 Å from S1 and the deepest hole is located at 0.36 Å from Br1.

Figures




Fig. 1. The molecular structure of the title compound, with 50% probability displacement ellipsoids and the atom-numbering scheme.





Fig. 2. The crystal packing of the title compound, viewed down the *b* axis. Hydrogen bonds are shown as dashed lines.

2-[*(E*)-2-(1*H*-Indol-3-yl)ethenyl]-1-methylpyridinium 4-bromobenzenesulfonate

Crystal data



$$F_{000} = 960$$

$$M_r = 471.36$$

$$D_x = 1.556 \text{ Mg m}^{-3}$$

Monoclinic, $P2_1/c$

Melting point = 508–510 K

Hall symbol: -P 2ybc

Mo $K\alpha$ radiation

$$\lambda = 0.71073 \text{ \AA}$$

$$a = 7.5188 (1) \text{ \AA}$$

Cell parameters from 10581 reflections

$$b = 13.3659 (2) \text{ \AA}$$

$$\theta = 2.0\text{--}37.5^\circ$$

$$c = 20.2670 (3) \text{ \AA}$$

$$\mu = 2.17 \text{ mm}^{-1}$$

$$\beta = 98.850 (1)^\circ$$

$$T = 100 \text{ K}$$

$$V = 2012.49 (5) \text{ \AA}^3$$

Block, yellow

$$Z = 4$$

$$0.31 \times 0.27 \times 0.16 \text{ mm}$$

Data collection

Bruker APEXII CCD area-detector diffractometer

10581 independent reflections

Radiation source: sealed tube	8073 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.033$
$T = 100$ K	$\theta_{\text{max}} = 37.5^\circ$
φ and ω scans	$\theta_{\text{min}} = 2.0^\circ$
Absorption correction: multi-scan (SADABS; Bruker, 2005)	$h = -11 \rightarrow 12$
$T_{\text{min}} = 0.548$, $T_{\text{max}} = 0.706$	$k = -22 \rightarrow 22$
67716 measured reflections	$l = -34 \rightarrow 34$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.033$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.085$	$w = 1/[\sigma^2(F_o^2) + (0.0369P)^2 + 0.9196P]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.03$	$(\Delta/\sigma)_{\text{max}} = 0.003$
10581 reflections	$\Delta\rho_{\text{max}} = 1.02 \text{ e \AA}^{-3}$
267 parameters	$\Delta\rho_{\text{min}} = -0.66 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

Special details

Experimental. The crystal was placed in the cold stream of an Oxford Cryosystems Cobra open-flow nitrogen cryostat [Cosier, J. & Glazer, A. M. (1986). *J. Appl. Cryst.* **19**, 105–107.] operating at 100.0 (1) K.

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\text{sigma}(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R- factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	0.437510 (18)	0.545613 (9)	0.317640 (6)	0.02094 (4)
S1	1.03248 (4)	0.87587 (2)	0.276111 (13)	0.01297 (5)
O1	1.08595 (14)	0.92518 (8)	0.33975 (5)	0.02417 (19)
O2	1.17976 (12)	0.82343 (7)	0.25255 (5)	0.02343 (19)
O3	0.93322 (13)	0.94158 (8)	0.22583 (5)	0.0253 (2)
C17	0.87426 (15)	0.78270 (8)	0.29047 (5)	0.01413 (18)
C18	0.92454 (16)	0.68277 (9)	0.30078 (6)	0.01618 (19)

supplementary materials

H18A	1.0438	0.6639	0.3012	0.019*
C19	0.79634 (16)	0.61112 (9)	0.31052 (6)	0.0171 (2)
H19A	0.8285	0.5442	0.3170	0.021*
C20	0.61934 (16)	0.64165 (8)	0.31029 (6)	0.01558 (19)
C21	0.56765 (15)	0.74130 (9)	0.30144 (6)	0.01596 (19)
H21A	0.4491	0.7603	0.3023	0.019*
C22	0.69622 (15)	0.81182 (8)	0.29127 (6)	0.01531 (18)
H22A	0.6637	0.8787	0.2850	0.018*
N1	0.75652 (13)	0.79242 (8)	0.58104 (5)	0.01574 (17)
N2	0.77966 (15)	1.20325 (8)	0.35483 (5)	0.01918 (19)
C1	0.77944 (17)	0.69502 (9)	0.59983 (6)	0.0196 (2)
H1A	0.7581	0.6761	0.6421	0.024*
C2	0.83291 (18)	0.62419 (9)	0.55835 (7)	0.0218 (2)
H2A	0.8490	0.5580	0.5721	0.026*
C3	0.86283 (18)	0.65363 (9)	0.49483 (6)	0.0210 (2)
H3A	0.8972	0.6067	0.4653	0.025*
C4	0.84120 (17)	0.75234 (9)	0.47621 (6)	0.0191 (2)
H4A	0.8621	0.7717	0.4340	0.023*
C5	0.78805 (15)	0.82472 (9)	0.51967 (5)	0.01522 (18)
C6	0.76564 (17)	0.92976 (9)	0.50349 (6)	0.01701 (19)
H6A	0.7261	0.9722	0.5346	0.020*
C7	0.79955 (16)	0.96932 (9)	0.44504 (6)	0.01598 (19)
H7A	0.8379	0.9250	0.4148	0.019*
C8	0.78260 (15)	1.07202 (9)	0.42496 (5)	0.01529 (18)
C9	0.80825 (16)	1.10341 (9)	0.36157 (6)	0.0178 (2)
H9A	0.8404	1.0618	0.3286	0.021*
C10	0.73615 (16)	1.24110 (9)	0.41377 (6)	0.0181 (2)
C11	0.69668 (18)	1.33969 (10)	0.42910 (7)	0.0234 (2)
H11A	0.6958	1.3909	0.3980	0.028*
C12	0.65886 (19)	1.35765 (10)	0.49284 (8)	0.0270 (3)
H12A	0.6305	1.4223	0.5047	0.032*
C13	0.66243 (19)	1.28055 (11)	0.53972 (7)	0.0263 (3)
H13A	0.6384	1.2953	0.5823	0.032*
C14	0.70096 (18)	1.18259 (10)	0.52424 (6)	0.0210 (2)
H14A	0.7027	1.1321	0.5559	0.025*
C15	0.73730 (16)	1.16115 (9)	0.45962 (6)	0.01610 (19)
C16	0.69856 (17)	0.86295 (10)	0.62973 (6)	0.0191 (2)
H16A	0.6809	0.8272	0.6693	0.029*
H16B	0.5877	0.8943	0.6105	0.029*
H16C	0.7894	0.9132	0.6410	0.029*
H1N2	0.796 (2)	1.2377 (15)	0.3209 (10)	0.028 (5)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Br1	0.02422 (6)	0.01477 (5)	0.02501 (6)	-0.00464 (4)	0.00755 (4)	-0.00167 (4)
S1	0.01221 (11)	0.01335 (10)	0.01371 (10)	-0.00058 (8)	0.00315 (8)	-0.00260 (9)
O1	0.0282 (5)	0.0241 (4)	0.0212 (4)	-0.0076 (4)	0.0070 (3)	-0.0102 (4)

O2	0.0167 (4)	0.0236 (4)	0.0319 (5)	0.0008 (3)	0.0096 (3)	-0.0090 (4)
O3	0.0191 (4)	0.0244 (5)	0.0314 (5)	-0.0025 (3)	0.0008 (3)	0.0130 (4)
C17	0.0150 (5)	0.0134 (4)	0.0139 (4)	0.0011 (3)	0.0021 (3)	-0.0009 (3)
C18	0.0155 (5)	0.0147 (4)	0.0185 (4)	0.0037 (4)	0.0031 (3)	0.0000 (4)
C19	0.0210 (5)	0.0130 (4)	0.0177 (4)	0.0029 (4)	0.0044 (4)	0.0003 (4)
C20	0.0187 (5)	0.0128 (4)	0.0156 (4)	-0.0010 (4)	0.0035 (3)	-0.0006 (4)
C21	0.0150 (5)	0.0147 (4)	0.0183 (4)	0.0008 (4)	0.0030 (3)	0.0006 (4)
C22	0.0161 (5)	0.0118 (4)	0.0181 (4)	0.0018 (3)	0.0032 (3)	0.0004 (4)
N1	0.0177 (4)	0.0157 (4)	0.0134 (4)	-0.0026 (3)	0.0012 (3)	0.0000 (3)
N2	0.0209 (5)	0.0179 (4)	0.0184 (4)	-0.0003 (4)	0.0023 (3)	0.0052 (4)
C1	0.0216 (5)	0.0172 (5)	0.0191 (5)	-0.0032 (4)	0.0004 (4)	0.0039 (4)
C2	0.0241 (6)	0.0150 (5)	0.0253 (5)	-0.0007 (4)	0.0004 (4)	0.0023 (4)
C3	0.0229 (6)	0.0165 (5)	0.0230 (5)	0.0023 (4)	0.0020 (4)	-0.0021 (4)
C4	0.0234 (6)	0.0175 (5)	0.0164 (4)	0.0021 (4)	0.0032 (4)	-0.0001 (4)
C5	0.0159 (5)	0.0156 (4)	0.0138 (4)	-0.0009 (4)	0.0011 (3)	0.0007 (4)
C6	0.0207 (5)	0.0151 (4)	0.0157 (4)	0.0013 (4)	0.0044 (4)	0.0006 (4)
C7	0.0180 (5)	0.0153 (4)	0.0145 (4)	0.0004 (4)	0.0020 (3)	0.0002 (4)
C8	0.0157 (5)	0.0152 (4)	0.0148 (4)	0.0000 (4)	0.0017 (3)	0.0008 (4)
C9	0.0180 (5)	0.0186 (5)	0.0166 (4)	0.0000 (4)	0.0025 (4)	0.0017 (4)
C10	0.0161 (5)	0.0164 (5)	0.0210 (5)	-0.0002 (4)	0.0006 (4)	0.0012 (4)
C11	0.0200 (6)	0.0151 (5)	0.0338 (6)	0.0013 (4)	0.0004 (5)	0.0019 (5)
C12	0.0230 (6)	0.0178 (5)	0.0402 (7)	0.0017 (5)	0.0045 (5)	-0.0072 (5)
C13	0.0269 (7)	0.0238 (6)	0.0290 (6)	0.0002 (5)	0.0073 (5)	-0.0085 (5)
C14	0.0241 (6)	0.0197 (5)	0.0197 (5)	-0.0005 (4)	0.0054 (4)	-0.0030 (4)
C15	0.0154 (5)	0.0152 (4)	0.0173 (4)	-0.0001 (4)	0.0016 (3)	0.0002 (4)
C16	0.0233 (6)	0.0200 (5)	0.0143 (4)	-0.0028 (4)	0.0043 (4)	-0.0022 (4)

Geometric parameters (\AA , $^\circ$)

Br1—C20	1.8978 (12)	C3—C4	1.3750 (18)
S1—O1	1.4492 (9)	C3—H3A	0.9300
S1—O2	1.4519 (9)	C4—C5	1.4070 (16)
S1—O3	1.4604 (10)	C4—H4A	0.9300
S1—C17	1.7768 (12)	C5—C6	1.4458 (16)
C17—C18	1.3952 (16)	C6—C7	1.3568 (16)
C17—C22	1.3966 (16)	C6—H6A	0.9300
C18—C19	1.3942 (17)	C7—C8	1.4318 (16)
C18—H18A	0.9300	C7—H7A	0.9300
C19—C20	1.3913 (17)	C8—C9	1.3929 (16)
C19—H19A	0.9300	C8—C15	1.4498 (16)
C20—C21	1.3913 (16)	C9—H9A	0.9300
C21—C22	1.3881 (16)	C10—C11	1.3961 (18)
C21—H21A	0.9300	C10—C15	1.4153 (16)
C22—H22A	0.9300	C11—C12	1.386 (2)
N1—C1	1.3599 (16)	C11—H11A	0.9300
N1—C5	1.3712 (15)	C12—C13	1.399 (2)
N1—C16	1.4782 (16)	C12—H12A	0.9300
N2—C9	1.3552 (16)	C13—C14	1.3874 (19)
N2—C10	1.3821 (16)	C13—H13A	0.9300

supplementary materials

N2—H1N2	0.850 (19)	C14—C15	1.4079 (17)
C1—C2	1.3669 (19)	C14—H14A	0.9300
C1—H1A	0.9300	C16—H16A	0.9600
C2—C3	1.3971 (19)	C16—H16B	0.9600
C2—H2A	0.9300	C16—H16C	0.9600
O1—S1—O2	113.02 (6)	C5—C4—H4A	119.3
O1—S1—O3	112.93 (6)	N1—C5—C4	117.17 (10)
O2—S1—O3	113.24 (6)	N1—C5—C6	118.76 (10)
O1—S1—C17	105.91 (6)	C4—C5—C6	124.07 (10)
O2—S1—C17	106.22 (6)	C7—C6—C5	123.06 (11)
O3—S1—C17	104.63 (5)	C7—C6—H6A	118.5
C18—C17—C22	120.21 (11)	C5—C6—H6A	118.5
C18—C17—S1	121.51 (9)	C6—C7—C8	126.96 (11)
C22—C17—S1	118.27 (8)	C6—C7—H7A	116.5
C19—C18—C17	120.10 (11)	C8—C7—H7A	116.5
C19—C18—H18A	120.0	C9—C8—C7	122.14 (11)
C17—C18—H18A	119.9	C9—C8—C15	105.99 (10)
C20—C19—C18	118.68 (10)	C7—C8—C15	131.86 (10)
C20—C19—H19A	120.7	N2—C9—C8	110.32 (11)
C18—C19—H19A	120.7	N2—C9—H9A	124.8
C21—C20—C19	121.96 (11)	C8—C9—H9A	124.8
C21—C20—Br1	117.86 (9)	N2—C10—C11	128.65 (12)
C19—C20—Br1	120.10 (9)	N2—C10—C15	108.22 (10)
C22—C21—C20	118.82 (11)	C11—C10—C15	123.14 (12)
C22—C21—H21A	120.6	C12—C11—C10	116.88 (12)
C20—C21—H21A	120.6	C12—C11—H11A	121.6
C21—C22—C17	120.21 (10)	C10—C11—H11A	121.6
C21—C22—H22A	119.9	C11—C12—C13	121.35 (12)
C17—C22—H22A	119.9	C11—C12—H12A	119.3
C1—N1—C5	121.55 (10)	C13—C12—H12A	119.3
C1—N1—C16	117.47 (10)	C14—C13—C12	121.62 (13)
C5—N1—C16	120.98 (10)	C14—C13—H13A	119.2
C9—N2—C10	109.17 (10)	C12—C13—H13A	119.2
C9—N2—H1N2	125.1 (13)	C13—C14—C15	118.67 (12)
C10—N2—H1N2	125.5 (13)	C13—C14—H14A	120.7
N1—C1—C2	121.83 (11)	C15—C14—H14A	120.7
N1—C1—H1A	119.1	C14—C15—C10	118.33 (11)
C2—C1—H1A	119.1	C14—C15—C8	135.36 (11)
C1—C2—C3	118.42 (12)	C10—C15—C8	106.29 (10)
C1—C2—H2A	120.8	N1—C16—H16A	109.5
C3—C2—H2A	120.8	N1—C16—H16B	109.5
C4—C3—C2	119.61 (12)	H16A—C16—H16B	109.5
C4—C3—H3A	120.2	N1—C16—H16C	109.5
C2—C3—H3A	120.2	H16A—C16—H16C	109.5
C3—C4—C5	121.40 (11)	H16B—C16—H16C	109.5
C3—C4—H4A	119.3		
O1—S1—C17—C18	-98.92 (10)	C3—C4—C5—C6	179.02 (12)
O2—S1—C17—C18	21.50 (11)	N1—C5—C6—C7	177.69 (11)

O3—S1—C17—C18	141.54 (10)	C4—C5—C6—C7	-2.13 (19)
O1—S1—C17—C22	81.05 (10)	C5—C6—C7—C8	-179.49 (11)
O2—S1—C17—C22	-158.53 (9)	C6—C7—C8—C9	-175.13 (12)
O3—S1—C17—C22	-38.49 (10)	C6—C7—C8—C15	3.9 (2)
C22—C17—C18—C19	1.37 (17)	C10—N2—C9—C8	0.64 (14)
S1—C17—C18—C19	-178.65 (9)	C7—C8—C9—N2	178.35 (11)
C17—C18—C19—C20	-0.57 (17)	C15—C8—C9—N2	-0.92 (14)
C18—C19—C20—C21	-0.76 (17)	C9—N2—C10—C11	179.84 (13)
C18—C19—C20—Br1	175.97 (9)	C9—N2—C10—C15	-0.08 (14)
C19—C20—C21—C22	1.26 (17)	N2—C10—C11—C12	-179.37 (13)
Br1—C20—C21—C22	-175.54 (8)	C15—C10—C11—C12	0.53 (19)
C20—C21—C22—C17	-0.44 (17)	C10—C11—C12—C13	0.8 (2)
C18—C17—C22—C21	-0.86 (17)	C11—C12—C13—C14	-1.1 (2)
S1—C17—C22—C21	179.16 (9)	C12—C13—C14—C15	0.1 (2)
C5—N1—C1—C2	-0.78 (18)	C13—C14—C15—C10	1.18 (18)
C16—N1—C1—C2	-179.93 (11)	C13—C14—C15—C8	179.66 (13)
N1—C1—C2—C3	-0.56 (19)	N2—C10—C15—C14	178.40 (11)
C1—C2—C3—C4	1.17 (19)	C11—C10—C15—C14	-1.52 (18)
C2—C3—C4—C5	-0.48 (19)	N2—C10—C15—C8	-0.48 (13)
C1—N1—C5—C4	1.44 (16)	C11—C10—C15—C8	179.60 (12)
C16—N1—C5—C4	-179.43 (11)	C9—C8—C15—C14	-177.76 (14)
C1—N1—C5—C6	-178.40 (11)	C7—C8—C15—C14	3.1 (2)
C16—N1—C5—C6	0.73 (16)	C9—C8—C15—C10	0.84 (13)
C3—C4—C5—N1	-0.81 (18)	C7—C8—C15—C10	-178.33 (12)

Hydrogen-bond geometry (\AA , °)

$D\cdots H$	$H\cdots A$	$D\cdots A$	$D—H\cdots A$
N2—H1N2···O2 ⁱ	0.85 (2)	1.91 (2)	2.7593 (14) 175.2 (17)
C1—H1A···O3 ⁱⁱ	0.93	2.53	3.2067 (16) 130
C7—H7A···O1	0.93	2.58	3.3095 (16) 136
C9—H9A···O1	0.93	2.58	3.2426 (16) 128
C14—H14A···O1 ⁱⁱⁱ	0.93	2.56	3.2987 (16) 137
C16—H16C···O1 ⁱⁱⁱ	0.96	2.36	3.2739 (17) 158
C19—H19A···O3 ^{iv}	0.93	2.51	3.2052 (16) 131
C21—H21A···O2 ^v	0.93	2.28	3.1310 (15) 152
C4—H4A···Cg3	0.93	2.82	3.5579 (13) 137
C16—H16A···Cg3 ^{vi}	0.96	2.69	3.5731 (13) 154
C16—H16B···Cg1 ^{vii}	0.96	2.74	3.4836 (14) 135

Symmetry codes: (i) $-x+2, y+1/2, -z+1/2$; (ii) $x, -y+3/2, z+1/2$; (iii) $-x+2, -y+2, -z+1$; (iv) $-x+2, y-1/2, -z+1/2$; (v) $x-1, y, z$; (vi) $x, -y+1/2, z-1/2$; (vii) $-x+1, -y+2, -z+1$.

supplementary materials

Fig. 1




Fig. 2

