

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

4,4,5,5-Tetramethyl-2-[4-(2-pyridyl)-phenyl]-3,4-dihydroimidazole-1-oxyl-3-oxide

Xiang-Yang Qin, Ping-An Wang and Xiao-Li Sun*

Department of Chemistry, School of Pharmacy, Fourth Military Medical University, Changle West Road 17, 710032 Xi-An, People's Republic of China

Correspondence e-mail: sun_xiaoli@yahoo.cn


Received 6 April 2009; accepted 7 April 2009

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.070; wR factor = 0.146; data-to-parameter ratio = 13.3.

In the title compound, $\text{C}_{18}\text{H}_{20}\text{N}_3\text{O}_2$, the pyridine and phenyl rings are coplanar [dihedral angle = $3.5(3)^\circ$]. The phenyl ring makes a dihedral angle of $29.6(1)^\circ$ with the imidazole ring. The crystal structure is stabilized by intermolecular $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds.

Related literature

For the preparation of the title compound see: Ullman *et al.* (1974). For recent synthetic use of the title compound and its derivatives, see: Li *et al.* (2009); Xu *et al.* (2008); Masuda *et al.* (2009); Train *et al.* (2009).



Experimental

Crystal data

$\text{C}_{18}\text{H}_{20}\text{N}_3\text{O}_2$
 $M_r = 310.37$
 Monoclinic, $P2_1/c$
 $a = 8.5150(17)$ Å
 $b = 22.286(5)$ Å

$c = 9.1360(18)$ Å
 $\beta = 109.45(3)^\circ$
 $V = 1634.8(6)$ Å³
 $Z = 4$
 Mo $K\alpha$ radiation

$\mu = 0.08$ mm⁻¹
 $T = 293$ K

0.20 × 0.20 × 0.20 mm

Data collection

Bruker SMART APEX CCD diffractometer
 Absorption correction: multi-scan (SADABS; Bruker, 2005)
 $T_{\min} = 0.983$, $T_{\max} = 0.983$

12953 measured reflections
 2819 independent reflections
 1896 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.063$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.070$
 $wR(F^2) = 0.146$
 $S = 1.09$
 2819 reflections

212 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.21$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.20$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{C12}-\text{H12}\cdots\text{O2}^{\ddagger}$	0.93	2.43	3.322 (4)	161

Symmetry code: (i) $-x, y - \frac{1}{2}, -z + \frac{3}{2}$.

Data collection: APEX2 (Bruker, 2007); cell refinement: SAINT (Bruker, 2007); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: Mercury (Macrae *et al.*, 2006) and CAMERON (Watkin *et al.*, 1996).

We thank the Natural Science Foundation of China (grant No. 20802092) for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT2927).

References

- Bruker (2005). SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
 Bruker (2007). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
 Li, R., Li, L., Xing, X. & Liao, D. (2009). *Inorg. Chim. Acta*. In the press. doi:10.1016/j.ica.2008.10.017.
 Macrae, C. F., Edgington, P. R., McCabe, P., Pidcock, E., Shields, G. P., Taylor, R., Towler, M. & van de Streek, J. (2006). *J. Appl. Cryst.* **39**, 453–457.
 Masuda, Y., Kurats, M., Suzuki, S., Kozaki, M., Shiomi, D., Sato, K., Takui, T., Hosokoshi, Y., Lan, X., Miyazaki, Y., Inada, A. & Okada, K. (2009). *J. Am. Chem. Soc.* **131**, 4670–4673.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Train, C., Norel, L. & Baumgarten, M. (2009). *Coord. Chem. Rev.* In the press. doi:10.1016/j.ccr.2008.10.004.
 Ullman, E. F., Osiecki, J. H., Boocock, D. G. B. & Darcy, R. (1974). *J. Am. Chem. Soc.* **96**, 7049–7053.
 Watkin, D. J., Prout, C. K. & Pearce, L. J. (1996). CAMERON. Chemical Crystallography Laboratory, Oxford, England.
 Xu, J., Ma, Y., Xu, G., Wang, C., Liao, D., Jiang, Z., Yan, S. & Li, L. (2008). *Inorg. Chem. Commun.* **11**, 1356–1358.

supplementary materials

Acta Cryst. (2009). E65, o1031 [doi:10.1107/S1600536809013221]

4,4,5,5-Tetramethyl-2-[4-(2-pyridyl)phenyl]-3,4-dihydroimidazole-1-oxyl-3-oxide

X.-Y. Qin, P.-A. Wang and X.-L. Sun

Comment

The title radical compound was obtained the oxidation of 4,4,5,5- tetramethyl-2-(4-(pyridin-2-yl)phenyl)imidazolidine-1,3-diol, which was prepared by the condensation of 4-(pyridin-2-yl)benzaldehyde with 2,3-Dimethyl-2,3-bis(hydroxylamino)butane. The title compound was used for coordination with many metal cations, such as Mn^{2+} , Cu^{2+} , Ni^{2+} and Zn^{2+} , in order to form some molecule-based magnetic materials (Train *et al.*, 2009; Masuda *et al.*, 2009).

In the crystal structure of the title compound, the pyridine ring and the phenyl ring are in one same plane, and this aromatic ring system is twisted with respect to the imidazole ring with a dihedral angle of $29.6(1)^\circ$, and the packing of molecules in the crystal structure is stabilized by intermolecular $C-H\cdots O$ hydrogen bonds. In the imidazole ring, the length of $N1-O1$ is $1.284(3) \text{ \AA}$, while the length of $N2-O2$ is $1.274(3) \text{ \AA}$.

Experimental

The title compound (I) was prepared according to the method reported by Ullman *et al.* (1974). 2,3-Dimethyl-2,3-bis(hydroxylamino)butane (1.48 g, 10.0 mmol) and 4-(pyridin-2-yl)benzaldehyde (1.83 g, 10.0 mmol) were dissolved in a methanol solution (20.0 ml), which was stirred for 3 h at room temperature, and then filtered, the cake was washed by methanol (5.0 ml) for twice. This product was dried under vaccum, then, it was suspended in dichloromethane (100.0 ml) and this reaction mixture was cooled at ice bath for 10 min, the water solution (30.0 ml) of NaO_4 (1.7 g,) was added dropwise to the above suspension and stirred for 20 min at this temperature, the organic layer was seperated and the aqueous phase was extracted by dichloromethane (30.0 ml) for twice. The combined organic layer was dried over Na_2SO_4 and the solvent was removed to give a dark blue residue which was purified by a flash column chromatography (eluent, ether and petroleum ether, the ratio of volume is 4 to 1) to yield the title compound (I) as a dark blue powder. Single crystals of (I) were obtained from the mixed solution of *n*-heptane and dichloromethane (the ratio of volume is 4 to 1).

Refinement

In both structures all the H atoms were discernible in the difference Fourier maps. However, they were constrained by riding model approximation. $C-H_{methyl}=0.96 \text{ \AA}$; $C-H_{aryl}=0.93 \text{ \AA}$; $U_{isoH_{methyl}}$ and $U_{isoH_{aryl}}$ are $1.5 U_{eq}(C)$ and $1.2 U_{eq}(C)$, respectively.

Figures




Fig. 1. The molecular structure of the title compound, showing the atomic labelling scheme. The displacement ellipsoids are drawn at the 50% probability level. The hydrogen atoms are drawn as spheres of arbitrary radius.




Fig. 2. The packing of the title molecules, viewed down the *a* axis. Dotted lines indicate hydrogen bonds.

4,4,5,5-Tetramethyl-2-[4-(2-pyridyl)phenyl]-3,4-dihydroimidazole-1-oxyl-3-oxide

Crystal data

$C_{18}H_{20}N_3O_2$	$F_{000} = 660$
$M_r = 310.37$	$D_x = 1.261 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
$a = 8.5150 (17) \text{ \AA}$	$\lambda = 0.71073 \text{ \AA}$
$b = 22.286 (5) \text{ \AA}$	Cell parameters from 2819 reflections
$c = 9.1360 (18) \text{ \AA}$	$\theta = 3.0\text{--}25.0^\circ$
$\beta = 109.45 (3)^\circ$	$\mu = 0.08 \text{ mm}^{-1}$
$V = 1634.8 (6) \text{ \AA}^3$	$T = 293 \text{ K}$
$Z = 4$	Block, blue
	$0.20 \times 0.20 \times 0.20 \text{ mm}$

Data collection

Bruker SMART APEX CCD diffractometer	2819 independent reflections
Radiation source: fine-focus sealed tube	1896 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.063$
Detector resolution: 0 pixels mm^{-1}	$\theta_{\text{max}} = 25.0^\circ$
$T = 293 \text{ K}$	$\theta_{\text{min}} = 3.0^\circ$
ω scans	$h = -10 \rightarrow 10$
Absorption correction: Multi-scan (<i>SADABS</i> ; Bruker, 2005)	$k = -26 \rightarrow 26$
$T_{\text{min}} = 0.983$, $T_{\text{max}} = 0.983$	$l = -10 \rightarrow 10$
12953 measured reflections	

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.070$	H-atom parameters constrained
$wR(F^2) = 0.146$	$w = 1/[\sigma^2(F_o^2) + (0.0512P)^2 + 0.5834P]$
$S = 1.09$	where $P = (F_o^2 + 2F_c^2)/3$
2819 reflections	$(\Delta/\sigma)_{\text{max}} < 0.001$
212 parameters	$\Delta\rho_{\text{max}} = 0.21 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	$\Delta\rho_{\text{min}} = -0.20 \text{ e \AA}^{-3}$
	Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.0559 (3)	0.57868 (12)	0.7773 (3)	0.0406 (7)
C2	0.3900 (3)	0.72221 (12)	0.7712 (3)	0.0370 (6)
C4	0.3958 (3)	0.66754 (12)	0.6990 (3)	0.0432 (7)
H4	0.4731	0.6620	0.6483	0.052*
C5	0.5033 (3)	0.77122 (11)	0.7688 (3)	0.0377 (6)
C6	0.2755 (3)	0.72855 (12)	0.8480 (3)	0.0469 (7)
H6	0.2695	0.7646	0.8972	0.056*
C7	0.1736 (3)	0.62798 (11)	0.7782 (3)	0.0365 (6)
C8	0.2889 (3)	0.62182 (12)	0.7019 (3)	0.0440 (7)
H8	0.2939	0.5859	0.6516	0.053*
C9	0.1706 (3)	0.68223 (12)	0.8522 (3)	0.0482 (8)
H9	0.0959	0.6873	0.9058	0.058*
C10	0.7488 (3)	0.82333 (12)	0.7781 (3)	0.0422 (7)
C11	0.6108 (3)	0.86972 (12)	0.7722 (3)	0.0412 (7)
C12	-0.1592 (4)	0.48645 (15)	0.7597 (4)	0.0699 (10)
H12	-0.2267	0.4533	0.7561	0.084*
C13	0.8852 (4)	0.81846 (14)	0.9347 (4)	0.0612 (9)
H13A	0.8362	0.8120	1.0140	0.092*
H13B	0.9490	0.8549	0.9558	0.092*
H13C	0.9569	0.7854	0.9333	0.092*
C14	0.8263 (4)	0.82919 (15)	0.6510 (4)	0.0671 (10)
H14A	0.9077	0.7981	0.6628	0.101*
H14B	0.8789	0.8677	0.6585	0.101*
H14C	0.7411	0.8254	0.5514	0.101*
C15	-0.0669 (5)	0.58456 (16)	0.8416 (5)	0.0849 (13)
H15	-0.0773	0.6201	0.8908	0.102*
C16	0.6523 (4)	0.91381 (15)	0.9054 (4)	0.0726 (11)
H16A	0.5576	0.9390	0.8945	0.109*
H16B	0.7450	0.9382	0.9045	0.109*
H16C	0.6808	0.8923	1.0017	0.109*
C18	0.5432 (4)	0.90355 (16)	0.6188 (4)	0.0720 (10)
H18A	0.5157	0.8755	0.5343	0.108*
H18B	0.6262	0.9310	0.6094	0.108*

supplementary materials

H18C	0.4452	0.9254	0.6161	0.108*
C19	-0.0414 (4)	0.48541 (14)	0.6919 (4)	0.0678 (10)
H19	-0.0351	0.4514	0.6352	0.081*
C20	-0.1762 (6)	0.53756 (18)	0.8335 (6)	0.1057 (16)
H20	-0.2595	0.5410	0.8779	0.127*
N1	0.6539 (3)	0.76539 (10)	0.7555 (3)	0.0437 (6)
N2	0.4734 (3)	0.82976 (10)	0.7821 (3)	0.0455 (6)
N3	0.0669 (3)	0.52930 (11)	0.6995 (3)	0.0597 (7)
O1	0.7257 (3)	0.71598 (9)	0.7424 (3)	0.0689 (7)
O2	0.3395 (3)	0.85231 (9)	0.7914 (3)	0.0739 (7)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0425 (16)	0.0374 (17)	0.0416 (17)	0.0011 (12)	0.0136 (14)	0.0010 (13)
C2	0.0350 (15)	0.0354 (16)	0.0390 (16)	0.0010 (12)	0.0101 (13)	0.0015 (12)
C4	0.0427 (16)	0.0451 (18)	0.0463 (18)	-0.0018 (13)	0.0209 (14)	-0.0052 (13)
C5	0.0364 (15)	0.0377 (17)	0.0398 (17)	0.0032 (12)	0.0139 (13)	-0.0007 (12)
C6	0.0516 (18)	0.0376 (17)	0.0558 (19)	-0.0034 (14)	0.0236 (16)	-0.0099 (14)
C7	0.0369 (15)	0.0352 (16)	0.0352 (16)	-0.0006 (11)	0.0089 (13)	0.0004 (12)
C8	0.0514 (18)	0.0369 (16)	0.0439 (18)	-0.0014 (14)	0.0162 (15)	-0.0075 (13)
C9	0.0455 (17)	0.0437 (18)	0.063 (2)	-0.0035 (13)	0.0289 (16)	-0.0064 (14)
C10	0.0409 (16)	0.0403 (16)	0.0478 (18)	-0.0064 (12)	0.0180 (14)	-0.0031 (13)
C11	0.0434 (16)	0.0317 (15)	0.0478 (18)	-0.0064 (12)	0.0141 (14)	-0.0004 (13)
C12	0.076 (2)	0.049 (2)	0.096 (3)	-0.0207 (18)	0.044 (2)	-0.0010 (19)
C13	0.0499 (18)	0.063 (2)	0.063 (2)	-0.0012 (15)	0.0082 (17)	0.0014 (17)
C14	0.086 (3)	0.065 (2)	0.068 (2)	-0.0102 (18)	0.049 (2)	-0.0028 (17)
C15	0.117 (3)	0.056 (2)	0.119 (3)	-0.034 (2)	0.090 (3)	-0.033 (2)
C16	0.059 (2)	0.067 (2)	0.089 (3)	-0.0061 (17)	0.021 (2)	-0.036 (2)
C18	0.068 (2)	0.067 (2)	0.081 (3)	0.0073 (18)	0.025 (2)	0.0286 (19)
C19	0.065 (2)	0.0379 (19)	0.106 (3)	-0.0094 (16)	0.036 (2)	-0.0121 (18)
C20	0.138 (4)	0.079 (3)	0.147 (4)	-0.049 (3)	0.110 (4)	-0.036 (3)
N1	0.0433 (14)	0.0372 (14)	0.0532 (16)	-0.0005 (11)	0.0194 (12)	-0.0037 (11)
N2	0.0397 (14)	0.0350 (14)	0.0647 (17)	0.0016 (11)	0.0212 (12)	0.0003 (11)
N3	0.0569 (17)	0.0402 (16)	0.090 (2)	-0.0082 (12)	0.0358 (15)	-0.0159 (13)
O1	0.0569 (14)	0.0418 (13)	0.121 (2)	0.0004 (10)	0.0471 (14)	-0.0134 (12)
O2	0.0542 (14)	0.0416 (13)	0.139 (2)	0.0080 (11)	0.0504 (14)	0.0038 (12)

Geometric parameters (\AA , $^\circ$)

C1—N3	1.331 (3)	C12—C19	1.343 (4)
C1—C15	1.365 (4)	C12—C20	1.356 (5)
C1—C7	1.486 (4)	C12—H12	0.9300
C2—C6	1.385 (3)	C13—H13A	0.9600
C2—C4	1.394 (4)	C13—H13B	0.9600
C2—C5	1.463 (4)	C13—H13C	0.9600
C4—C8	1.373 (4)	C14—H14A	0.9600
C4—H4	0.9300	C14—H14B	0.9600
C5—N1	1.334 (3)	C14—H14C	0.9600

C5—N2	1.343 (3)	C15—C20	1.387 (5)
C6—C9	1.374 (4)	C15—H15	0.9300
C6—H6	0.9300	C16—H16A	0.9600
C7—C8	1.388 (4)	C16—H16B	0.9600
C7—C9	1.389 (4)	C16—H16C	0.9600
C8—H8	0.9300	C18—H18A	0.9600
C9—H9	0.9300	C18—H18B	0.9600
C10—N1	1.501 (3)	C18—H18C	0.9600
C10—C13	1.517 (4)	C19—N3	1.331 (4)
C10—C14	1.520 (4)	C19—H19	0.9300
C10—C11	1.552 (4)	C20—H20	0.9300
C11—N2	1.497 (3)	N1—O1	1.284 (3)
C11—C16	1.512 (4)	N2—O2	1.274 (3)
C11—C18	1.525 (4)		
N3—C1—C15	120.7 (3)	C10—C13—H13B	109.5
N3—C1—C7	116.5 (2)	H13A—C13—H13B	109.5
C15—C1—C7	122.6 (3)	C10—C13—H13C	109.5
C6—C2—C4	118.1 (2)	H13A—C13—H13C	109.5
C6—C2—C5	120.7 (2)	H13B—C13—H13C	109.5
C4—C2—C5	121.2 (2)	C10—C14—H14A	109.5
C8—C4—C2	120.8 (3)	C10—C14—H14B	109.5
C8—C4—H4	119.6	H14A—C14—H14B	109.5
C2—C4—H4	119.6	C10—C14—H14C	109.5
N1—C5—N2	108.7 (2)	H14A—C14—H14C	109.5
N1—C5—C2	126.0 (2)	H14B—C14—H14C	109.5
N2—C5—C2	125.3 (2)	C1—C15—C20	120.1 (3)
C9—C6—C2	120.8 (3)	C1—C15—H15	120.0
C9—C6—H6	119.6	C20—C15—H15	120.0
C2—C6—H6	119.6	C11—C16—H16A	109.5
C8—C7—C9	117.5 (2)	C11—C16—H16B	109.5
C8—C7—C1	120.9 (2)	H16A—C16—H16B	109.5
C9—C7—C1	121.6 (2)	C11—C16—H16C	109.5
C4—C8—C7	121.4 (2)	H16A—C16—H16C	109.5
C4—C8—H8	119.3	H16B—C16—H16C	109.5
C7—C8—H8	119.3	C11—C18—H18A	109.5
C6—C9—C7	121.5 (3)	C11—C18—H18B	109.5
C6—C9—H9	119.3	H18A—C18—H18B	109.5
C7—C9—H9	119.3	C11—C18—H18C	109.5
N1—C10—C13	106.1 (2)	H18A—C18—H18C	109.5
N1—C10—C14	108.6 (2)	H18B—C18—H18C	109.5
C13—C10—C14	109.6 (2)	N3—C19—C12	125.1 (3)
N1—C10—C11	101.6 (2)	N3—C19—H19	117.5
C13—C10—C11	114.5 (2)	C12—C19—H19	117.5
C14—C10—C11	115.6 (2)	C12—C20—C15	118.7 (3)
N2—C11—C16	108.4 (2)	C12—C20—H20	120.7
N2—C11—C18	106.6 (2)	C15—C20—H20	120.7
C16—C11—C18	109.5 (3)	O1—N1—C5	126.4 (2)
N2—C11—C10	101.5 (2)	O1—N1—C10	120.0 (2)
C16—C11—C10	115.7 (2)	C5—N1—C10	113.2 (2)

supplementary materials

C18—C11—C10	114.3 (2)	O2—N2—C5	126.2 (2)
C19—C12—C20	117.6 (3)	O2—N2—C11	120.3 (2)
C19—C12—H12	121.2	C5—N2—C11	113.4 (2)
C20—C12—H12	121.2	C19—N3—C1	117.7 (3)
C10—C13—H13A	109.5		
C6—C2—C4—C8	1.1 (4)	C7—C1—C15—C20	178.2 (4)
C5—C2—C4—C8	-180.0 (2)	C20—C12—C19—N3	4.3 (6)
C6—C2—C5—N1	152.3 (3)	C19—C12—C20—C15	-3.0 (7)
C4—C2—C5—N1	-26.6 (4)	C1—C15—C20—C12	-0.7 (7)
C6—C2—C5—N2	-26.5 (4)	N2—C5—N1—O1	179.1 (2)
C4—C2—C5—N2	154.6 (3)	C2—C5—N1—O1	0.1 (4)
C4—C2—C6—C9	-0.1 (4)	N2—C5—N1—C10	6.8 (3)
C5—C2—C6—C9	-179.1 (3)	C2—C5—N1—C10	-172.3 (2)
N3—C1—C7—C8	-0.4 (4)	C13—C10—N1—O1	-65.1 (3)
C15—C1—C7—C8	-175.3 (3)	C14—C10—N1—O1	52.6 (3)
N3—C1—C7—C9	178.3 (3)	C11—C10—N1—O1	175.0 (2)
C15—C1—C7—C9	3.4 (4)	C13—C10—N1—C5	107.8 (3)
C2—C4—C8—C7	-1.0 (4)	C14—C10—N1—C5	-134.5 (3)
C9—C7—C8—C4	-0.2 (4)	C11—C10—N1—C5	-12.1 (3)
C1—C7—C8—C4	178.5 (3)	N1—C5—N2—O2	177.4 (3)
C2—C6—C9—C7	-1.0 (4)	C2—C5—N2—O2	-3.6 (4)
C8—C7—C9—C6	1.1 (4)	N1—C5—N2—C11	2.2 (3)
C1—C7—C9—C6	-177.5 (3)	C2—C5—N2—C11	-178.8 (2)
N1—C10—C11—N2	11.7 (2)	C16—C11—N2—O2	52.8 (3)
C13—C10—C11—N2	-102.1 (2)	C18—C11—N2—O2	-65.0 (3)
C14—C10—C11—N2	129.1 (2)	C10—C11—N2—O2	175.1 (2)
N1—C10—C11—C16	128.8 (3)	C16—C11—N2—C5	-131.6 (3)
C13—C10—C11—C16	15.0 (4)	C18—C11—N2—C5	110.5 (3)
C14—C10—C11—C16	-113.7 (3)	C10—C11—N2—C5	-9.4 (3)
N1—C10—C11—C18	-102.6 (3)	C12—C19—N3—C1	-1.6 (5)
C13—C10—C11—C18	143.6 (3)	C15—C1—N3—C19	-2.5 (5)
C14—C10—C11—C18	14.8 (3)	C7—C1—N3—C19	-177.5 (3)
N3—C1—C15—C20	3.5 (6)		

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C12—H12 \cdots O2 ⁱ	0.93	2.43	3.322 (4)	161

Symmetry codes: (i) $-x, y-1/2, -z+3/2$.

Fig. 1




Fig. 2

