

## Phenyl *N*-(4-fluorophenyl)carbamate

**Zhao Yang and Zhi-Xiang Wang\***

 Department of Pharmaceutical Engineering, China Pharmaceutical University,  
 Tongjiaxiang No. 24 Nanjing, Nanjing 210009, People's Republic of China  
 Correspondence e-mail: yzcpu@163.com


Received 7 April 2009; accepted 8 April 2009

 Key indicators: single-crystal X-ray study;  $T = 294$  K; mean  $\sigma(\text{C}-\text{C}) = 0.007$  Å;  
 $R$  factor = 0.073;  $wR$  factor = 0.157; data-to-parameter ratio = 13.4.

The asymmetric unit of the title compound,  $\text{C}_{13}\text{H}_{10}\text{FNO}_2$ , contains two crystallographically independent molecules. The aromatic rings are oriented at dihedral angles of  $61.77$  (3) and  $53.94$  (3)° in the two molecules. An  $\text{N}-\text{H}\cdots\text{O}$  hydrogen bond links the molecules. In the crystal structure, intermolecular  $\text{N}-\text{H}\cdots\text{O}$  hydrogen bonds link the molecules into chains. Weak  $\text{C}-\text{H}\cdots\pi$  interactions are also present.

### Related literature

For a related structure, see: Hynes *et al.* (2008). For bond-length data, see: Allen *et al.* (1987).



### Experimental

#### Crystal data

$\text{C}_{13}\text{H}_{10}\text{FNO}_2$	$\gamma = 91.19$ (3)°
$M_r = 231.22$	$V = 1135.3$ (4) Å <sup>3</sup>
Triclinic, $P\bar{1}$	$Z = 4$
$a = 5.8860$ (12) Å	Mo $K\alpha$ radiation
$b = 7.8540$ (16) Å	$\mu = 0.10$ mm <sup>-1</sup>
$c = 24.761$ (5) Å	$T = 294$ K
$\alpha = 96.62$ (3)°	$0.10 \times 0.10 \times 0.08$ mm
$\beta = 92.82$ (3)°	

#### Data collection

Enraf–Nonius CAD-4 diffractometer	4110 independent reflections
Absorption correction: $\psi$ scan (North <i>et al.</i> , 1968)	1662 reflections with $I > 2\sigma(I)$
$T_{\text{min}} = 0.990$ , $T_{\text{max}} = 0.992$	$R_{\text{int}} = 0.060$
4548 measured reflections	3 standard reflections
	frequency: 120 min
	intensity decay: 1%

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.073$	307 parameters
$wR(F^2) = 0.157$	H-atom parameters constrained
$S = 1.00$	$\Delta\rho_{\text{max}} = 0.13$ e Å <sup>-3</sup>
4110 reflections	$\Delta\rho_{\text{min}} = -0.13$ e Å <sup>-3</sup>

**Table 1**

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N1}-\text{H1A}\cdots\text{O4}^{\text{i}}$	0.86	2.32	3.044 (4)	142
$\text{N2}-\text{H2C}\cdots\text{O2}$	0.86	2.08	2.931 (4)	171
$\text{C19}-\text{H19A}\cdots\text{Cg2}^{\text{ii}}$	0.93	2.94	3.644 (4)	134
$\text{C23}-\text{H23A}\cdots\text{Cg1}^{\text{iii}}$	0.93	2.97	3.710 (5)	138

Symmetry codes: (i)  $x - 1, y - 1, z$ ; (ii)  $x, y + 1, z$ ; (iii)  $x + 1, y + 1, z$ .  $\text{Cg1}$  and  $\text{Cg2}$  are the centroids of the  $\text{C1}-\text{C6}$  and  $\text{C8}-\text{C13}$  rings, respectively.

Data collection: *CAD-4 Software* (Enraf–Nonius, 1989); cell refinement: *CAD-4 Software*; data reduction: *XCAD4* (Harms & Wocadlo, 1995); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2009); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008).

The authors thank the Center of Testing and Analysis, Nanjing University, for support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HK2663).

### References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Enraf–Nonius (1989). *CAD-4 Software*. Enraf–Nonius, Delft, The Netherlands.
- Harms, K. & Wocadlo, S. (1995). *XCAD4*. University of Marburg, Germany.
- Hynes, J. J., Dyckman, A. J., Lin, S. W. & Stephen, T. (2008). *J. Med. Chem.* **51**, 4–16.
- North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst.* **A24**, 351–359.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

**supplementary materials**

*Acta Cryst.* (2009). E65, o1036 [ doi:10.1107/S1600536809013312 ]

## Phenyl *N*-(4-fluorophenyl)carbamate

Z. Yang and Z.-X. Wang

### Comment

Some derivatives of aniline are important chemical materials. We report herein the crystal structure of the title compound.

The asymmetric unit of the title compound contains two crystallographically independent molecules (Fig. 1), in which the bond lengths (Allen *et al.*, 1987) and angles are within normal ranges. Rings A (C1-C6), B (C8-C13) and A' (C14-C19), B' (C21-C26) are, of course, planar, and they are oriented at dihedral angles of A/B = 61.77 (3) and A'/B' = 53.94 (3) °. Intramolecular N-H...O hydrogen bond (Table 1) links the molecules.

In the crystal structure, intra- and intermolecular N-H...O hydrogen bonds (Table 1) link the molecules into chains (Fig. 2), in which they may be effective in the stabilization of the structure. There also exist weak C—H... $\pi$  interactions (Table 1).

### Experimental

For the preparation of the title compound, phenyl chloroformate (1.0 ml) was added slowly to a cold solution of 4-fluorbenzenamine (1.0 g) and triethylamine (0.8 ml) in methylene chloride (10 ml) at 273 K. The mixture was then warmed and stirred for 1 h at room temperature. It was washed with water (20 ml), dried and concentrated to give the title compound (yield; 1.3 g) (Hynes *et al.*, 2008). Crystals suitable for X-ray analysis were obtained by slow evaporation of a methanol solution.

### Refinement

H atoms were positioned geometrically, with N-H = 0.86 Å (for NH) and C-H = 0.93 Å for aromatic H and constrained to ride on their parent atoms, with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C,N})$ .

### Figures




Fig. 1. The molecular structure of the title molecule, with the atom-numbering scheme. Hydrogen bond is shown as dashed line.




Fig. 2. A partial packing diagram of the title compound. Hydrogen bonds are shown as dashed lines.

### Phenyl *N*-(4-fluorophenyl)carbamate

#### Crystal data

$C_{13}H_{10}FNO_2$

$M_r = 231.22$

Triclinic,  $P\bar{1}$

Hall symbol:  $-P\ 1$

$a = 5.8860\ (12)\ \text{\AA}$

$b = 7.8540\ (16)\ \text{\AA}$

$c = 24.761\ (5)\ \text{\AA}$

$\alpha = 96.62\ (3)^\circ$

$\beta = 92.82\ (3)^\circ$

$\gamma = 91.19\ (3)^\circ$

$V = 1135.3\ (4)\ \text{\AA}^3$

$Z = 4$

$F_{000} = 480$

$D_x = 1.353\ \text{Mg m}^{-3}$

Mo  $K\alpha$  radiation

$\lambda = 0.71073\ \text{\AA}$

Cell parameters from 25 reflections

$\theta = 8\text{--}12^\circ$

$\mu = 0.10\ \text{mm}^{-1}$

$T = 294\ \text{K}$

Block, colorless

$0.10 \times 0.10 \times 0.08\ \text{mm}$

#### Data collection

Enraf–Nonius CAD-4  
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 294\ \text{K}$

$\omega/2\theta$  scans

Absorption correction:  $\psi$  scan  
(North *et al.*, 1968)

$T_{\min} = 0.990$ ,  $T_{\max} = 0.992$

4548 measured reflections

4110 independent reflections

1662 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.060$

$\theta_{\max} = 25.3^\circ$

$\theta_{\min} = 1.7^\circ$

$h = 0 \rightarrow 7$

$k = -9 \rightarrow 9$

$l = -29 \rightarrow 29$

3 standard reflections

every 120 min

intensity decay: 1%

#### Refinement

Refinement on  $F^2$

Least-squares matrix: full

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

$$R[F^2 > 2\sigma(F^2)] = 0.073$$

$$wR(F^2) = 0.157$$

$$S = 1.00$$

4110 reflections

307 parameters

Primary atom site location: structure-invariant direct methods

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.052P)^2]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} < 0.001$$

$$\Delta\rho_{\max} = 0.13 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.13 \text{ e } \text{\AA}^{-3}$$

Extinction correction: none

### Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
N1	0.0390 (6)	-0.6473 (4)	0.26993 (14)	0.0831 (11)
H1A	-0.0691	-0.7213	0.2599	0.100*
N2	0.6939 (5)	-0.2644 (4)	0.22504 (13)	0.0755 (11)
H2C	0.6083	-0.3507	0.2300	0.091*
F1	-0.0366 (5)	-0.3165 (3)	0.47653 (11)	0.1065 (9)
F2	1.1428 (5)	-0.3284 (4)	0.03482 (12)	0.1253 (11)
O1	0.1510 (5)	-0.7508 (4)	0.19019 (13)	0.1045 (12)
O2	0.3575 (5)	-0.5355 (4)	0.23766 (11)	0.0916 (10)
O3	0.5300 (5)	-0.1536 (4)	0.29759 (12)	0.0938 (11)
O4	0.8164 (4)	-0.0021 (3)	0.26628 (10)	0.0724 (8)
C1	0.3835 (12)	-0.7830 (7)	0.0572 (2)	0.1033 (18)
H1B	0.3416	-0.7637	0.0218	0.124*
C2	0.2332 (9)	-0.7392 (6)	0.0984 (3)	0.0934 (16)
H2A	0.0941	-0.6896	0.0917	0.112*
C3	0.3045 (9)	-0.7740 (6)	0.1487 (2)	0.0763 (14)
C4	0.5077 (11)	-0.8420 (6)	0.1599 (2)	0.0894 (16)
H4A	0.5516	-0.8605	0.1953	0.107*
C5	0.6465 (9)	-0.8828 (6)	0.1186 (3)	0.1024 (18)
H5A	0.7849	-0.9327	0.1259	0.123*
C6	0.5889 (11)	-0.8528 (7)	0.0673 (3)	0.108 (2)
H6A	0.6871	-0.8792	0.0394	0.130*
C7	0.1980 (8)	-0.6340 (6)	0.23447 (19)	0.0860 (15)
C8	0.0241 (7)	-0.5556 (5)	0.32213 (17)	0.0698 (12)
C9	-0.1636 (7)	-0.5849 (5)	0.35086 (18)	0.0740 (12)

## supplementary materials

---

H9A	-0.2782	-0.6603	0.3349	0.089*
C10	-0.1859 (8)	-0.5051 (6)	0.4028 (2)	0.0861 (14)
H10A	-0.3134	-0.5253	0.4221	0.103*
C11	-0.0157 (8)	-0.3963 (5)	0.42483 (19)	0.0776 (13)
C12	0.1691 (8)	-0.3626 (5)	0.39833 (18)	0.0772 (13)
H12A	0.2818	-0.2868	0.4150	0.093*
C13	0.1923 (7)	-0.4407 (5)	0.34595 (18)	0.0784 (13)
H13A	0.3192	-0.4165	0.3269	0.094*
C14	0.4753 (10)	0.1478 (6)	0.4438 (2)	0.0860 (15)
H14A	0.4594	0.2114	0.4774	0.103*
C15	0.6668 (9)	0.0580 (6)	0.4346 (2)	0.0882 (15)
H15A	0.7815	0.0618	0.4620	0.106*
C16	0.6942 (7)	-0.0391 (5)	0.3851 (2)	0.0744 (13)
H16A	0.8265	-0.0994	0.3790	0.089*
C17	0.5254 (8)	-0.0445 (5)	0.34616 (18)	0.0647 (11)
C18	0.3329 (8)	0.0490 (6)	0.35436 (18)	0.0775 (13)
H18A	0.2203	0.0478	0.3266	0.093*
C19	0.3070 (9)	0.1455 (6)	0.4044 (2)	0.0829 (14)
H19A	0.1761	0.2075	0.4107	0.099*
C20	0.6969 (7)	-0.1294 (6)	0.26345 (17)	0.0763 (13)
C21	0.8171 (7)	-0.2801 (5)	0.17706 (16)	0.0588 (10)
C22	1.0269 (7)	-0.2011 (5)	0.17460 (18)	0.0738 (12)
H22A	1.0954	-0.1378	0.2054	0.089*
C23	1.1343 (7)	-0.2167 (5)	0.12616 (19)	0.0798 (13)
H23A	1.2738	-0.1612	0.1237	0.096*
C24	1.0365 (9)	-0.3124 (6)	0.08250 (19)	0.0840 (14)
C25	0.8320 (8)	-0.3995 (5)	0.08437 (18)	0.0813 (13)
H25A	0.7693	-0.4681	0.0540	0.098*
C26	0.7227 (7)	-0.3818 (5)	0.13273 (17)	0.0717 (12)
H26A	0.5845	-0.4393	0.1352	0.086*

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
N1	0.078 (3)	0.080 (3)	0.086 (3)	-0.031 (2)	0.023 (2)	-0.013 (2)
N2	0.082 (3)	0.062 (2)	0.080 (3)	-0.038 (2)	0.023 (2)	-0.0024 (19)
F1	0.120 (2)	0.102 (2)	0.094 (2)	-0.0065 (17)	0.0279 (17)	-0.0139 (16)
F2	0.140 (3)	0.121 (2)	0.112 (2)	-0.0283 (19)	0.061 (2)	-0.0174 (18)
O1	0.105 (3)	0.109 (3)	0.091 (2)	-0.056 (2)	0.030 (2)	-0.027 (2)
O2	0.094 (2)	0.085 (2)	0.091 (2)	-0.0490 (18)	0.0269 (18)	-0.0119 (17)
O3	0.099 (2)	0.086 (2)	0.089 (2)	-0.0449 (18)	0.0397 (19)	-0.0265 (18)
O4	0.0745 (19)	0.0645 (18)	0.0752 (19)	-0.0264 (15)	0.0099 (15)	-0.0018 (14)
C1	0.112 (5)	0.125 (5)	0.071 (4)	-0.036 (4)	-0.012 (4)	0.014 (3)
C2	0.081 (4)	0.089 (4)	0.112 (5)	-0.014 (3)	-0.010 (4)	0.030 (3)
C3	0.065 (3)	0.070 (3)	0.090 (4)	-0.027 (3)	0.014 (3)	-0.005 (3)
C4	0.099 (4)	0.083 (4)	0.083 (4)	-0.035 (3)	-0.008 (3)	0.008 (3)
C5	0.080 (4)	0.063 (3)	0.163 (6)	-0.007 (3)	-0.009 (5)	0.013 (4)
C6	0.081 (4)	0.102 (5)	0.134 (6)	-0.029 (4)	0.032 (4)	-0.025 (4)

C7	0.081 (4)	0.093 (4)	0.077 (3)	-0.033 (3)	0.015 (3)	-0.015 (3)
C8	0.069 (3)	0.062 (3)	0.079 (3)	-0.017 (2)	0.015 (2)	0.008 (2)
C9	0.066 (3)	0.067 (3)	0.087 (3)	-0.016 (2)	0.010 (2)	0.002 (2)
C10	0.083 (3)	0.078 (3)	0.097 (4)	-0.013 (3)	0.025 (3)	0.006 (3)
C11	0.081 (4)	0.061 (3)	0.089 (4)	0.005 (3)	0.024 (3)	-0.007 (3)
C12	0.081 (3)	0.074 (3)	0.076 (3)	-0.015 (3)	0.000 (3)	0.009 (3)
C13	0.075 (3)	0.077 (3)	0.082 (3)	-0.030 (2)	0.009 (2)	0.006 (3)
C14	0.093 (4)	0.086 (4)	0.078 (4)	-0.010 (3)	0.020 (3)	0.002 (3)
C15	0.090 (4)	0.098 (4)	0.077 (4)	-0.012 (3)	0.002 (3)	0.018 (3)
C16	0.055 (3)	0.064 (3)	0.108 (4)	-0.001 (2)	0.015 (3)	0.020 (3)
C17	0.065 (3)	0.057 (3)	0.072 (3)	-0.014 (2)	0.012 (3)	0.005 (2)
C18	0.077 (4)	0.078 (3)	0.079 (3)	-0.020 (3)	-0.004 (3)	0.021 (3)
C19	0.080 (4)	0.069 (3)	0.102 (4)	0.013 (3)	0.022 (3)	0.008 (3)
C20	0.081 (3)	0.070 (3)	0.076 (3)	-0.028 (3)	0.017 (3)	-0.003 (3)
C21	0.059 (3)	0.048 (2)	0.068 (3)	-0.007 (2)	0.003 (2)	0.002 (2)
C22	0.054 (3)	0.073 (3)	0.090 (3)	-0.018 (2)	0.002 (2)	-0.005 (2)
C23	0.068 (3)	0.070 (3)	0.099 (4)	-0.010 (2)	0.029 (3)	-0.007 (3)
C24	0.099 (4)	0.070 (3)	0.082 (4)	-0.006 (3)	0.032 (3)	-0.006 (3)
C25	0.093 (4)	0.073 (3)	0.074 (3)	-0.006 (3)	0.009 (3)	-0.008 (2)
C26	0.072 (3)	0.065 (3)	0.074 (3)	-0.018 (2)	0.007 (2)	-0.007 (2)

*Geometric parameters (Å, °)*

N1—C7	1.325 (5)	C9—H9A	0.9300
N1—C8	1.413 (4)	C10—C11	1.354 (5)
N1—H1A	0.8600	C10—H10A	0.9300
N2—C20	1.340 (4)	C11—C12	1.335 (5)
N2—C21	1.417 (4)	C12—C13	1.384 (5)
N2—H2C	0.8600	C12—H12A	0.9300
F1—C11	1.371 (4)	C13—H13A	0.9300
F2—C24	1.358 (4)	C14—C19	1.354 (6)
O1—C3	1.399 (5)	C14—C15	1.357 (6)
O1—C7	1.359 (4)	C14—H14A	0.9300
O2—C7	1.198 (4)	C15—C16	1.385 (6)
O3—C17	1.396 (5)	C15—H15A	0.9300
O3—C20	1.351 (4)	C16—C17	1.347 (5)
O4—C20	1.204 (4)	C16—H16A	0.9300
C1—C6	1.361 (7)	C17—C18	1.374 (5)
C1—C2	1.400 (6)	C18—C19	1.393 (6)
C1—H1B	0.9300	C18—H18A	0.9300
C2—C3	1.355 (6)	C19—H19A	0.9300
C2—H2A	0.9300	C21—C26	1.368 (5)
C3—C4	1.349 (6)	C21—C22	1.377 (5)
C4—C5	1.353 (6)	C22—C23	1.377 (5)
C4—H4A	0.9300	C22—H22A	0.9300
C5—C6	1.348 (7)	C23—C24	1.341 (5)
C5—H5A	0.9300	C23—H23A	0.9300
C6—H6A	0.9300	C24—C25	1.377 (6)
C8—C9	1.373 (5)	C25—C26	1.381 (5)

## supplementary materials

---

C8—C13	1.388 (5)	C25—H25A	0.9300
C9—C10	1.379 (5)	C26—H26A	0.9300
C7—N1—C8	128.3 (4)	C13—C12—H12A	120.1
C7—N1—H1A	115.8	C12—C13—C8	119.2 (4)
C8—N1—H1A	115.8	C12—C13—H13A	120.4
C20—N2—C21	126.8 (3)	C8—C13—H13A	120.4
C20—N2—H2C	116.6	C19—C14—C15	120.5 (5)
C21—N2—H2C	116.6	C19—C14—H14A	119.7
C7—O1—C3	120.4 (3)	C15—C14—H14A	119.7
C20—O3—C17	118.5 (3)	C14—C15—C16	121.0 (5)
C6—C1—C2	122.3 (5)	C14—C15—H15A	119.5
C6—C1—H1B	118.9	C16—C15—H15A	119.5
C2—C1—H1B	118.9	C17—C16—C15	118.7 (4)
C3—C2—C1	115.4 (5)	C17—C16—H16A	120.6
C3—C2—H2A	122.3	C15—C16—H16A	120.6
C1—C2—H2A	122.3	C16—C17—C18	120.9 (4)
C4—C3—C2	123.5 (5)	C16—C17—O3	121.9 (5)
C4—C3—O1	118.5 (6)	C18—C17—O3	117.0 (4)
C2—C3—O1	117.8 (6)	C17—C18—C19	119.7 (4)
C3—C4—C5	118.9 (5)	C17—C18—H18A	120.1
C3—C4—H4A	120.6	C19—C18—H18A	120.1
C5—C4—H4A	120.6	C14—C19—C18	119.0 (5)
C6—C5—C4	121.5 (6)	C14—C19—H19A	120.5
C6—C5—H5A	119.2	C18—C19—H19A	120.5
C4—C5—H5A	119.2	O4—C20—N2	127.1 (4)
C5—C6—C1	118.4 (6)	O4—C20—O3	124.4 (4)
C5—C6—H6A	120.8	N2—C20—O3	108.3 (3)
C1—C6—H6A	120.8	C26—C21—C22	120.3 (4)
O2—C7—N1	128.5 (4)	C26—C21—N2	117.4 (3)
O2—C7—O1	122.5 (4)	C22—C21—N2	122.3 (4)
N1—C7—O1	109.0 (4)	C21—C22—C23	119.4 (4)
C9—C8—C13	118.8 (4)	C21—C22—H22A	120.3
C9—C8—N1	118.1 (4)	C23—C22—H22A	120.3
C13—C8—N1	123.0 (4)	C24—C23—C22	119.6 (4)
C8—C9—C10	121.4 (4)	C24—C23—H23A	120.2
C8—C9—H9A	119.3	C22—C23—H23A	120.2
C10—C9—H9A	119.3	C23—C24—F2	119.6 (4)
C11—C10—C9	117.7 (4)	C23—C24—C25	122.3 (4)
C11—C10—H10A	121.2	F2—C24—C25	118.1 (4)
C9—C10—H10A	121.2	C24—C25—C26	118.1 (4)
C12—C11—C10	123.0 (4)	C24—C25—H25A	121.0
C12—C11—F1	118.9 (4)	C26—C25—H25A	121.0
C10—C11—F1	118.1 (4)	C21—C26—C25	120.2 (4)
C11—C12—C13	119.8 (4)	C21—C26—H26A	119.9
C11—C12—H12A	120.1	C25—C26—H26A	119.9
C6—C1—C2—C3	-1.2 (7)	C19—C14—C15—C16	-0.6 (7)
C1—C2—C3—C4	1.7 (7)	C14—C15—C16—C17	-0.7 (6)
C1—C2—C3—O1	-173.0 (4)	C15—C16—C17—C18	2.3 (6)



C7—O1—C3—C4	67.3 (6)	C15—C16—C17—O3	-173.0 (3)
C7—O1—C3—C2	-117.7 (5)	C20—O3—C17—C16	-63.8 (6)
C2—C3—C4—C5	-2.0 (7)	C20—O3—C17—C18	120.8 (4)
O1—C3—C4—C5	172.7 (4)	C16—C17—C18—C19	-2.6 (6)
C3—C4—C5—C6	1.8 (7)	O3—C17—C18—C19	172.9 (4)
C4—C5—C6—C1	-1.3 (8)	C15—C14—C19—C18	0.3 (7)
C2—C1—C6—C5	1.1 (8)	C17—C18—C19—C14	1.3 (6)
C8—N1—C7—O2	-4.2 (9)	C21—N2—C20—O4	-5.8 (8)
C8—N1—C7—O1	177.7 (4)	C21—N2—C20—O3	170.2 (4)
C3—O1—C7—O2	7.6 (8)	C17—O3—C20—O4	-15.0 (7)
C3—O1—C7—N1	-174.2 (4)	C17—O3—C20—N2	168.9 (4)
C7—N1—C8—C9	176.8 (5)	C20—N2—C21—C26	-150.6 (4)
C7—N1—C8—C13	-4.7 (7)	C20—N2—C21—C22	31.2 (6)
C13—C8—C9—C10	-1.1 (7)	C26—C21—C22—C23	4.0 (6)
N1—C8—C9—C10	177.5 (4)	N2—C21—C22—C23	-177.8 (4)
C8—C9—C10—C11	-0.1 (7)	C21—C22—C23—C24	-1.9 (7)
C9—C10—C11—C12	0.6 (7)	C22—C23—C24—F2	-179.7 (4)
C9—C10—C11—F1	-180.0 (4)	C22—C23—C24—C25	-1.2 (8)
C10—C11—C12—C13	0.0 (8)	C23—C24—C25—C26	2.1 (7)
F1—C11—C12—C13	-179.4 (4)	F2—C24—C25—C26	-179.4 (4)
C11—C12—C13—C8	-1.1 (7)	C22—C21—C26—C25	-3.1 (6)
C9—C8—C13—C12	1.6 (7)	N2—C21—C26—C25	178.6 (4)
N1—C8—C13—C12	-176.9 (4)	C24—C25—C26—C21	0.1 (7)

Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ )

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N1—H1A $\cdots$ O4 <sup>i</sup>	0.86	2.32	3.044 (4)	142
N2—H2C $\cdots$ O2	0.86	2.08	2.931 (4)	171
C19—H19A $\cdots$ Cg2 <sup>ii</sup>	0.93	2.94	3.644 (4)	134
C23—H23A $\cdots$ Cg1 <sup>iii</sup>	0.93	2.97	3.710 (5)	138

Symmetry codes: (i)  $x-1, y-1, z$ ; (ii)  $x, y+1, z$ ; (iii)  $x+1, y+1, z$ .

Fig. 1




Fig. 2

