

2-[3-Acetyl-5-(2-chloro-3-pyridyl)-2-methyl-2,3-dihydro-1,3,4-oxadiazol-2-yl]-4-fluorophenyl acetate

Quan Qin, Li Juan Xu, Li Fang Pan, Shui Qing Chen and
Qing Bao Song*

The State Key Laboratory Breeding Base of Green Chemistry–Synthesis Technology, College of Chemical Engineering and Materials Science, Zhejiang University of Technology, Hangzhou 310014, People's Republic of China
Correspondence e-mail: qbSong@zjut.edu.cn


Received 30 March 2009; accepted 24 April 2009

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$;
 R factor = 0.043; wR factor = 0.121; data-to-parameter ratio = 13.7.

In the title compound, $\text{C}_{18}\text{H}_{15}\text{ClFN}_3\text{O}_4$, the dihedral angle between the substituted pyridine ring and the oxadiazoline ring is $9.73(19)^\circ$ and the acyl group is coplanar with the oxadiazoline ring [$\text{O}-\text{C}-\text{N}-\text{C}$ torsion angle = $-2.1(3)^\circ$]. Furthermore, the substituted benzene ring is almost orthogonal with the oxadiazoline ring, the dihedral angle between them being $87.56(18)^\circ$.

Related literature

For background to 1,3,4-oxadiazoline derivatives and related structures, see: Song *et al.* (2006a,b); Pan *et al.* (2007). For the pharmacological properties of 2,5-disubstituted 1,3,4-oxadiazolines, see: Chimirri *et al.* (1994, 1996); Dogan *et al.* (1998).



Experimental

Crystal data

$\text{C}_{18}\text{H}_{15}\text{ClFN}_3\text{O}_4$	$V = 1831.8(6)\text{ \AA}^3$
$M_r = 391.78$	$Z = 4$
Monoclinic, $P2_1/n$	Mo $K\alpha$ radiation
$a = 10.120(2)\text{ \AA}$	$\mu = 0.25\text{ mm}^{-1}$
$b = 13.900(3)\text{ \AA}$	$T = 293\text{ K}$
$c = 13.320(3)\text{ \AA}$	$0.12 \times 0.10 \times 0.08\text{ mm}$
$\beta = 102.14(3)^\circ$	

Data collection

Bruker APEXII CCD area-detector diffractometer	9882 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 2000)	3403 independent reflections
$T_{\min} = 0.971$, $T_{\max} = 0.981$	2394 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.026$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.043$	248 parameters
$wR(F^2) = 0.121$	H-atom parameters not refined
$S = 1.00$	$\Delta\rho_{\max} = 0.22\text{ e \AA}^{-3}$
3403 reflections	$\Delta\rho_{\min} = -0.22\text{ e \AA}^{-3}$

Data collection: *APEX2* (Bruker, 2000); cell refinement: *SAINT* (Bruker, 2000); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

This work was supported by the Key Discipline of Applied Chemistry, Zhejiang Province and the State Key Laboratory Breeding Base of Green Chemistry–Synthesis Technology, Zhejiang University of Technology (People's Republic of China).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: TK2411).

References

- Bruker (2000). *APEX2, SAINT* and *SADABS*. Bruker AXS inc., Madison, Wisconsin, USA.
- Chimirri, A., Grasso, S., Monforte, A. M., Monforte, P., Zappala, M. & Carotti, A. (1994). *Farmaco*, **49**, 509–511.
- Chimirri, A., Grasso, S., Montforte, A. M., Rao, A. & Zappala, M. (1996). *Farmaco*, **51**, 125–129.
- Dogan, H. N., Duran, A., Rollas, S., Sener, G., Armutak, Y. & Keyer-Uysal, M. (1998). *Med. Sci. Res.*, **26**, 755–758.
- Pan, L.-F., Lu, Y.-Q., Qin, Q., Qi, C.-Z. & Song, Q.-B. (2007). *Acta Cryst. E63*, o3988.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Song, Q.-B., Zhang, J., Dong, Y. & Tiekkink, E. R. T. (2006a). *Acta Cryst. E62*, o4388–o4390.
- Song, Q.-B., Zhang, J. & Tiekkink, E. R. T. (2006b). *Acta Cryst. E62*, o4115–o4117.

supplementary materials

Acta Cryst. (2009). E65, o1181 [doi:10.1107/S1600536809015323]

2-[3-Acetyl-5-(2-chloro-3-pyridyl)-2-methyl-2,3-dihydro-1,3,4-oxadiazol-2-yl]-4-fluorophenyl acetate

Q. Qin, L. J. Xu, L. F. Pan, S. Q. Chen and Q. B. Song

Comment

In continuation of our study of 1,3,4-oxadiazoline derivatives (Song *et al.*, 2006a,b; Pan *et al.*, 2007), which possess a wide range of pharmaceutical activities (Chimirri *et al.*, 1994, 1996; Dogan *et al.*, 1998), a series of new 1,3,4-oxadiazoline derivatives have been prepared. We present herein the crystal structure of the title compound, (I).

In (I), Fig. 1, the molecule is twisted about the C8—C9 bond. Within the five-membered oxadiazoline ring, there is a formal C13==N1 double bond (1.282 (3) Å). The bond distance of C13—O4 (1.360 (2) Å) is considerably shorter than that of C9—O4 bond (1.452 (2) Å), suggesting some delocalization of π -electron density over the O4—C13—N1 chromophore.

Experimental

A solution of 2-chloro-*N'*-(1-(5-fluoro-2-hydroxyphenyl)ethylidene) nicotinohydrazide (0.5 g, 1.62 mmol) in acetic anhydride (10 ml) was refluxed until the reaction was finished. The acetic anhydride was distilled under vacuum. The residue was recrystallized from ethanol (10 ml). Colorless crystals (0.46 g) were obtained by slow evaporation of an ethanol solution of (I) after 2 days at room temperature.

Refinement

All H atoms were placed in calculated positions and allowed to ride on their parent atoms, with C—H = 0.93–0.96 Å and refined in the riding model approximation with $U_{\text{iso}}(\text{H}) = 1.2\text{--}1.5U_{\text{eq}}(\text{C})$.

Figures




Fig. 1. View of (I) showing the atom-labelling scheme. Displacement ellipsoids are drawn at the 40% probability level. H atoms are represented by circles of arbitrary size.

supplementary materials

2-[3-Acetyl-5-(2-chloro-3-pyridyl)-2-methyl-2,3-dihydro-1,3,4-oxadiazol-2-yl]-4-fluorophenyl acetate

Crystal data

C ₁₈ H ₁₅ ClFN ₃ O ₄	$F_{000} = 808$
$M_r = 391.78$	$D_x = 1.421 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/n$	Mo $K\alpha$ radiation
	$\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2yn	Cell parameters from 3403 reflections
$a = 10.120 (2) \text{ \AA}$	$\theta = 2.1\text{--}25.5^\circ$
$b = 13.900 (3) \text{ \AA}$	$\mu = 0.25 \text{ mm}^{-1}$
$c = 13.320 (3) \text{ \AA}$	$T = 293 \text{ K}$
$\beta = 102.14 (3)^\circ$	Block, colorless
$V = 1831.8 (6) \text{ \AA}^3$	$0.12 \times 0.10 \times 0.08 \text{ mm}$
$Z = 4$	

Data collection

Bruker APEXII CCD area-detector diffractometer	3403 independent reflections
Radiation source: fine-focus sealed tube	2394 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.026$
$T = 293 \text{ K}$	$\theta_{\text{max}} = 25.5^\circ$
φ and ω scans	$\theta_{\text{min}} = 2.1^\circ$
Absorption correction: multi-scan (SADABS; Bruker, 2000)	$h = -12 \rightarrow 7$
$T_{\text{min}} = 0.971$, $T_{\text{max}} = 0.981$	$k = -16 \rightarrow 16$
9882 measured reflections	$l = -16 \rightarrow 16$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.043$	H-atom parameters not refined
$wR(F^2) = 0.121$	$w = 1/[\sigma^2(F_o^2) + (0.06P)^2 + 0.3383P]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.00$	$(\Delta/\sigma)_{\text{max}} < 0.001$
3403 reflections	$\Delta\rho_{\text{max}} = 0.22 \text{ e \AA}^{-3}$
248 parameters	$\Delta\rho_{\text{min}} = -0.22 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.0490 (3)	0.1369 (2)	0.5790 (2)	0.0842 (8)
H1A	-0.0049	0.1943	0.5694	0.126*
H1B	0.0010	0.0859	0.5383	0.126*
H1C	0.1328	0.1485	0.5580	0.126*
C2	0.0772 (2)	0.10902 (17)	0.6890 (2)	0.0686 (6)
C3	0.1358 (2)	-0.02657 (15)	0.79837 (16)	0.0566 (5)
C4	0.0293 (3)	-0.04387 (16)	0.8454 (2)	0.0702 (7)
H4	-0.0576	-0.0247	0.8140	0.084*
C5	0.0520 (3)	-0.08956 (18)	0.9390 (2)	0.0778 (7)
H5	-0.0187	-0.1021	0.9718	0.093*
C6	0.1815 (3)	-0.11600 (17)	0.98264 (17)	0.0729 (7)
C7	0.2892 (3)	-0.09887 (15)	0.93740 (16)	0.0627 (6)
H7	0.3758	-0.1177	0.9699	0.075*
C8	0.2675 (2)	-0.05317 (14)	0.84270 (15)	0.0523 (5)
C9	0.3835 (2)	-0.03837 (14)	0.78801 (14)	0.0517 (5)
C10	0.5212 (2)	-0.07353 (16)	0.84381 (18)	0.0655 (6)
H10A	0.5183	-0.1419	0.8534	0.098*
H10B	0.5454	-0.0425	0.9095	0.098*
H10C	0.5872	-0.0585	0.8039	0.098*
C11	0.4183 (3)	0.23549 (16)	0.76258 (19)	0.0731 (7)
H11A	0.3303	0.2645	0.7478	0.110*
H11B	0.4494	0.2268	0.6999	0.110*
H11C	0.4801	0.2765	0.8080	0.110*
C12	0.4110 (2)	0.14036 (15)	0.81247 (16)	0.0551 (5)
C13	0.35464 (19)	-0.02406 (16)	0.61505 (14)	0.0512 (5)
C14	0.3289 (2)	-0.06543 (17)	0.51150 (15)	0.0561 (5)
C15	0.3246 (2)	-0.16524 (19)	0.50291 (18)	0.0692 (6)
H15	0.3420	-0.2032	0.5617	0.083*
C16	0.2944 (2)	-0.2080 (2)	0.4075 (2)	0.0806 (8)
H16	0.2913	-0.2746	0.4009	0.097*
C17	0.2691 (3)	-0.1500 (3)	0.3232 (2)	0.0896 (9)
H17	0.2472	-0.1791	0.2590	0.107*
C18	0.3038 (2)	-0.0137 (2)	0.41987 (16)	0.0655 (6)

supplementary materials

C1	0.30866 (7)	0.11042 (5)	0.41760 (5)	0.0845 (3)
F1	0.2064 (2)	-0.16130 (13)	1.07494 (10)	0.1092 (6)
N1	0.37927 (17)	0.06330 (13)	0.64394 (12)	0.0543 (4)
N2	0.39054 (17)	0.06139 (12)	0.75037 (12)	0.0525 (4)
N3	0.2739 (2)	-0.0537 (2)	0.32705 (14)	0.0793 (6)
O1	0.10784 (15)	0.01308 (10)	0.69914 (11)	0.0612 (4)
O2	0.42063 (17)	0.13026 (11)	0.90474 (11)	0.0697 (5)
O3	0.0761 (2)	0.15939 (13)	0.76138 (16)	0.0960 (6)
O4	0.34983 (15)	-0.08897 (10)	0.69061 (10)	0.0569 (4)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0707 (17)	0.0814 (18)	0.097 (2)	0.0185 (13)	0.0106 (14)	0.0250 (15)
C2	0.0516 (13)	0.0587 (15)	0.0921 (19)	0.0046 (11)	0.0072 (12)	-0.0006 (14)
C3	0.0683 (15)	0.0453 (11)	0.0562 (12)	-0.0010 (10)	0.0132 (11)	-0.0100 (9)
C4	0.0750 (16)	0.0595 (15)	0.0802 (17)	-0.0053 (12)	0.0258 (13)	-0.0181 (12)
C5	0.098 (2)	0.0677 (16)	0.0785 (18)	-0.0171 (15)	0.0424 (16)	-0.0228 (14)
C6	0.116 (2)	0.0610 (15)	0.0452 (13)	-0.0158 (14)	0.0260 (14)	-0.0094 (10)
C7	0.0875 (17)	0.0523 (13)	0.0470 (12)	-0.0097 (11)	0.0112 (12)	-0.0046 (10)
C8	0.0686 (14)	0.0409 (11)	0.0462 (11)	-0.0038 (9)	0.0097 (10)	-0.0070 (8)
C9	0.0636 (13)	0.0448 (11)	0.0434 (11)	0.0005 (9)	0.0036 (9)	0.0021 (8)
C10	0.0662 (14)	0.0631 (14)	0.0630 (14)	0.0084 (11)	0.0038 (11)	0.0152 (11)
C11	0.0900 (18)	0.0493 (13)	0.0749 (16)	-0.0044 (12)	0.0058 (13)	0.0103 (11)
C12	0.0573 (13)	0.0493 (12)	0.0531 (13)	0.0000 (9)	-0.0012 (10)	0.0032 (9)
C13	0.0449 (11)	0.0609 (13)	0.0457 (11)	0.0075 (10)	0.0048 (9)	0.0037 (10)
C14	0.0425 (11)	0.0771 (15)	0.0474 (12)	0.0085 (10)	0.0065 (9)	-0.0007 (10)
C15	0.0611 (14)	0.0862 (18)	0.0574 (14)	0.0141 (12)	0.0058 (11)	-0.0118 (12)
C16	0.0746 (17)	0.095 (2)	0.0675 (17)	0.0166 (14)	0.0046 (13)	-0.0188 (14)
C17	0.0727 (18)	0.132 (3)	0.0597 (17)	0.0181 (18)	0.0046 (13)	-0.0288 (17)
C18	0.0463 (12)	0.0983 (18)	0.0509 (13)	0.0108 (11)	0.0077 (10)	0.0046 (12)
C11	0.0910 (5)	0.0999 (5)	0.0607 (4)	0.0095 (4)	0.0117 (3)	0.0222 (3)
F1	0.1671 (18)	0.1102 (13)	0.0567 (9)	-0.0254 (11)	0.0381 (10)	0.0077 (8)
N1	0.0541 (10)	0.0628 (12)	0.0431 (9)	0.0031 (8)	0.0033 (8)	0.0088 (8)
N2	0.0628 (11)	0.0474 (10)	0.0429 (9)	0.0000 (8)	0.0011 (8)	0.0065 (7)
N3	0.0658 (13)	0.123 (2)	0.0473 (12)	0.0129 (13)	0.0071 (9)	-0.0021 (11)
O1	0.0643 (9)	0.0546 (9)	0.0614 (9)	0.0060 (7)	0.0060 (7)	-0.0009 (7)
O2	0.0934 (12)	0.0583 (9)	0.0497 (9)	-0.0047 (8)	-0.0023 (8)	0.0010 (7)
O3	0.1075 (15)	0.0635 (11)	0.1116 (16)	0.0128 (10)	0.0107 (12)	-0.0142 (11)
O4	0.0735 (10)	0.0501 (8)	0.0461 (8)	0.0030 (7)	0.0102 (7)	-0.0016 (6)

Geometric parameters (\AA , $^\circ$)

C1—C2	1.484 (4)	C10—H10B	0.9600
C1—H1A	0.9600	C10—H10C	0.9600
C1—H1B	0.9600	C11—C12	1.489 (3)
C1—H1C	0.9600	C11—H11A	0.9600
C2—O3	1.194 (3)	C11—H11B	0.9600
C2—O1	1.369 (3)	C11—H11C	0.9600

C3—C4	1.377 (3)	C12—O2	1.220 (2)
C3—C8	1.390 (3)	C12—N2	1.364 (3)
C3—O1	1.405 (2)	C13—N1	1.282 (3)
C4—C5	1.375 (3)	C13—O4	1.360 (2)
C4—H4	0.9300	C13—C14	1.466 (3)
C5—C6	1.367 (4)	C14—C15	1.392 (3)
C5—H5	0.9300	C14—C18	1.393 (3)
C6—F1	1.357 (3)	C15—C16	1.377 (3)
C6—C7	1.373 (3)	C15—H15	0.9300
C7—C8	1.388 (3)	C16—C17	1.362 (4)
C7—H7	0.9300	C16—H16	0.9300
C8—C9	1.520 (3)	C17—N3	1.340 (4)
C9—O4	1.452 (2)	C17—H17	0.9300
C9—N2	1.481 (2)	C18—N3	1.331 (3)
C9—C10	1.516 (3)	C18—Cl1	1.727 (3)
C10—H10A	0.9600	N1—N2	1.398 (2)
C2—C1—H1A	109.5	H10A—C10—H10C	109.5
C2—C1—H1B	109.5	H10B—C10—H10C	109.5
H1A—C1—H1B	109.5	C12—C11—H11A	109.5
C2—C1—H1C	109.5	C12—C11—H11B	109.5
H1A—C1—H1C	109.5	H11A—C11—H11B	109.5
H1B—C1—H1C	109.5	C12—C11—H11C	109.5
O3—C2—O1	122.1 (2)	H11A—C11—H11C	109.5
O3—C2—C1	127.7 (2)	H11B—C11—H11C	109.5
O1—C2—C1	110.2 (2)	O2—C12—N2	119.22 (19)
C4—C3—C8	122.2 (2)	O2—C12—C11	123.4 (2)
C4—C3—O1	118.3 (2)	N2—C12—C11	117.36 (19)
C8—C3—O1	119.31 (19)	N1—C13—O4	116.17 (17)
C5—C4—C3	119.7 (3)	N1—C13—C14	129.68 (19)
C5—C4—H4	120.1	O4—C13—C14	114.15 (18)
C3—C4—H4	120.1	C15—C14—C18	116.4 (2)
C6—C5—C4	118.2 (2)	C15—C14—C13	117.68 (19)
C6—C5—H5	120.9	C18—C14—C13	125.8 (2)
C4—C5—H5	120.9	C16—C15—C14	120.2 (2)
F1—C6—C5	119.3 (3)	C16—C15—H15	119.9
F1—C6—C7	117.7 (3)	C14—C15—H15	119.9
C5—C6—C7	123.0 (2)	C17—C16—C15	118.2 (3)
C6—C7—C8	119.4 (2)	C17—C16—H16	120.9
C6—C7—H7	120.3	C15—C16—H16	120.9
C8—C7—H7	120.3	N3—C17—C16	124.1 (2)
C7—C8—C3	117.5 (2)	N3—C17—H17	118.0
C7—C8—C9	120.5 (2)	C16—C17—H17	118.0
C3—C8—C9	121.92 (18)	N3—C18—C14	124.3 (3)
O4—C9—N2	99.78 (14)	N3—C18—Cl1	113.72 (19)
O4—C9—C10	107.50 (17)	C14—C18—Cl1	122.02 (18)
N2—C9—C10	111.29 (17)	C13—N1—N2	104.80 (16)
O4—C9—C8	107.64 (16)	C12—N2—N1	124.70 (17)
N2—C9—C8	112.70 (16)	C12—N2—C9	124.06 (16)
C10—C9—C8	116.38 (17)	N1—N2—C9	111.19 (15)

supplementary materials

C9—C10—H10A	109.5	C18—N3—C17	116.8 (2)
C9—C10—H10B	109.5	C2—O1—C3	118.16 (18)
H10A—C10—H10B	109.5	C13—O4—C9	107.56 (15)
C9—C10—H10C	109.5		
C8—C3—C4—C5	-0.5 (3)	C15—C14—C18—Cl1	-178.79 (16)
O1—C3—C4—C5	174.88 (19)	C13—C14—C18—Cl1	3.5 (3)
C3—C4—C5—C6	0.3 (3)	O4—C13—N1—N2	-1.3 (2)
C4—C5—C6—F1	-179.9 (2)	C14—C13—N1—N2	177.94 (19)
C4—C5—C6—C7	0.1 (4)	O2—C12—N2—N1	-179.27 (18)
F1—C6—C7—C8	179.65 (18)	C11—C12—N2—N1	1.6 (3)
C5—C6—C7—C8	-0.4 (3)	O2—C12—N2—C9	-2.1 (3)
C6—C7—C8—C3	0.2 (3)	C11—C12—N2—C9	178.74 (19)
C6—C7—C8—C9	-176.51 (19)	C13—N1—N2—C12	-177.06 (19)
C4—C3—C8—C7	0.2 (3)	C13—N1—N2—C9	5.5 (2)
O1—C3—C8—C7	-175.12 (17)	O4—C9—N2—C12	175.46 (18)
C4—C3—C8—C9	176.90 (19)	C10—C9—N2—C12	-71.3 (3)
O1—C3—C8—C9	1.5 (3)	C8—C9—N2—C12	61.5 (2)
C7—C8—C9—O4	119.08 (18)	O4—C9—N2—N1	-7.0 (2)
C3—C8—C9—O4	-57.5 (2)	C10—C9—N2—N1	106.21 (19)
C7—C8—C9—N2	-131.88 (18)	C8—C9—N2—N1	-120.95 (17)
C3—C8—C9—N2	51.6 (2)	C14—C18—N3—C17	-0.5 (3)
C7—C8—C9—C10	-1.6 (3)	C11—C18—N3—C17	179.67 (18)
C3—C8—C9—C10	-178.13 (18)	C16—C17—N3—C18	-0.8 (4)
N1—C13—C14—C15	170.8 (2)	O3—C2—O1—C3	1.4 (3)
O4—C13—C14—C15	-10.0 (3)	C1—C2—O1—C3	-179.18 (19)
N1—C13—C14—C18	-11.5 (3)	C4—C3—O1—C2	75.7 (2)
O4—C13—C14—C18	167.70 (19)	C8—C3—O1—C2	-108.7 (2)
C18—C14—C15—C16	-1.0 (3)	N1—C13—O4—C9	-3.4 (2)
C13—C14—C15—C16	176.9 (2)	C14—C13—O4—C9	177.29 (16)
C14—C15—C16—C17	-0.1 (4)	N2—C9—O4—C13	5.97 (19)
C15—C16—C17—N3	1.1 (4)	C10—C9—O4—C13	-110.18 (18)
C15—C14—C18—N3	1.4 (3)	C8—C9—O4—C13	123.72 (16)
C13—C14—C18—N3	-176.3 (2)		

Fig. 1

