

## 6-Bromo-1-methyl-1*H*-2,1-benzothiazin-4(3*H*)-one 2,2-dioxide

Muhammad Shafiq,<sup>a</sup> M. Nawaz Tahir,<sup>b\*</sup> Islam Ullah Khan,<sup>a</sup> Muhammad Nadeem Arshad<sup>a</sup> and Muhammad Nadeem Asghar<sup>a</sup>

<sup>a</sup>Government College University, Department of Chemistry, Lahore, Pakistan, and

<sup>b</sup>University of Sargodha, Department of Physics, Sargodha, Pakistan

Correspondence e-mail: dmntahir\_uos@yahoo.com


Received 6 March 2009; accepted 28 April 2009

Key indicators: single-crystal X-ray study;  $T = 296\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$ ;  $R$  factor = 0.030;  $wR$  factor = 0.072; data-to-parameter ratio = 16.3.

In the crystal structure of the title compound,  $\text{C}_9\text{H}_8\text{BrNO}_3\text{S}$ , the thiazine ring is in the twisted form. In the crystal, pairs of intermolecular  $\text{C}-\text{H}\cdots\text{O}$  hydrogen bonds form inversion dimers with an  $R_2^2(8)$  ring motif. Weak intermolecular  $\text{C}-\text{H}\cdots\text{Br}$  and  $\text{C}-\text{H}\cdots\pi$  interactions are also present.

### Related literature

For the structures of benzothiazine derivatives, see: Arshad *et al.* (2008); Shafiq *et al.* (2008a,b); Tahir *et al.* (2008). For the related structure, 6-bromo-1-methyl-1*H*-benzo[*c*][1,2]thiazin-4(3*H*)-one 2,2-dioxide, see: Shafiq *et al.* (2009). For hydrogen-bond motifs, see: Bernstein *et al.* (1995). For puckering parameters, see: Cremer & Pople (1975). For the synthesis, see: Lombardino (1972).



### Experimental

#### Crystal data

$\text{C}_9\text{H}_8\text{BrNO}_3\text{S}$

$M_r = 290.13$

Monoclinic,  $P2_{1}/n$

$a = 5.4577(3)\text{ \AA}$

$b = 12.6400(8)\text{ \AA}$

$c = 15.1258(10)\text{ \AA}$

$\beta = 96.204(2)^\circ$

$V = 1037.35(11)\text{ \AA}^3$

$Z = 4$

Mo  $K\alpha$  radiation

$\mu = 4.15\text{ mm}^{-1}$

$T = 296\text{ K}$

$0.20 \times 0.17 \times 0.15\text{ mm}$

#### Data collection

Bruker Kappa APEXII CCD

diffractometer

Absorption correction: multi-scan  
(*SADABS*; Bruker, 2005)

$T_{\min} = 0.439$ ,  $T_{\max} = 0.540$

11077 measured reflections

2234 independent reflections

1709 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.032$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.030$

$wR(F^2) = 0.072$

$S = 1.04$

2234 reflections

137 parameters

H-atom parameters constrained

$\Delta\rho_{\text{max}} = 0.41\text{ e \AA}^{-3}$

$\Delta\rho_{\text{min}} = -0.35\text{ e \AA}^{-3}$

**Table 1**  
Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C3—H3 $\cdots$ O3 <sup>i</sup>	0.93	2.54	3.308 (4)	140
C8—H8A $\cdots$ O2 <sup>ii</sup>	0.97	2.54	3.470 (3)	162
C9—H9B $\cdots$ O3	0.96	2.41	2.824 (3)	106
C5—H5 $\cdots$ Br1 <sup>iii</sup>	0.93	2.94	3.871 (3)	175
C9—H9A $\cdots$ Br1 <sup>iv</sup>	0.96	3.01	3.871 (2)	150
C9—H9C $\cdots$ Cg1 <sup>v</sup>	0.96	2.83	3.449 (3)	123

Symmetry codes: (i)  $-x + \frac{3}{2}, y + \frac{1}{2}, -z + \frac{1}{2}$ ; (ii)  $-x + 1, -y, -z + 1$ ; (iii)  $-x, -y + 1, -z + 1$ ; (iv)  $-x + \frac{1}{2}, y - \frac{1}{2}, -z + \frac{1}{2}$ ; (v)  $x + 1, y, z$ . Cg1 is the centroid of the C1–C6 ring.

Data collection: *APEX2* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997) and *PLATON* (Spek, 2009); software used to prepare material for publication: *WinGX* (Farrugia, 1999) and *PLATON*.

MS gratefully acknowledges the Higher Education Commission, Islamabad, Pakistan, for providing a Scholarship under the Indigenous PhD Program (PIN 042–120567-PS2–276).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: FB2142).

### References

- Arshad, M. N., Tahir, M. N., Khan, I. U., Shafiq, M. & Siddiqui, W. A. (2008). *Acta Cryst. E64*, o2045.
- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Bruker (2005). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2007). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Cremer, D. & Pople, J. A. (1975). *J. Am. Chem. Soc.* **97**, 1354–1358.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Lombardino, J. G. (1972). *J. Heterocycl. Chem.* **9**, 315–317.
- Shafiq, M., Khan, I. U., Tahir, M. N. & Siddiqui, W. A. (2008a). *Acta Cryst. E64*, o558.
- Shafiq, M., Tahir, M. N., Khan, I. U., Ahmad, S. & Siddiqui, W. A. (2008b). *Acta Cryst. E64*, o1270.
- Shafiq, M., Tahir, M. N., Khan, I. U., Arshad, M. N. & Safdar, M. (2009). *Acta Cryst. E65*, o393.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D65*, 148–155.
- Tahir, M. N., Shafiq, M., Khan, I. U., Siddiqui, W. A. & Arshad, M. N. (2008). *Acta Cryst. E64*, o557.

## **supplementary materials**

Acta Cryst. (2009). E65, o1182 [doi:10.1107/S1600536809015980]

## 6-Bromo-1-methyl-1*H*-2,1-benzothiazin-4(3*H*)-one 2,2-dioxide

M. Shafiq, M. N. Tahir, I. U. Khan, M. N. Arshad and M. N. Asghar

### Comment

We have reported crystal structures of the synthesized derivatives of the benzothiazine molecule (Shafiq *et al.*, 2008a; Shafiq *et al.*, 2008b; Tahir *et al.*, 2008; Arshad *et al.*, 2008). Here we report the title compound (I), (Fig. 1), that belongs to this series of the structures.

(I) is closely related to the crystal structure of 6-bromo-1-methyl-1*H*-benzo[*c*][1,2]thiazin-4(3*H*)-one 2,2-dioxide, (II), (Shafiq *et al.*, 2009). (I) and (II) differ by the presence of the methyl and ethyl groups at the N-atom, respectively. The bromo-substituted benzene ring A (C1—C6) is planar with Br deviated by 0.064 (4) Å from the mean plane. The thiazine ring B (S1/N1/C1/C6—C8) is in the twisted form, with the maximum puckering amplitude  $Q_T = 0.577$  (2) Å (Cremer & Pople, 1975). The title molecules form dimers interconnected by a pair of the intermolecular H-bonds C8—H8A···O2<sup>i</sup> [symmetry code:  $i = -x + 1, -y, -z + 1$ ] with the  $R_2^2(8)$  ring motif (Bernstein *et al.*, 1995), (Tab. 1, Fig. 2). The dimers are linked to each other forming helices through the other intermolecular H-bonding C3—H3···O3<sup>ii</sup> [symmetry code:  $ii = -x + 3/2, y + 1/2, -z + 1/2$ ]. The molecules are also stabilized due to C—H···π-electron interaction with the benzene group and intermolecular C—H···Br interactions (Tab. 1).

### Experimental

The title compound was prepared in a three step scheme following the reported procedure (Lombardino, 1972). In the first step, methyl-2-amino-5-bromobenzoate (92 mg, 4 mmol) was put in dichloromethane (10 ml) and this mixture was introduced into a round bottom flask. A solution of methanesulfonyl chloride (550 mg, 4.8 mmol) in dichloromethane (10 ml) was slowly added (10–15 minutes) to this mixture. The mixture was stirred at 60–70 °C for 2–3 days keeping pH of the mixture alkaline by triethylamine. After the completion of the reaction, the solvent was evaporated under reduced pressure to get methyl-5-bromo-2-[(methylsulfonyl)amino] benzoate.

In the second step, methyl-5-bromo-2-[(methylsulfonyl)amino] benzoate (1.02 g, 3.3 mmol) was introduced into 5 ml of *N,N*-dimethylformamide (DMF). The mixture was added to a suspension of NaH (158.38 mg, 6.6 mmol) in DMF (10 ml). The mixture was stirred at room temperature for 14–16 h. After that, methyl-5-bromo-2-[methyl(methylsulfonyl)amino]benzoate was obtained.

In the third step methyl-5-bromo-2-[methyl(methylsulfonyl)amino]benzoate was cyclized. Therefore methyl-5-bromo-2-[methyl(methylsulfonyl)amino]benzoate (418.83 mg, 1.3 mmol) was introduced in DMF (5 ml) and added to the suspension of NaH (59.99 mg, 2.5 mmol) in DMF (10 ml). The mixture was stirred at room temperature for 3–4 h. Then the reaction mixture was poured into ice and clear solution was obtained. The pH of this solution was adjusted between 5–6. The precipitated crude product was recrystallized from ethanol. Yellow needle-shaped crystals of the title compound of suitable size for structure analysis were grown in this way.

# supplementary materials

---

## Refinement

Though all the hydrogens were discernible in the difference electron density map, the H-atoms were situated into idealized positions, with C-H = 0.93, 0.96 and 0.97 Å for aryl, methyl and methylene H, respectively, and constrained to ride on their parent atoms, with  $U_{\text{iso}}(\text{H}) = xU_{\text{eq}}(\text{C})$ , where  $x = 1.5$  for methyl and 1.2 for other carrier atoms.

## Figures




Fig. 1. The title compound, with the atom-numbering scheme. The displacement ellipsoids are drawn at the 50% probability level. The H-atoms are shown by small circles of arbitrary radius. The dotted lines show the intramolecular H-bonds.




Fig. 2. A section of the title structure showing the dimers bind by the hydrogen bonds.

## 6-Bromo-1-methyl-1*H*-2,1-benzothiazin-4(3*H*)-one 2,2-dioxide

### Crystal data

C <sub>9</sub> H <sub>8</sub> BrNO <sub>3</sub> S	$F_{000} = 576$
$M_r = 290.13$	$D_x = 1.858 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/n$	Mo $K\alpha$ radiation
Hall symbol: -P 2yn	$\lambda = 0.71073 \text{ \AA}$
$a = 5.4577 (3) \text{ \AA}$	Cell parameters from 2234 reflections
$b = 12.6400 (8) \text{ \AA}$	$\theta = 2.1\text{--}27.0^\circ$
$c = 15.1258 (10) \text{ \AA}$	$\mu = 4.15 \text{ mm}^{-1}$
$\beta = 96.204 (2)^\circ$	$T = 296 \text{ K}$
$V = 1037.35 (11) \text{ \AA}^3$	Prism, yellow
$Z = 4$	$0.20 \times 0.17 \times 0.15 \text{ mm}$

### Data collection

Bruker Kappa APEXII CCD diffractometer	2234 independent reflections
Radiation source: fine-focus sealed tube	1709 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.032$
Detector resolution: 7.40 pixels $\text{mm}^{-1}$	$\theta_{\text{max}} = 27.0^\circ$

$T = 296 \text{ K}$                                    $\theta_{\min} = 2.1^\circ$   
 $\omega$  scans     $h = -6 \rightarrow 6$   
 Absorption correction: multi-scan  
 (SADABS; Bruker, 2005)                           $k = -16 \rightarrow 16$   
 $T_{\min} = 0.439, T_{\max} = 0.540$                            $l = -18 \rightarrow 19$   
 11077 measured reflections

### Refinement

Refinement on  $F^2$     Secondary atom site location: difference Fourier map  
 Least-squares matrix: full                                  Hydrogen site location: difference Fourier map  
 $R[F^2 > 2\sigma(F^2)] = 0.030$                                   H-atom parameters constrained  
 $wR(F^2) = 0.072$      $w = 1/[\sigma^2(F_o^2) + (0.0309P)^2 + 0.4349P]$   
 $S = 1.04$     where  $P = (F_o^2 + 2F_c^2)/3$   
 2234 reflections     $(\Delta/\sigma)_{\max} = 0.001$   
 137 parameters     $\Delta\rho_{\max} = 0.41 \text{ e \AA}^{-3}$   
 31 constraints     $\Delta\rho_{\min} = -0.35 \text{ e \AA}^{-3}$   
 Extinction correction: none  
 Primary atom site location: structure-invariant direct  
 methods

### Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	0.06150 (6)	0.57510 (2)	0.36727 (2)	0.05871 (14)
S1	0.74473 (12)	0.10243 (5)	0.40740 (4)	0.03706 (17)
O1	0.3779 (4)	0.23972 (17)	0.56688 (14)	0.0606 (6)
O2	0.5116 (4)	0.05733 (15)	0.37692 (14)	0.0529 (5)
O3	0.9592 (4)	0.03989 (16)	0.40469 (14)	0.0555 (6)
N1	0.7895 (4)	0.21290 (17)	0.35343 (14)	0.0385 (5)
C1	0.6129 (4)	0.29366 (19)	0.35581 (16)	0.0326 (6)
C2	0.5631 (5)	0.3622 (2)	0.28382 (19)	0.0403 (6)
H2	0.6430	0.3525	0.2332	0.048*
C3	0.3980 (5)	0.4437 (2)	0.2866 (2)	0.0430 (7)
H3	0.3650	0.4882	0.2378	0.052*
C4	0.2814 (5)	0.4595 (2)	0.36193 (19)	0.0409 (6)

## supplementary materials

---

C5	0.3200 (5)	0.3919 (2)	0.43312 (19)	0.0416 (6)
H5	0.2368	0.4024	0.4829	0.050*
C6	0.4846 (4)	0.3073 (2)	0.43087 (17)	0.0362 (6)
C7	0.5129 (5)	0.2357 (2)	0.50864 (18)	0.0407 (6)
C8	0.7204 (5)	0.1553 (2)	0.51326 (17)	0.0419 (6)
H8A	0.6884	0.0989	0.5540	0.050*
H8B	0.8745	0.1891	0.5355	0.050*
C9	0.9601 (4)	0.2116 (2)	0.28524 (19)	0.0415 (6)
H9A	0.8798	0.1812	0.2316	0.062*
H9B	1.1024	0.1701	0.3057	0.062*
H9C	1.0102	0.2826	0.2737	0.062*

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Br1	0.0680 (2)	0.0468 (2)	0.0609 (2)	0.02306 (14)	0.00505 (16)	0.00177 (15)
S1	0.0439 (3)	0.0358 (4)	0.0327 (4)	0.0072 (3)	0.0098 (3)	0.0042 (3)
O1	0.0815 (14)	0.0670 (14)	0.0380 (12)	0.0261 (11)	0.0282 (11)	0.0127 (11)
O2	0.0603 (12)	0.0441 (12)	0.0541 (14)	-0.0125 (9)	0.0050 (10)	0.0006 (10)
O3	0.0652 (12)	0.0557 (13)	0.0489 (13)	0.0284 (10)	0.0214 (10)	0.0133 (10)
N1	0.0420 (11)	0.0415 (13)	0.0350 (13)	0.0070 (9)	0.0179 (10)	0.0092 (10)
C1	0.0345 (12)	0.0319 (14)	0.0317 (15)	-0.0013 (10)	0.0052 (10)	0.0010 (11)
C2	0.0446 (14)	0.0398 (15)	0.0382 (17)	0.0001 (11)	0.0127 (12)	0.0070 (12)
C3	0.0523 (15)	0.0346 (15)	0.0423 (17)	-0.0010 (12)	0.0061 (13)	0.0102 (12)
C4	0.0426 (13)	0.0343 (14)	0.0453 (18)	0.0060 (11)	0.0026 (12)	-0.0001 (13)
C5	0.0483 (15)	0.0431 (15)	0.0347 (16)	0.0084 (12)	0.0104 (12)	-0.0017 (13)
C6	0.0415 (13)	0.0386 (15)	0.0288 (15)	0.0030 (11)	0.0051 (11)	0.0017 (11)
C7	0.0502 (14)	0.0431 (16)	0.0296 (15)	0.0072 (12)	0.0080 (12)	0.0006 (12)
C8	0.0523 (15)	0.0464 (17)	0.0274 (15)	0.0104 (12)	0.0058 (12)	0.0057 (12)
C9	0.0403 (13)	0.0444 (16)	0.0425 (17)	-0.0026 (11)	0.0171 (12)	-0.0018 (13)

### Geometric parameters ( $\text{\AA}$ , $^\circ$ )

Br1—C4	1.898 (3)	C3—C4	1.379 (4)
S1—O3	1.4169 (19)	C3—H3	0.9300
S1—O2	1.424 (2)	C4—C5	1.372 (4)
S1—N1	1.649 (2)	C5—C6	1.400 (3)
S1—C8	1.753 (3)	C5—H5	0.9300
O1—C7	1.209 (3)	C6—C7	1.479 (4)
N1—C1	1.407 (3)	C7—C8	1.518 (3)
N1—C9	1.463 (3)	C8—H8A	0.9700
C1—C2	1.395 (3)	C8—H8B	0.9700
C1—C6	1.407 (3)	C9—H9A	0.9600
C2—C3	1.373 (4)	C9—H9B	0.9600
C2—H2	0.9300	C9—H9C	0.9600
O3—S1—O2	118.63 (13)	C4—C5—C6	120.1 (2)
O3—S1—N1	106.93 (11)	C4—C5—H5	120.0
O2—S1—N1	110.70 (12)	C6—C5—H5	120.0

O3—S1—C8	112.54 (13)	C5—C6—C1	119.3 (2)
O2—S1—C8	107.17 (13)	C5—C6—C7	117.4 (2)
N1—S1—C8	99.15 (12)	C1—C6—C7	123.2 (2)
C1—N1—C9	121.1 (2)	O1—C7—C6	122.3 (2)
C1—N1—S1	117.60 (16)	O1—C7—C8	120.3 (2)
C9—N1—S1	118.62 (17)	C6—C7—C8	117.4 (2)
C2—C1—N1	120.5 (2)	C7—C8—S1	110.06 (18)
C2—C1—C6	118.8 (2)	C7—C8—H8A	109.6
N1—C1—C6	120.8 (2)	S1—C8—H8A	109.6
C3—C2—C1	121.1 (2)	C7—C8—H8B	109.6
C3—C2—H2	119.5	S1—C8—H8B	109.6
C1—C2—H2	119.5	H8A—C8—H8B	108.2
C2—C3—C4	119.8 (3)	N1—C9—H9A	109.5
C2—C3—H3	120.1	N1—C9—H9B	109.5
C4—C3—H3	120.1	H9A—C9—H9B	109.5
C5—C4—C3	120.9 (2)	N1—C9—H9C	109.5
C5—C4—Br1	119.3 (2)	H9A—C9—H9C	109.5
C3—C4—Br1	119.8 (2)	H9B—C9—H9C	109.5
O3—S1—N1—C1	−172.74 (19)	Br1—C4—C5—C6	−179.0 (2)
O2—S1—N1—C1	56.7 (2)	C4—C5—C6—C1	1.2 (4)
C8—S1—N1—C1	−55.7 (2)	C4—C5—C6—C7	−178.3 (2)
O3—S1—N1—C9	25.6 (2)	C2—C1—C6—C5	−3.1 (4)
O2—S1—N1—C9	−105.0 (2)	N1—C1—C6—C5	176.5 (2)
C8—S1—N1—C9	142.7 (2)	C2—C1—C6—C7	176.5 (2)
C9—N1—C1—C2	12.8 (4)	N1—C1—C6—C7	−3.9 (4)
S1—N1—C1—C2	−148.4 (2)	C5—C6—C7—O1	9.9 (4)
C9—N1—C1—C6	−166.8 (2)	C1—C6—C7—O1	−169.7 (3)
S1—N1—C1—C6	32.0 (3)	C5—C6—C7—C8	−170.1 (2)
N1—C1—C2—C3	−177.5 (2)	C1—C6—C7—C8	10.3 (4)
C6—C1—C2—C3	2.0 (4)	O1—C7—C8—S1	139.9 (2)
C1—C2—C3—C4	0.9 (4)	C6—C7—C8—S1	−40.0 (3)
C2—C3—C4—C5	−2.8 (4)	O3—S1—C8—C7	170.50 (19)
C2—C3—C4—Br1	177.9 (2)	O2—S1—C8—C7	−57.3 (2)
C3—C4—C5—C6	1.7 (4)	N1—S1—C8—C7	57.8 (2)

*Hydrogen-bond geometry (Å, °)*

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
C3—H3···O3 <sup>i</sup>	0.93	2.54	3.308 (4)	140
C8—H8A···O2 <sup>ii</sup>	0.97	2.54	3.470 (3)	162
C9—H9B···O3	0.96	2.41	2.824 (3)	106
C5—H5···Br1 <sup>iii</sup>	0.93	2.94	3.871 (3)	175
C9—H9A···Br1 <sup>iv</sup>	0.96	3.01	3.871 (2)	150
C9—H9C···Cg1 <sup>v</sup>	0.96	2.83	3.449 (3)	123

Symmetry codes: (i)  $-x+3/2, y+1/2, -z+1/2$ ; (ii)  $-x+1, -y, -z+1$ ; (iii)  $-x, -y+1, -z+1$ ; (iv)  $-x+1/2, y-1/2, -z+1/2$ ; (v)  $x+1, y, z$ .

## supplementary materials

---

Fig. 1




Fig. 2

