

Diazidobis[2,4-diamino-6-(2-pyridyl)-1,3,5-triazine- $\kappa^2 N^1, N^6$]zinc(II)

Qi-Hua Zhao,* Ai-Ling Fan, Li-Nan Li and Ming-Jing Xie

School of Chemical Science and Technology, Key Laboratory of Medicinal Chemistry for Natural Resources, Ministry of Education, Yunnan University, Kunming 650091, People's Republic of China

Correspondence e-mail: qhzao@ynu.edu.cn


Received 12 April 2009; accepted 29 April 2009

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.005\text{ \AA}$; R factor = 0.045; wR factor = 0.111; data-to-parameter ratio = 16.2.

In the title mononuclear complex, $[\text{Zn}(\text{N}_3)_2(\text{C}_8\text{H}_8\text{N}_6)_2]$, the Zn^{II} atom, lying on a twofold rotation axis, is six-coordinated in a distorted octahedral environment by four N atoms from two 2,4-diamino-6-(2-pyridyl)-1,3,5-triazine ligands and two N atoms from two end-on-coordinated azide ions. $\text{N}-\text{H}\cdots\text{N}$ hydrogen bonds between the ligand and azide ion link the complex molecules into a three-dimensional network.

Related literature

For general background to organic–inorganic hybrid complexes with azide ligands, see: Carranza *et al.* (2008); Gadad *et al.* (2000, 2004); Sun & Du (2005).



Experimental

Crystal data

$[\text{Zn}(\text{N}_3)_2(\text{C}_8\text{H}_8\text{N}_6)_2]$
 $M_r = 525.86$
Monoclinic, $C2/c$
 $a = 18.288(9)\text{ \AA}$
 $b = 14.231(7)\text{ \AA}$
 $c = 9.144(4)\text{ \AA}$
 $\beta = 115.382(5)^\circ$

$V = 2150.2(18)\text{ \AA}^3$
 $Z = 4$
Mo $K\alpha$ radiation
 $\mu = 1.19\text{ mm}^{-1}$
 $T = 293\text{ K}$
 $0.20 \times 0.18 \times 0.08\text{ mm}$

Data collection

Bruker APEXII CCD diffractometer
Absorption correction: multi-scan (*SADABS*; Sheldrick, 1996)
 $T_{\min} = 0.588$, $T_{\max} = 0.841$
(expected range = 0.636–0.909)

9145 measured reflections
2569 independent reflections
1766 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.056$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.045$
 $wR(F^2) = 0.111$
 $S = 1.00$
2569 reflections
159 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.51\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.41\text{ e \AA}^{-3}$

Table 1
Selected bond lengths (\AA).

Zn1–N7	2.153 (3)	Zn1–N1	2.202 (2)
Zn1–N4	2.166 (2)		

Table 2
Hydrogen-bond geometry (\AA , $^\circ$).

D–H \cdots A	D–H	H \cdots A	D \cdots A	D–H \cdots A
N5–H5A \cdots N3 ⁱ	0.86	2.19	3.042 (4)	175
N5–H5B \cdots N7 ⁱⁱ	0.86	2.31	3.060 (4)	147
N6–H6A \cdots N9 ⁱⁱⁱ	0.86	2.34	3.025 (4)	137
N6–H6B \cdots N7 ^{iv}	0.86	2.10	2.939 (4)	164

Symmetry codes: (i) $-x + \frac{3}{2}, -y + \frac{1}{2}, -z + 3$; (ii) $-x + \frac{3}{2}, -y + \frac{1}{2}, -z + 2$; (iii) $x, -y + 1, z + \frac{1}{2}$; (iv) $-x + 1, y, -z + \frac{3}{2}$.

Data collection: *APEX2* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

We acknowledge the National Natural Science Foundation of China (grant No. 20761005) for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HY2194).

References

- Bruker (2007). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Carranza, J., Julve, M. & Sletten, J. (2008). *Inorg. Chim. Acta*, **361**, 2499–2507.
- Gadad, A. K., Mahajanshetti, C. S., Nimbalkar, S. & Raichurkar, A. (2000). *Eur. J. Med. Chem.* **35**, 853–857.
- Gadad, A. K., Noolvi, M. N. & Karpoormath, R. V. (2004). *Bioorg. Med. Chem.* **12**, 5651–5659.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Sun, B.-W. & Du, L. (2005). *Acta Cryst. E* **61**, m1099–m1100.

supplementary materials

Acta Cryst. (2009). E65, m622 [doi:10.1107/S1600536809016055]

Diazidobis[2,4-diamino-6-(2-pyridyl)-1,3,5-triazine- κ^2N^1,N^6]zinc(II)

Q.-H. Zhao, A.-L. Fan, L.-N. Li and M.-J. Xie

Comment

At present, the design and synthesis of organic–inorganic hybrid compounds have drawn considerable attention. It was reported that 1*H*-1,3,5-triazole derivatives are a type of heterocyclic compounds with widely biological activity and outstanding capability. Meanwhile, a large number of complexes with azide ligands have been structurally and magnetically characterized (Carranza *et al.*, 2008; Gadad *et al.*, 2000, 2004; Sun & Du, 2005). Herein we report the synthesis and crystal structure of the title compound.

As shown in Fig. 1, the central Zn^{II} atom is coordinated by six N atoms and assumes a distorted octahedral geometry. The Zn—N(amide) bond [2.166 (2) Å] and the Zn—N(azide) bond [2.153 (3) Å] are significantly shorter than the Zn—N(pyridine) bond [2.202 (2) Å] (Table 1). The pyridyl and triazine rings of the ligand are essentially planar. The C3 atom shows an evident deviation of 0.1264 (3) Å from the mean plane through all atoms of the ligand. The two ligands coordinated to the Zn atom form a dihedral angle of 74.80 (7)°. The molecules are connected into a three-dimensional network through N—H···N hydrogen bonds (Table 2 and Fig. 2).

Experimental

All chemicals used (reagent grade) were commercially available. The compound was synthesized by heating a mixture of Zn(CH₃COO)₂ (30 mg, 0.15 mmol), 2,4-diamino-6-pyridyl-1,3,5-triazine (17.6 mg, 0.1 mmol), CH₃OH (5 ml) and H₂O (15 ml) at 338 K in stirring condition. After a few minutes, an aqueous solution (2 ml) of sodium azide (6.5 mg, 0.1 mmol) was added to it and stirred for 30 min. Colourless single crystals suitable for X-ray analysis were obtained by slow evaporation at room temperature for three weeks.

Refinement

H atoms were positioned geometrically and treated as riding on their parent atoms, with N—H = 0.86 and C—H = 0.93 Å and with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C}, \text{N})$.

supplementary materials

Figures




Fig. 1. The molecular structure of the title compound. Displacement ellipsoids are drawn at the 30% probability level. [Symmetry code: (i) -x+1, y, -z+3/2.]




Fig. 2. The crystal packing diagram of the title compound, showing hydrogen bonds (dashed lines).

Diazidobis[2,4-diamino-6-(2-pyridyl)-1,3,5-triazine- $\kappa^2\text{N}^1,\text{N}^6$]zinc(II)

Crystal data

[Zn(N ₃) ₂ (C ₈ H ₈ N ₆) ₂]	$F_{000} = 1072$
$M_r = 525.86$	$D_x = 1.624 \text{ Mg m}^{-3}$
Monoclinic, C2/c	Mo $K\alpha$ radiation
Hall symbol: -C 2yc	$\lambda = 0.71073 \text{ \AA}$
$a = 18.288 (9) \text{ \AA}$	Cell parameters from 2266 reflections
$b = 14.231 (7) \text{ \AA}$	$\theta = 1.9\text{--}28.6^\circ$
$c = 9.144 (4) \text{ \AA}$	$\mu = 1.19 \text{ mm}^{-1}$
$\beta = 115.382 (5)^\circ$	$T = 293 \text{ K}$
$V = 2150.2 (18) \text{ \AA}^3$	Block, colourless
$Z = 4$	$0.20 \times 0.18 \times 0.08 \text{ mm}$

Data collection

Bruker APEXII CCD diffractometer	2569 independent reflections
Radiation source: fine-focus sealed tube	1766 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.056$
$T = 293 \text{ K}$	$\theta_{\text{max}} = 28.6^\circ$
φ and ω scans	$\theta_{\text{min}} = 1.9^\circ$
Absorption correction: multi-scan	$h = -24\text{--}24$

(SADABS; Sheldrick, 1996)

$T_{\min} = 0.588$, $T_{\max} = 0.841$

9145 measured reflections

$k = -18 \rightarrow 19$

$l = -12 \rightarrow 12$

Refinement

Refinement on F^2

Secondary atom site location: difference Fourier map

Least-squares matrix: full

Hydrogen site location: inferred from neighbouring sites

$R[F^2 > 2\sigma(F^2)] = 0.045$

H-atom parameters constrained

$wR(F^2) = 0.111$

$$w = 1/[\sigma^2(F_o^2) + (0.0537P)^2 + 0.3944P]$$

where $P = (F_o^2 + 2F_c^2)/3$

$S = 1.00$

$(\Delta/\sigma)_{\max} < 0.001$

2569 reflections

$\Delta\rho_{\max} = 0.51 \text{ e \AA}^{-3}$

159 parameters

$\Delta\rho_{\min} = -0.41 \text{ e \AA}^{-3}$

Primary atom site location: structure-invariant direct methods

Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Zn1	0.5000	0.26762 (3)	0.7500	0.03110 (17)
N1	0.57688 (14)	0.16446 (17)	0.7022 (3)	0.0345 (6)
N2	0.72454 (13)	0.16338 (16)	1.1155 (3)	0.0317 (6)
N3	0.67379 (15)	0.25546 (16)	1.2722 (3)	0.0335 (6)
N4	0.59865 (13)	0.24286 (16)	0.9857 (3)	0.0295 (6)
N5	0.79367 (15)	0.17410 (19)	1.3913 (3)	0.0466 (7)
H5A	0.7994	0.1940	1.4845	0.056*
H5B	0.8297	0.1380	1.3842	0.056*
N6	0.55659 (15)	0.33821 (19)	1.1393 (3)	0.0463 (7)
H6A	0.5622	0.3608	1.2308	0.056*
H6B	0.5157	0.3542	1.0517	0.056*
N7	0.56565 (14)	0.37551 (19)	0.6915 (3)	0.0370 (6)
N8	0.58831 (15)	0.43984 (19)	0.7856 (3)	0.0372 (6)
N9	0.61108 (18)	0.5009 (2)	0.8783 (4)	0.0554 (8)
C1	0.5666 (2)	0.1303 (2)	0.5574 (4)	0.0486 (9)
H1A	0.5178	0.1421	0.4682	0.058*
C2	0.6245 (2)	0.0791 (3)	0.5352 (4)	0.0536 (10)
H2A	0.6149	0.0572	0.4327	0.064*
C3	0.6968 (2)	0.0602 (2)	0.6659 (4)	0.0452 (8)
H3A	0.7370	0.0257	0.6537	0.054*

supplementary materials

C4	0.70815 (19)	0.0941 (2)	0.8161 (4)	0.0375 (7)
H4A	0.7560	0.0818	0.9071	0.045*
C5	0.64771 (17)	0.14623 (19)	0.8293 (3)	0.0294 (6)
C6	0.65790 (16)	0.18691 (19)	0.9882 (3)	0.0276 (6)
C7	0.72914 (17)	0.1992 (2)	1.2585 (3)	0.0320 (7)
C8	0.61087 (17)	0.2789 (2)	1.1335 (3)	0.0311 (7)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Zn1	0.0264 (3)	0.0377 (3)	0.0228 (3)	0.000	0.00450 (19)	0.000
N1	0.0326 (13)	0.0405 (15)	0.0225 (13)	0.0043 (11)	0.0043 (11)	-0.0033 (10)
N2	0.0282 (13)	0.0402 (15)	0.0218 (13)	0.0042 (10)	0.0060 (10)	0.0004 (10)
N3	0.0307 (13)	0.0454 (16)	0.0206 (12)	0.0050 (11)	0.0075 (10)	0.0004 (10)
N4	0.0244 (12)	0.0399 (15)	0.0198 (12)	0.0021 (10)	0.0052 (10)	-0.0012 (10)
N5	0.0393 (15)	0.070 (2)	0.0202 (13)	0.0202 (14)	0.0027 (11)	-0.0020 (13)
N6	0.0393 (15)	0.070 (2)	0.0212 (13)	0.0195 (14)	0.0050 (12)	-0.0045 (12)
N7	0.0326 (14)	0.0448 (16)	0.0321 (14)	-0.0050 (12)	0.0123 (11)	-0.0025 (12)
N8	0.0285 (14)	0.0446 (17)	0.0338 (15)	-0.0006 (12)	0.0089 (11)	0.0082 (13)
N9	0.061 (2)	0.0489 (18)	0.0433 (17)	-0.0132 (15)	0.0104 (15)	-0.0083 (15)
C1	0.0435 (19)	0.063 (2)	0.0272 (17)	0.0092 (17)	0.0041 (15)	-0.0105 (16)
C2	0.059 (2)	0.065 (2)	0.0317 (18)	0.0114 (19)	0.0149 (17)	-0.0161 (16)
C3	0.051 (2)	0.046 (2)	0.042 (2)	0.0076 (16)	0.0222 (17)	-0.0088 (15)
C4	0.0379 (17)	0.0382 (18)	0.0335 (17)	0.0037 (14)	0.0125 (14)	-0.0006 (14)
C5	0.0316 (15)	0.0283 (15)	0.0265 (15)	-0.0020 (12)	0.0108 (12)	-0.0022 (12)
C6	0.0265 (14)	0.0303 (15)	0.0244 (15)	-0.0014 (12)	0.0094 (12)	0.0012 (12)
C7	0.0291 (15)	0.0372 (17)	0.0246 (15)	0.0008 (13)	0.0066 (12)	0.0047 (12)
C8	0.0284 (15)	0.0384 (17)	0.0231 (15)	0.0021 (13)	0.0078 (12)	0.0020 (12)

Geometric parameters (\AA , $^\circ$)

Zn1—N7	2.153 (3)	N5—H5B	0.8600
Zn1—N7 ⁱ	2.153 (3)	N6—C8	1.322 (4)
Zn1—N4 ⁱ	2.166 (2)	N6—H6A	0.8600
Zn1—N4	2.166 (2)	N6—H6B	0.8600
Zn1—N1	2.202 (2)	N7—N8	1.202 (4)
Zn1—N1 ⁱ	2.202 (2)	N8—N9	1.159 (4)
N1—C5	1.343 (3)	C1—C2	1.371 (5)
N1—C1	1.346 (4)	C1—H1A	0.9300
N2—C6	1.318 (3)	C2—C3	1.377 (5)
N2—C7	1.372 (4)	C2—H2A	0.9300
N3—C7	1.338 (4)	C3—C4	1.385 (4)
N3—C8	1.339 (4)	C3—H3A	0.9300
N4—C6	1.337 (3)	C4—C5	1.379 (4)
N4—C8	1.371 (4)	C4—H4A	0.9300
N5—C7	1.329 (3)	C5—C6	1.500 (4)
N5—H5A	0.8600		
N7—Zn1—N7 ⁱ	89.01 (14)	H6A—N6—H6B	120.0

N7—Zn1—N4 ⁱ	100.61 (9)	N8—N7—Zn1	115.1 (2)
N7 ⁱ —Zn1—N4 ⁱ	92.75 (9)	N9—N8—N7	178.8 (3)
N7—Zn1—N4	92.75 (9)	N1—C1—C2	123.1 (3)
N7 ⁱ —Zn1—N4	100.61 (9)	N1—C1—H1A	118.4
N4 ⁱ —Zn1—N4	161.28 (13)	C2—C1—H1A	118.4
N7—Zn1—N1	87.41 (10)	C1—C2—C3	119.4 (3)
N7 ⁱ —Zn1—N1	174.96 (9)	C1—C2—H2A	120.3
N4 ⁱ —Zn1—N1	91.39 (9)	C3—C2—H2A	120.3
N4—Zn1—N1	76.04 (9)	C2—C3—C4	118.2 (3)
N7—Zn1—N1 ⁱ	174.96 (9)	C2—C3—H3A	120.9
N7 ⁱ —Zn1—N1 ⁱ	87.41 (10)	C4—C3—H3A	120.9
N4 ⁱ —Zn1—N1 ⁱ	76.04 (9)	C5—C4—C3	119.3 (3)
N4—Zn1—N1 ⁱ	91.39 (9)	C5—C4—H4A	120.4
N1—Zn1—N1 ⁱ	96.38 (14)	C3—C4—H4A	120.4
C5—N1—C1	117.3 (3)	N1—C5—C4	122.7 (3)
C5—N1—Zn1	114.71 (18)	N1—C5—C6	115.9 (2)
C1—N1—Zn1	127.1 (2)	C4—C5—C6	121.4 (3)
C6—N2—C7	113.8 (2)	N2—C6—N4	127.0 (2)
C7—N3—C8	115.9 (2)	N2—C6—C5	116.2 (2)
C6—N4—C8	114.7 (2)	N4—C6—C5	116.8 (2)
C6—N4—Zn1	115.90 (17)	N5—C7—N3	119.1 (3)
C8—N4—Zn1	129.38 (19)	N5—C7—N2	116.0 (3)
C7—N5—H5A	120.0	N3—C7—N2	124.9 (2)
C7—N5—H5B	120.0	N6—C8—N3	118.4 (3)
H5A—N5—H5B	120.0	N6—C8—N4	118.1 (2)
C8—N6—H6A	120.0	N3—C8—N4	123.5 (3)
C8—N6—H6B	120.0		

Symmetry codes: (i) $-x+1, y, -z+3/2$.

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
N5—H5A···N3 ⁱⁱ	0.86	2.19	3.042 (4)	175
N5—H5B···N7 ⁱⁱⁱ	0.86	2.31	3.060 (4)	147
N6—H6A···N9 ^{iv}	0.86	2.34	3.025 (4)	137
N6—H6B···N7 ⁱ	0.86	2.10	2.939 (4)	164

Symmetry codes: (ii) $-x+3/2, -y+1/2, -z+3$; (iii) $-x+3/2, -y+1/2, -z+2$; (iv) $x, -y+1, z+1/2$; (i) $-x+1, y, -z+3/2$.

supplementary materials

Fig. 1




Fig. 2

