

Poly[(μ_2 -azido- $\kappa^2 N^1:N^1$)[μ_2 -5-(8-quinolyloxymethyl)tetrazolato- $\kappa^4 N^1,O,N^5:N^4$]zinc(II)]

Hong-ling Cai

Ordered Matter Science Research Center, Southeast University, Nanjing 210096, People's Republic of China

Correspondence e-mail: seuwangwei@gmail.com


Received 14 April 2009; accepted 5 May 2009

 Key indicators: single-crystal X-ray study; $T = 294$ K; mean $\sigma(C-C) = 0.007$ Å; R factor = 0.047; wR factor = 0.183; data-to-parameter ratio = 14.3.

In the title compound, $[Zn(C_{11}H_8N_5O)(N_3)]_n$, the Zn atom is hexacoordinated by five N atoms and one O atom in a distorted octahedral geometry. The chelating 5-(8-quinolyloxymethyl)tetrazolate ligands are approximately planar, with a dihedral angle of $3.6(2)^\circ$ between the quinoline and tetrazole planes. Adjacent Zn atoms are linked by two bridging azide ligands across a centre of inversion, and further coordination by one N atom of an adjacent tetrazole unit forms two-dimensional frameworks in (100). C—H...N interactions exist between ligands in neighbouring layers.

Related literature

For the use of tetrazole derivatives in coordination chemistry, see: Wang *et al.* (2005); Xiong *et al.* (2002). For details of the synthesis, see: Luo & Ye (2008). For related structures, see: Wang & Ye (2008); Chen & Ye (2008).



Experimental

Crystal data

$[Zn(C_{11}H_8N_5O)(N_3)]$
 $M_r = 333.64$
 Monoclinic, $P2_1/c$
 $a = 10.352(8)$ Å
 $b = 14.108(9)$ Å
 $c = 8.626(8)$ Å
 $\beta = 90.31(2)^\circ$

$V = 1259.8(17)$ Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 1.96$ mm⁻¹
 $T = 294$ K
 $0.18 \times 0.12 \times 0.10$ mm

Data collection

Rigaku SCXmini CCD diffractometer
 Absorption correction: multi-scan *CrystalClear* (Rigaku, 2005)
 $T_{\min} = 0.714$, $T_{\max} = 0.821$

11540 measured reflections
 2714 independent reflections
 2261 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.059$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.047$
 $wR(F^2) = 0.183$
 $S = 1.19$
 2714 reflections

190 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.63$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.71$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C3—H3A...N2 ⁱ	0.93	2.49	3.411 (8)	170
C11—H11B...N8 ⁱⁱ	0.97	2.54	3.252 (7)	130

 Symmetry codes: (i) $x - 1, y, z$; (ii) $-x + 2, y - \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: *CrystalClear* (Rigaku, 2005); cell refinement: *CrystalClear*; data reduction: *CrystalClear*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

This work was supported by a start-up grant from Southeast University.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: B12367).

References

- Chen, F. & Ye, H.-Y. (2008). *Acta Cryst.* **E64**, m1060.
 Luo, H.-Z. & Ye, H.-Y. (2008). *Acta Cryst.* **E64**, o136.
 Rigaku (2005). *CrystalClear*. Rigaku Corporation, Tokyo, Japan.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Wang, X.-S., Tang, Y.-Z., Huang, X.-F., Qu, Z.-R., Che, C.-M., Chan, C. W. H. & Xiong, R.-G. (2005). *Inorg. Chem.* **44**, 5278–5285.
 Wang, G.-X. & Ye, H.-Y. (2008). *Acta Cryst.* **E64**, m1006.
 Xiong, R.-G., Xue, X., Zhao, H., You, X.-Z., Abrahams, B. F. & Xue, Z.-L. (2002). *Angew. Chem. Int. Ed.* **41**, 3800–3803.

supplementary materials

Acta Cryst. (2009). E65, m632 [doi:10.1107/S1600536809016924]

Poly[(μ_2 -azido- $\kappa^2 N^1:N^1$)[μ_2 -5-(8-quinolyloxymethyl)tetrazolato- $\kappa^4 N^1,O,N^5:N^4$]zinc(II)]

H. Cai

Comment

As shown in Fig. 1, the Zn atom adopts a distorted octahedral coordination geometry, coordinated by one N atom and one O atom from 8-hydroxyquinoline, two N atoms from two different tetrazole groups and two N atoms from two bridging azide groups. Thus, 8-[(1*H*-tetrazol-5-yl)methoxy]quinoline acts as a tetradentate linker while the azide groups bridge between Zn atoms to form centrosymmetric rhombic units. Two-dimensional frameworks are formed in the (100) planes, and C—H \cdots N interactions exist between ligands in neighbouring planes.

Experimental

A mixture of quinolin-8-ol (1.45 g, 10 mmol), 1.38 g K₂CO₃, 30 ml acetone and 2-bromoacetonitrile (1.32 g, 11 mmol) was refluxed overnight. After cooling, the resulting dark mixture was extracted with ether (30 ml) and the solvent was removed at reduced pressure to give a crude product. Recrystallization from ethanol gave the pure ligand 2-(quinolin-8-yloxy)acetonitrile as a white powder. A mixture of this ligand (0.037 g, 0.2 mmol), ZnCl₂ (0.026 g, 0.2 mmol) and water (1 ml) was sealed in a glass tube and maintained at 383 K. Yellow crystals of the title compound suitable for X-ray analysis were obtained after 3 d.

Refinement

H atoms were placed geometrically with C—H = 0.93 or 0.97 Å and allowed to ride during refinement with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$.

Figures




Fig. 1. Molecular structure with displacement ellipsoids drawn at the 30% probability level. H atoms are omitted. Symmetry code: (A) $2 - x, 2 - y, -z$.




Fig. 2. Packing diagram viewed along the *a* axis.

Poly[(μ_2 -azido- κ^2 N¹:N¹)[μ_2 -5-(8-quinolyloxymethyl)tetrazolato- κ^4 N¹,O,N⁵:N⁴]zinc(II)]

Crystal data

[Zn(C₁₁H₈N₅O)(N₃)

M_r = 333.64

Monoclinic, *P*2₁/*c*

Hall symbol: -*P* 2ybc

a = 10.352 (8) Å

b = 14.108 (9) Å

c = 8.626 (8) Å

β = 90.31 (2)°

V = 1259.8 (17) Å³

Z = 4

*F*₀₀₀ = 672

D_x = 1.759 Mg m⁻³

Mo *K*α radiation

λ = 0.71073 Å

Cell parameters from 3428 reflections

θ = 2.0–27.3°

μ = 1.96 mm⁻¹

T = 294 K

Block, pale yellow

0.18 × 0.12 × 0.10 mm

Data collection

Rigaku SCXmini CCD
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

T = 295 K

ω scans

Absorption correction: multi-scan

CrystalClear (Rigaku, 2005)

*T*_{min} = 0.714, *T*_{max} = 0.821

11540 measured reflections

2714 independent reflections

2261 reflections with *I* > 2σ(*I*)

*R*_{int} = 0.059

θ _{max} = 27.3°

θ _{min} = 2.0°

h = -13→13

k = -18→18

l = -11→11

Refinement

Refinement on *F*²

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.047$

$wR(F^2) = 0.183$

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0764P)^2]$

where $P = (F_o^2 + 2F_c^2)/3$

$S = 1.19$ $(\Delta/\sigma)_{\max} < 0.001$
 2714 reflections $\Delta\rho_{\max} = 0.63 \text{ e } \text{Å}^{-3}$
 190 parameters $\Delta\rho_{\min} = -0.71 \text{ e } \text{Å}^{-3}$
 Primary atom site location: structure-invariant direct methods Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Zn1	0.95242 (4)	0.90443 (3)	0.09995 (5)	0.0298 (2)
N5	0.7775 (3)	0.9622 (3)	0.1958 (4)	0.0347 (8)
N1	1.0142 (3)	0.7920 (3)	-0.0386 (4)	0.0374 (8)
O1	0.7733 (3)	0.8144 (2)	0.0151 (4)	0.0420 (8)
C9	0.6528 (4)	0.8458 (3)	0.0556 (5)	0.0343 (9)
C8	0.5420 (4)	0.9627 (4)	0.2162 (6)	0.0447 (11)
C7	0.7806 (5)	1.0369 (3)	0.2914 (5)	0.0433 (11)
H7A	0.8605	1.0629	0.3173	0.052*
C6	0.6587 (4)	0.9259 (3)	0.1565 (5)	0.0326 (9)
C5	0.4240 (5)	0.9206 (4)	0.1716 (8)	0.0595 (15)
H5A	0.3469	0.9447	0.2100	0.071*
C4	0.5380 (4)	0.8062 (3)	0.0114 (6)	0.0471 (11)
H4A	0.5366	0.7553	-0.0571	0.056*
C3	0.4212 (5)	0.8432 (4)	0.0705 (8)	0.0621 (15)
H3A	0.3427	0.8161	0.0421	0.075*
C2	0.5503 (5)	1.0407 (4)	0.3162 (6)	0.0551 (14)
H2A	0.4755	1.0674	0.3566	0.066*
C11	0.7891 (4)	0.7426 (3)	-0.0998 (5)	0.0340 (9)
H11A	0.7493	0.7611	-0.1973	0.041*
H11B	0.7521	0.6829	-0.0661	0.041*
N2	1.1345 (4)	0.7624 (3)	-0.0749 (5)	0.0488 (10)
C10	0.9352 (4)	0.7353 (3)	-0.1149 (5)	0.0319 (8)
N6	1.0849 (3)	1.0102 (3)	0.1218 (4)	0.0351 (8)
N3	1.1247 (4)	0.6921 (3)	-0.1722 (5)	0.0504 (11)
C1	0.6690 (6)	1.0780 (4)	0.3545 (7)	0.0538 (13)
H1A	0.6753	1.1296	0.4212	0.065*
N7	1.1303 (4)	1.0413 (3)	0.2394 (4)	0.0382 (8)

supplementary materials

N8	1.1778 (6)	1.0723 (4)	0.3487 (6)	0.0664 (14)
N4	0.9978 (3)	0.6719 (3)	-0.1977 (4)	0.0371 (8)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Zn1	0.0276 (3)	0.0278 (3)	0.0339 (3)	-0.00349 (16)	0.0000 (2)	-0.00067 (16)
N5	0.0353 (18)	0.0345 (18)	0.0345 (18)	0.0002 (15)	0.0042 (15)	0.0007 (15)
N1	0.0323 (18)	0.039 (2)	0.041 (2)	0.0005 (15)	-0.0040 (16)	-0.0074 (16)
O1	0.0300 (15)	0.0434 (17)	0.0526 (19)	-0.0026 (13)	0.0022 (14)	-0.0189 (14)
C9	0.029 (2)	0.034 (2)	0.040 (2)	-0.0011 (16)	0.0015 (17)	0.0019 (17)
C8	0.032 (2)	0.049 (3)	0.053 (3)	0.011 (2)	0.005 (2)	0.006 (2)
C7	0.046 (3)	0.041 (2)	0.044 (2)	-0.004 (2)	0.003 (2)	-0.011 (2)
C6	0.028 (2)	0.036 (2)	0.034 (2)	-0.0015 (17)	0.0012 (17)	0.0041 (17)
C5	0.030 (3)	0.057 (3)	0.092 (5)	0.008 (2)	0.007 (3)	0.003 (3)
C4	0.036 (2)	0.044 (3)	0.061 (3)	-0.006 (2)	-0.005 (2)	0.004 (2)
C3	0.030 (2)	0.069 (4)	0.087 (4)	-0.005 (2)	-0.005 (3)	-0.001 (3)
C2	0.044 (3)	0.061 (3)	0.061 (3)	0.016 (2)	0.015 (2)	-0.012 (3)
C11	0.038 (2)	0.031 (2)	0.033 (2)	-0.0013 (17)	0.0002 (17)	-0.0049 (16)
N2	0.037 (2)	0.050 (2)	0.059 (3)	0.0069 (18)	-0.0083 (19)	-0.011 (2)
C10	0.039 (2)	0.0271 (19)	0.0290 (18)	-0.0014 (16)	-0.0001 (16)	-0.0019 (15)
N6	0.0348 (18)	0.0364 (19)	0.0341 (18)	-0.0118 (15)	-0.0039 (15)	0.0024 (14)
N3	0.036 (2)	0.054 (3)	0.061 (3)	0.0023 (18)	-0.0036 (19)	-0.019 (2)
C1	0.055 (3)	0.049 (3)	0.057 (3)	0.002 (2)	0.012 (3)	-0.017 (3)
N7	0.041 (2)	0.0360 (19)	0.0371 (19)	-0.0044 (16)	-0.0013 (16)	-0.0009 (15)
N8	0.089 (4)	0.062 (3)	0.048 (3)	-0.012 (3)	-0.018 (3)	-0.013 (2)
N4	0.0325 (18)	0.0384 (19)	0.0403 (19)	0.0055 (15)	-0.0001 (15)	-0.0099 (16)

Geometric parameters (\AA , $^\circ$)

Zn1—N6	2.035 (4)	C5—C3	1.398 (9)
Zn1—N1	2.088 (4)	C5—H5A	0.930
Zn1—N4 ⁱ	2.102 (4)	C4—C3	1.415 (7)
Zn1—N5	2.155 (4)	C4—H4A	0.930
Zn1—N6 ⁱⁱ	2.291 (4)	C3—H3A	0.930
Zn1—O1	2.360 (4)	C2—C1	1.376 (8)
N5—C7	1.338 (6)	C2—H2A	0.930
N5—C6	1.373 (6)	C11—C10	1.522 (6)
N1—C10	1.318 (5)	C11—H11A	0.970
N1—N2	1.352 (5)	C11—H11B	0.970
O1—C9	1.371 (5)	N2—N3	1.302 (6)
O1—C11	1.428 (5)	C10—N4	1.318 (5)
C9—C4	1.366 (6)	N6—N7	1.199 (5)
C9—C6	1.426 (6)	N6—Zn1 ⁱⁱ	2.291 (4)
C8—C2	1.401 (7)	N3—N4	1.361 (6)
C8—C5	1.410 (8)	C1—H1A	0.930
C8—C6	1.415 (6)	N7—N8	1.147 (6)
C7—C1	1.405 (7)	N4—Zn1 ⁱⁱⁱ	2.102 (4)

C7—H7A	0.930		
N6—Zn1—N1	113.67 (17)	C8—C6—C9	118.6 (4)
N6—Zn1—N4 ⁱ	98.71 (15)	C3—C5—C8	120.9 (5)
N1—Zn1—N4 ⁱ	91.07 (17)	C3—C5—H5A	119.5
N6—Zn1—N5	104.72 (17)	C8—C5—H5A	119.5
N1—Zn1—N5	140.15 (14)	C9—C4—C3	119.5 (5)
N4 ⁱ —Zn1—N5	93.41 (15)	C9—C4—H4A	120.2
N6—Zn1—N6 ⁱⁱ	78.59 (15)	C3—C4—H4A	120.2
N1—Zn1—N6 ⁱⁱ	88.40 (17)	C5—C3—C4	119.9 (5)
N4 ⁱ —Zn1—N6 ⁱⁱ	176.76 (13)	C5—C3—H3A	120.1
N5—Zn1—N6 ⁱⁱ	89.04 (15)	C4—C3—H3A	120.1
N6—Zn1—O1	162.00 (14)	C1—C2—C8	120.0 (5)
N1—Zn1—O1	69.91 (14)	C1—C2—H2A	120.0
N4 ⁱ —Zn1—O1	98.84 (14)	C8—C2—H2A	120.0
N5—Zn1—O1	70.27 (15)	O1—C11—C10	103.0 (3)
N6 ⁱⁱ —Zn1—O1	83.98 (13)	O1—C11—H11A	111.2
C7—N5—C6	117.7 (4)	C10—C11—H11A	111.2
C7—N5—Zn1	121.1 (3)	O1—C11—H11B	111.2
C6—N5—Zn1	121.1 (3)	C10—C11—H11B	111.2
C10—N1—N2	105.5 (4)	H11A—C11—H11B	109.1
C10—N1—Zn1	123.8 (3)	N3—N2—N1	108.4 (4)
N2—N1—Zn1	130.7 (3)	N4—C10—N1	112.2 (4)
C9—O1—C11	120.9 (3)	N4—C10—C11	125.7 (4)
C9—O1—Zn1	117.5 (3)	N1—C10—C11	122.1 (4)
C11—O1—Zn1	120.3 (2)	N7—N6—Zn1	127.4 (3)
C4—C9—O1	126.0 (4)	N7—N6—Zn1 ⁱⁱ	125.3 (3)
C4—C9—C6	121.9 (4)	Zn1—N6—Zn1 ⁱⁱ	101.41 (15)
O1—C9—C6	112.0 (4)	N2—N3—N4	109.7 (4)
C2—C8—C5	123.3 (5)	C2—C1—C7	119.0 (5)
C2—C8—C6	117.6 (5)	C2—C1—H1A	120.5
C5—C8—C6	119.1 (5)	C7—C1—H1A	120.5
N5—C7—C1	123.1 (5)	N8—N7—N6	177.4 (5)
N5—C7—H7A	118.4	C10—N4—N3	104.3 (4)
C1—C7—H7A	118.4	C10—N4—Zn1 ⁱⁱⁱ	133.6 (3)
N5—C6—C8	122.5 (4)	N3—N4—Zn1 ⁱⁱⁱ	117.0 (3)
N5—C6—C9	118.8 (4)		
N6—Zn1—N5—C7	-19.6 (4)	C5—C8—C6—C9	-1.4 (7)
N1—Zn1—N5—C7	176.1 (3)	C4—C9—C6—N5	-179.8 (4)
N4 ⁱ —Zn1—N5—C7	80.3 (4)	O1—C9—C6—N5	1.8 (6)
N6 ⁱⁱ —Zn1—N5—C7	-97.6 (4)	C4—C9—C6—C8	2.3 (7)
O1—Zn1—N5—C7	178.5 (4)	O1—C9—C6—C8	-176.1 (4)
N6—Zn1—N5—C6	158.0 (3)	C2—C8—C5—C3	179.5 (5)
N1—Zn1—N5—C6	-6.3 (4)	C6—C8—C5—C3	0.4 (9)
N4 ⁱ —Zn1—N5—C6	-102.1 (3)	O1—C9—C4—C3	176.1 (5)
N6 ⁱⁱ —Zn1—N5—C6	80.0 (3)	C6—C9—C4—C3	-2.0 (7)

supplementary materials

O1—Zn1—N5—C6	-3.9 (3)	C8—C5—C3—C4	-0.1 (10)
N6—Zn1—N1—C10	-154.1 (3)	C9—C4—C3—C5	0.9 (9)
N4 ⁱ —Zn1—N1—C10	106.0 (4)	C5—C8—C2—C1	-180.0 (6)
N5—Zn1—N1—C10	9.3 (5)	C6—C8—C2—C1	-0.8 (8)
N6 ⁱⁱ —Zn1—N1—C10	-77.3 (4)	C9—O1—C11—C10	176.3 (4)
O1—Zn1—N1—C10	6.9 (3)	Zn1—O1—C11—C10	9.5 (4)
N6—Zn1—N1—N2	23.8 (5)	C10—N1—N2—N3	1.5 (5)
N4 ⁱ —Zn1—N1—N2	-76.2 (4)	Zn1—N1—N2—N3	-176.6 (3)
N5—Zn1—N1—N2	-172.8 (3)	N2—N1—C10—N4	-0.3 (5)
N6 ⁱⁱ —Zn1—N1—N2	100.6 (4)	Zn1—N1—C10—N4	178.0 (3)
O1—Zn1—N1—N2	-175.2 (4)	N2—N1—C10—C11	177.4 (4)
N6—Zn1—O1—C9	-71.8 (5)	Zn1—N1—C10—C11	-4.3 (6)
N1—Zn1—O1—C9	-176.7 (3)	O1—C11—C10—N4	173.4 (4)
N4 ⁱ —Zn1—O1—C9	95.4 (3)	O1—C11—C10—N1	-3.9 (5)
N5—Zn1—O1—C9	4.9 (3)	N1—Zn1—N6—N7	-123.2 (4)
N6 ⁱⁱ —Zn1—O1—C9	-86.2 (3)	N4 ⁱ —Zn1—N6—N7	-28.2 (4)
N6—Zn1—O1—C11	95.5 (5)	N5—Zn1—N6—N7	67.7 (4)
N1—Zn1—O1—C11	-9.5 (3)	N6 ⁱⁱ —Zn1—N6—N7	153.7 (5)
N4 ⁱ —Zn1—O1—C11	-97.4 (3)	O1—Zn1—N6—N7	139.0 (4)
N5—Zn1—O1—C11	172.1 (3)	N1—Zn1—N6—Zn1 ⁱⁱ	83.13 (19)
N6 ⁱⁱ —Zn1—O1—C11	81.0 (3)	N4 ⁱ —Zn1—N6—Zn1 ⁱⁱ	178.17 (15)
C11—O1—C9—C4	9.5 (7)	N5—Zn1—N6—Zn1 ⁱⁱ	-85.94 (17)
Zn1—O1—C9—C4	176.6 (4)	N6 ⁱⁱ —Zn1—N6—Zn1 ⁱⁱ	0.0
C11—O1—C9—C6	-172.2 (4)	O1—Zn1—N6—Zn1 ⁱⁱ	-14.7 (5)
Zn1—O1—C9—C6	-5.1 (5)	N1—N2—N3—N4	-2.1 (6)
C6—N5—C7—C1	1.2 (7)	C8—C2—C1—C7	0.3 (9)
Zn1—N5—C7—C1	178.9 (4)	N5—C7—C1—C2	-0.5 (9)
C7—N5—C6—C8	-1.8 (6)	N1—C10—N4—N3	-1.0 (5)
Zn1—N5—C6—C8	-179.4 (3)	C11—C10—N4—N3	-178.5 (4)
C7—N5—C6—C9	-179.6 (4)	N1—C10—N4—Zn1 ⁱⁱⁱ	-153.6 (3)
Zn1—N5—C6—C9	2.8 (5)	C11—C10—N4—Zn1 ⁱⁱⁱ	28.8 (7)
C2—C8—C6—N5	1.6 (7)	N2—N3—N4—C10	1.9 (5)
C5—C8—C6—N5	-179.2 (5)	N2—N3—N4—Zn1 ⁱⁱⁱ	160.0 (3)
C2—C8—C6—C9	179.4 (4)		

Symmetry codes: (i) $x, -y+3/2, z+1/2$; (ii) $-x+2, -y+2, -z$; (iii) $x, -y+3/2, z-1/2$.

Hydrogen-bond geometry ($\text{\AA}, ^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C3—H3A \cdots N2 ^{iv}	0.93	2.49	3.411 (8)	170
C11—H11B \cdots N8 ^v	0.97	2.54	3.252 (7)	130

Symmetry codes: (iv) $x-1, y, z$; (v) $-x+2, y-1/2, -z+1/2$.

Fig. 1




Fig. 2

