

Ethyl 6-(4-chlorophenyl)-4-(4-methoxyphenyl)-2-oxocyclohex-3-ene-1-carboxylate

Hoong-Kun Fun,^{a,*‡} Samuel Robinson Jebas,^{a,§}
K. S. Girish^b and Balakrishna Kalluraya^b

^aX-ray Crystallography Unit, School of Physics, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia, and ^bDepartment of Studies in Chemistry, Mangalore University, Mangalagangotri, Mangalore 574 199, India
Correspondence e-mail: hkfun@usm.my


Received 27 April 2009; accepted 30 April 2009

Key indicators: single-crystal X-ray study; $T = 100\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$; R factor = 0.072; wR factor = 0.180; data-to-parameter ratio = 30.5.

In the title compound, $\text{C}_{22}\text{H}_{21}\text{ClO}_4$, the cyclohex-3-ene unit adopts an envelope conformation in both independent molecules comprising the asymmetric unit. The two benzene rings are inclined to each other at a dihedral angle of $82.03(5)^\circ$ [$86.37(5)^\circ$]. In the crystal, the molecules interact via $\text{C}-\text{H}\cdots\text{O}$, $\text{C}-\text{H}\cdots\text{Cl}$ and $\text{C}-\text{H}\cdots\pi$ interactions.

Related literature

For the biological activity of cyclohexenones, see: Hamon *et al.* (1996); Honda (2002); Keil *et al.* (1996). For green chemistry, see: Hoel & Nielsen (1999); Larhed *et al.* (1999). For ring puckering analysis, see: Cremer & Pople (1975). For stability of the temperature controller used in the data collection, see: Cosier & Glazer (1986).



Experimental

Crystal data

$\text{C}_{22}\text{H}_{21}\text{ClO}_4$
 $M_r = 384.84$

Monoclinic, $P2_1/c$
 $a = 11.9729(3)\text{ \AA}$

‡ Thomson Reuters ResearcherID: A-3561-2009.

§ Thomson Reuters ResearcherID: A-5473-2009. Permanent address: Department of Physics, Karunya University, Karunya Nagar, Coimbatore 641 114, India.

$b = 8.1713(2)\text{ \AA}$
 $c = 39.2033(8)\text{ \AA}$
 $\beta = 98.990(1)^\circ$
 $V = 3788.31(15)\text{ \AA}^3$
 $Z = 8$

Mo $K\alpha$ radiation
 $\mu = 0.23\text{ mm}^{-1}$
 $T = 100\text{ K}$
 $0.57 \times 0.40 \times 0.17\text{ mm}$

Data collection

Bruker SMART APEXII CCD area-detector diffractometer
Absorption correction: multi-scan (*SADABS*; Bruker, 2005)
 $T_{\min} = 0.882$, $T_{\max} = 0.962$

60118 measured reflections
14422 independent reflections
10971 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.043$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.072$
 $wR(F^2) = 0.180$
 $S = 1.16$
14422 reflections

473 parameters
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 1.03\text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.96\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C11A—H11A \cdots O1B ⁱ	0.98	2.53	3.492 (2)	167
C11B—H11B \cdots O1A ⁱⁱ	0.98	2.53	3.501 (2)	170
C12A—H12B \cdots O2B	0.97	2.51	3.450 (2)	162
C12B—H12C \cdots O2A ⁱⁱⁱ	0.97	2.56	3.441 (2)	151
C15B—H15B \cdots O4B ^{iv}	0.93	2.59	3.485 (3)	163
C20B—H20D \cdots C11A ^{iv}	0.97	2.83	3.585 (2)	136
C22A—H22A \cdots Cg1 ⁱⁱ	0.97	2.83	3.666 (2)	146

Symmetry codes: (i) $-x + 1, y - \frac{1}{2}, -z + \frac{3}{2}$; (ii) $-x + 1, y + \frac{1}{2}, -z + \frac{3}{2}$; (iii) $x - 1, y, z$; (iv) $x, y + 1, z$.

Data collection: *APEX2* (Bruker, 2005); cell refinement: *SAINT* (Bruker, 2005); data reduction: *SAINT*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL* and *PLATON* (Spek, 2009).

HKF and SRJ thank the Malaysian Government and Universiti Sains Malaysia for the Science Fund grant No. 305/PFIZIK/613312. SRJ thanks Universiti Sains Malaysia for a post-doctoral research fellowship. HKF also thanks Universiti Sains Malaysia for the Research University Golden Goose grant No. 1001/PFIZIK/811012.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: TK2440).

References

- Bruker (2005). *APEX2, SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Cosier, J. & Glazer, A. M. (1986). *J. Appl. Cryst.* **19**, 105–107.
- Cremer, D. & Pople, J. A. (1975). *J. Am. Chem. Soc.* **97**, 1354–1358.
- Hamon, D. P. G., Hayball, P. J., Massy-Westropp, R. A., Newton, J. L. & Tamblyn, J. G. (1996). *Tetrahedron Asymmetry*, **7**, 263–272.
- Hoel, A. M. L. & Nielsen, J. (1999). *Tetrahedron Lett.* **40**, 3941–3944.
- Honda (2002). *J. Synth. Org. Chem. Jpn.* **60**, 1104–1111.
- Keil, M., Schirmer, U., Kolassa, D., Kast, J., Wuerzer, B. & Meyer, N. (1996). US Patent No. 5 554 582.
- Larhed, M., Lindeberg, G. & Hallberg, A. (1999). *Tetrahedron Lett.* **37**, 8219–8222.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.

supplementary materials

Acta Cryst. (2009). E65, o1235 [doi:10.1107/S1600536809016237]

Ethyl 6-(4-chlorophenyl)-4-(4-methoxyphenyl)-2-oxocyclohex-3-ene-1-carboxylate

H.-K. Fun, S. R. Jebas, K. S. Girish and B. Kalluraya

Comment

Cyclohexenones are important intermediates in the synthesis of a wide variety of biologically active simple, condensed and bridged heterocycles. These cyclohexenones are also found to possess various types of biological activity, e.g. herbicidal (Keil *et al.*, 1996), in vitro inhibition of human platelet cylooxygenase (Hamon *et al.*, 1996), as an HMG-CoA reductase inhibitor (Honda, 2002), and displays anti-obesity properties (Honda, 2002). Today green chemistry plays an important role in chemical research. The large number of publications clearly indicates the development of this area of chemistry (Hoel & Nielsen, 1999; Larhed *et al.*, 1999).

The asymmetric unit of (I) (Fig. 1) comprises of two crystallographically independent molecules (A & B) with almost similar geometries. The cyclohex-3-ene unit in both the molecules adopt an envelope conformation with puckering parameters $Q = 0.5107 (18)$ Å, $\theta = 124.6 (2)$ ° and $\phi = 47.7 (2)$ ° for molecule A, and $Q = 0.50 (18)$ Å, $\theta = 56.0 (2)$ ° and $\phi = 231.5 (2)$ ° for molecule B (Cremer & Pople, 1975). The two benzene rings are inclined to each other forming dihedral angles of $66.82 (6)$ ° (C1A—C6A:C13A—C18A) in molecule A and $73.68 (5)$ ° (C1B—C6B: C13B—C18B) in molecule B.

Globally, the crystal packing comprises layers stabilized by C—H···O and C—H···Cl contacts (Fig. 2) together with C—H···π interactions (Table 1).

Experimental

1-(*p*-Methoxyphenyl)-3-(*p*-chlorophenyl)-2-propene-1-one (0.01 mol), potassium carbonate (0.04 mol), ethyl acetoacetate (0.02 mol) were ground in a mortar using a pestle for uniform mixing. The paste formed was transferred to a 50 mL beaker and placed in a microwave oven operating at 160 W for 5 mins. The product (I) was poured into cold water, filtered, dried and recrystallized from ethanol-dioxane mixture; m. pt. = 411 - 412 K.

Refinement

H atoms were positioned geometrically [C—H = 0.93–0.98 Å] and refined using a riding model with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ and $1.5U_{\text{eq}}(\text{methyl C})$. A rotating-group model was used for the methyl groups. The maximum and minimum residual electron density peaks of 1.03 and -0.96 eÅ⁻³, respectively, were located 0.08 Å and 0.05 Å from the C20A and C21A atoms, respectively

supplementary materials

Figures




Fig. 1. The molecular structure of (I), showing 50% probability displacement ellipsoids and the atom numbering scheme.




Fig. 2. The crystal packing for (I), viewed along the b axis. Dashed lines indicate $\text{C}—\text{H}\cdots\text{O}$ and $\text{C}—\text{H}\cdots\text{Cl}$ contacts.

Ethyl 6-(4-chlorophenyl)-4-(4-methoxyphenyl)-2-oxocyclohex-3-ene-1-carboxylate

Crystal data

$\text{C}_{22}\text{H}_{21}\text{ClO}_4$	$F_{000} = 1616$
$M_r = 384.84$	$D_x = 1.349 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
Hall symbol: -P 2ybc	$\lambda = 0.71073 \text{ \AA}$
$a = 11.9729 (3) \text{ \AA}$	Cell parameters from 9875 reflections
$b = 8.1713 (2) \text{ \AA}$	$\theta = 2.7\text{--}33.0^\circ$
$c = 39.2033 (8) \text{ \AA}$	$\mu = 0.23 \text{ mm}^{-1}$
$\beta = 98.990 (1)^\circ$	$T = 100 \text{ K}$
$V = 3788.31 (15) \text{ \AA}^3$	Plate, colourless
$Z = 8$	$0.57 \times 0.40 \times 0.17 \text{ mm}$

Data collection

Bruker SMART APEXII CCD area-detector diffractometer	14422 independent reflections
Radiation source: fine-focus sealed tube	10971 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.043$
$T = 100 \text{ K}$	$\theta_{\text{max}} = 33.2^\circ$
φ and ω scans	$\theta_{\text{min}} = 2.6^\circ$
Absorption correction: multi-scan (SADABS; Bruker, 2005)	$h = -17\text{--}18$
$T_{\text{min}} = 0.882$, $T_{\text{max}} = 0.962$	$k = -12\text{--}11$
60118 measured reflections	$l = -60\text{--}60$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.072$	H-atom parameters constrained
$wR(F^2) = 0.180$	$w = 1/[\sigma^2(F_o^2) + (0.064P)^2 + 2.9459P]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.16$	$(\Delta/\sigma)_{\max} < 0.002$
14422 reflections	$\Delta\rho_{\max} = 1.03 \text{ e \AA}^{-3}$
473 parameters	$\Delta\rho_{\min} = -0.96 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

Special details

Experimental. The crystal was placed in the cold stream of an Oxford Cyrosystems Cobra open-flow nitrogen cryostat (Cosier & Glazer, 1986) operating at 100.0 (1) K.

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Cl1A	0.76330 (4)	0.14248 (6)	0.969549 (11)	0.02789 (11)
O1A	0.45467 (10)	0.75170 (17)	0.66944 (3)	0.0223 (2)
O2A	0.94698 (12)	1.01249 (18)	0.84718 (3)	0.0291 (3)
O3A	1.04004 (14)	0.7270 (2)	0.89626 (4)	0.0409 (4)
O4A	0.92250 (12)	0.85217 (18)	0.92726 (3)	0.0284 (3)
C1A	0.56825 (14)	0.6458 (2)	0.75979 (4)	0.0203 (3)
H1	0.5517	0.5806	0.7778	0.024*
C2A	0.49553 (14)	0.6428 (2)	0.72817 (4)	0.0202 (3)
H2	0.4320	0.5758	0.7252	0.024*
C3A	0.51899 (13)	0.7407 (2)	0.70113 (4)	0.0179 (3)
C4A	0.61617 (15)	0.8382 (2)	0.70575 (4)	0.0204 (3)
H3	0.6326	0.9031	0.6877	0.024*
C5A	0.68769 (15)	0.8387 (2)	0.73694 (4)	0.0199 (3)
H4	0.7525	0.9031	0.7395	0.024*
C6A	0.66492 (13)	0.7440 (2)	0.76511 (4)	0.0169 (3)
C7A	0.73704 (14)	0.7533 (2)	0.79924 (4)	0.0177 (3)

supplementary materials

C8A	0.81593 (15)	0.8723 (2)	0.80671 (4)	0.0205 (3)
H5	0.8307	0.9400	0.7889	0.025*
C9A	0.87892 (15)	0.9002 (2)	0.84112 (4)	0.0220 (3)
C10A	0.84923 (15)	0.7906 (2)	0.86985 (4)	0.0220 (3)
H10A	0.7841	0.8370	0.8788	0.026*
C11A	0.81864 (15)	0.6186 (2)	0.85548 (4)	0.0216 (3)
H11A	0.8830	0.5787	0.8452	0.026*
C12A	0.71788 (14)	0.6307 (2)	0.82654 (4)	0.0190 (3)
H12A	0.7036	0.5240	0.8159	0.023*
H12B	0.6513	0.6621	0.8363	0.023*
C13A	0.80016 (15)	0.4985 (2)	0.88364 (4)	0.0218 (3)
C14A	0.87730 (15)	0.3722 (2)	0.89192 (4)	0.0228 (3)
H14A	0.9380	0.3619	0.8799	0.027*
C15A	0.86513 (15)	0.2608 (2)	0.91800 (4)	0.0227 (3)
H15A	0.9168	0.1760	0.9232	0.027*
C16A	0.77543 (15)	0.2776 (2)	0.93598 (4)	0.0211 (3)
C17A	0.69563 (16)	0.4008 (3)	0.92803 (5)	0.0268 (4)
H17A	0.6348	0.4100	0.9400	0.032*
C18A	0.70848 (17)	0.5101 (3)	0.90174 (5)	0.0274 (4)
H18A	0.6552	0.5925	0.8961	0.033*
C19A	0.94980 (17)	0.7852 (2)	0.89875 (5)	0.0250 (3)
C20A	1.0027 (2)	0.8377 (3)	0.95884 (5)	0.0350 (3)
H20A	1.0111	0.9422	0.9707	0.042*
H20B	1.0760	0.8047	0.9536	0.042*
C21A	0.9589 (2)	0.7117 (3)	0.98119 (5)	0.0350 (3)
H21A	1.0087	0.7043	1.0028	0.052*
H21B	0.9551	0.6074	0.9698	0.052*
H21C	0.8847	0.7426	0.9853	0.052*
C22A	0.35991 (15)	0.6440 (3)	0.66210 (5)	0.0261 (4)
H22A	0.3232	0.6610	0.6388	0.039*
H22B	0.3075	0.6660	0.6777	0.039*
H22C	0.3853	0.5326	0.6648	0.039*
C11B	0.23527 (5)	1.53507 (8)	0.970110 (14)	0.04101 (15)
O1B	-0.03768 (10)	0.91313 (17)	0.67571 (3)	0.0228 (3)
O2B	0.45811 (13)	0.65728 (19)	0.85379 (4)	0.0322 (3)
O3B	0.55166 (14)	0.9617 (3)	0.89936 (5)	0.0459 (4)
O4B	0.45395 (15)	0.8167 (2)	0.93271 (4)	0.0410 (3)
C1B	0.07581 (14)	1.0158 (2)	0.76615 (4)	0.0208 (3)
H6	0.0588	1.0796	0.7843	0.025*
C2B	0.00365 (14)	1.0205 (2)	0.73455 (4)	0.0213 (3)
H7	-0.0596	1.0881	0.7317	0.026*
C3B	0.02715 (14)	0.9233 (2)	0.70740 (4)	0.0183 (3)
C4B	0.12421 (14)	0.8254 (2)	0.71188 (4)	0.0200 (3)
H8	0.1404	0.7607	0.6938	0.024*
C5B	0.19597 (14)	0.8244 (2)	0.74301 (4)	0.0196 (3)
H9	0.2610	0.7606	0.7454	0.023*
C6B	0.17279 (13)	0.9183 (2)	0.77137 (4)	0.0171 (3)
C7B	0.24481 (13)	0.9098 (2)	0.80548 (4)	0.0177 (3)
C8B	0.32596 (15)	0.7943 (2)	0.81310 (4)	0.0211 (3)

H10	0.3424	0.7277	0.7953	0.025*
C9B	0.38899 (15)	0.7684 (2)	0.84758 (5)	0.0233 (3)
C10B	0.36130 (15)	0.8793 (2)	0.87657 (4)	0.0230 (3)
H10B	0.2999	0.8306	0.8870	0.028*
C11B	0.32456 (15)	1.0481 (2)	0.86175 (4)	0.0224 (3)
H11B	0.3873	1.0929	0.8514	0.027*
C12B	0.22305 (14)	1.0301 (2)	0.83289 (4)	0.0199 (3)
H12C	0.1579	0.9941	0.8428	0.024*
H12D	0.2052	1.1361	0.8223	0.024*
C13B	0.30004 (16)	1.1681 (2)	0.88921 (5)	0.0238 (3)
C14B	0.36628 (16)	1.3074 (2)	0.89537 (4)	0.0249 (3)
H14B	0.4254	1.3244	0.8830	0.030*
C15B	0.34550 (17)	1.4219 (3)	0.91981 (5)	0.0277 (4)
H15B	0.3898	1.5156	0.9236	0.033*
C16B	0.25847 (17)	1.3951 (3)	0.93839 (5)	0.0284 (4)
C17B	0.19055 (19)	1.2581 (3)	0.93277 (6)	0.0389 (5)
H17B	0.1315	1.2420	0.9452	0.047*
C18B	0.21188 (19)	1.1445 (3)	0.90817 (6)	0.0362 (5)
H18B	0.1668	1.0515	0.9043	0.043*
C19B	0.46667 (17)	0.8930 (3)	0.90359 (5)	0.0275 (4)
C20B	0.5512 (2)	0.8290 (3)	0.96065 (6)	0.0410 (3)
H20C	0.6171	0.7776	0.9537	0.049*
H20D	0.5688	0.9429	0.9659	0.049*
C21B	0.5210 (2)	0.7456 (3)	0.99117 (6)	0.0410 (3)
H21D	0.5807	0.7601	1.0104	0.062*
H21E	0.5104	0.6310	0.9863	0.062*
H21F	0.4523	0.7914	0.9967	0.062*
C22B	-0.12906 (15)	1.0268 (3)	0.66798 (5)	0.0269 (4)
H22D	-0.1659	1.0104	0.6447	0.040*
H22E	-0.1002	1.1364	0.6705	0.040*
H22F	-0.1824	1.0097	0.6836	0.040*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cl1A	0.0339 (2)	0.0278 (2)	0.02181 (18)	-0.00302 (18)	0.00375 (15)	0.00647 (16)
O1A	0.0201 (6)	0.0265 (7)	0.0197 (5)	-0.0030 (5)	0.0011 (4)	0.0015 (5)
O2A	0.0337 (7)	0.0281 (7)	0.0243 (6)	-0.0102 (6)	0.0009 (5)	-0.0009 (5)
O3A	0.0383 (9)	0.0536 (11)	0.0294 (7)	0.0118 (8)	0.0007 (6)	-0.0097 (7)
O4A	0.0331 (7)	0.0305 (8)	0.0202 (6)	0.0055 (6)	-0.0002 (5)	0.0003 (5)
C1A	0.0223 (7)	0.0186 (8)	0.0208 (7)	-0.0019 (6)	0.0058 (6)	0.0031 (6)
C2A	0.0188 (7)	0.0208 (8)	0.0218 (7)	-0.0027 (6)	0.0053 (5)	0.0007 (6)
C3A	0.0181 (7)	0.0171 (7)	0.0189 (7)	0.0018 (6)	0.0042 (5)	-0.0003 (5)
C4A	0.0238 (8)	0.0202 (8)	0.0176 (7)	-0.0035 (6)	0.0047 (5)	0.0017 (6)
C5A	0.0221 (7)	0.0196 (8)	0.0187 (7)	-0.0041 (6)	0.0047 (5)	-0.0004 (6)
C6A	0.0187 (7)	0.0151 (7)	0.0175 (6)	0.0002 (5)	0.0047 (5)	-0.0001 (5)
C7A	0.0194 (7)	0.0167 (7)	0.0175 (6)	0.0029 (6)	0.0047 (5)	0.0008 (5)
C8A	0.0246 (8)	0.0191 (8)	0.0182 (7)	-0.0012 (6)	0.0041 (6)	0.0015 (6)

supplementary materials

C9A	0.0263 (8)	0.0206 (8)	0.0188 (7)	-0.0007 (6)	0.0030 (6)	-0.0007 (6)
C10A	0.0254 (8)	0.0204 (8)	0.0202 (7)	0.0028 (6)	0.0041 (6)	-0.0007 (6)
C11A	0.0235 (8)	0.0229 (9)	0.0187 (7)	0.0020 (6)	0.0040 (6)	0.0011 (6)
C12A	0.0210 (7)	0.0171 (8)	0.0189 (7)	0.0011 (6)	0.0035 (5)	0.0017 (6)
C13A	0.0236 (8)	0.0215 (8)	0.0193 (7)	-0.0003 (6)	0.0005 (6)	0.0026 (6)
C14A	0.0230 (8)	0.0245 (9)	0.0209 (7)	0.0004 (6)	0.0032 (6)	0.0017 (6)
C15A	0.0225 (8)	0.0221 (9)	0.0228 (7)	0.0031 (6)	0.0015 (6)	0.0026 (6)
C16A	0.0251 (8)	0.0211 (8)	0.0166 (6)	-0.0017 (6)	0.0013 (6)	0.0019 (6)
C17A	0.0267 (9)	0.0311 (10)	0.0237 (8)	0.0059 (7)	0.0076 (6)	0.0046 (7)
C18A	0.0279 (9)	0.0282 (10)	0.0264 (8)	0.0088 (7)	0.0054 (7)	0.0076 (7)
C19A	0.0312 (9)	0.0228 (9)	0.0199 (7)	0.0003 (7)	0.0009 (6)	0.0000 (6)
C20A	0.0417 (8)	0.0371 (9)	0.0238 (6)	-0.0014 (7)	-0.0021 (5)	0.0038 (6)
C21A	0.0417 (8)	0.0371 (9)	0.0238 (6)	-0.0014 (7)	-0.0021 (5)	0.0038 (6)
C22A	0.0185 (7)	0.0328 (10)	0.0266 (8)	-0.0043 (7)	0.0025 (6)	-0.0035 (7)
Cl1B	0.0409 (3)	0.0450 (3)	0.0356 (3)	0.0091 (2)	0.0014 (2)	-0.0205 (2)
O1B	0.0202 (6)	0.0274 (7)	0.0204 (5)	0.0026 (5)	0.0024 (4)	0.0002 (5)
O2B	0.0327 (7)	0.0336 (8)	0.0290 (7)	0.0127 (6)	0.0011 (5)	0.0021 (6)
O3B	0.0327 (8)	0.0601 (12)	0.0425 (9)	-0.0162 (8)	-0.0013 (7)	0.0143 (8)
O4B	0.0416 (6)	0.0488 (7)	0.0299 (5)	-0.0111 (5)	-0.0030 (4)	0.0040 (5)
C1B	0.0216 (7)	0.0197 (8)	0.0218 (7)	0.0028 (6)	0.0058 (6)	-0.0030 (6)
C2B	0.0187 (7)	0.0230 (8)	0.0228 (7)	0.0044 (6)	0.0055 (6)	-0.0004 (6)
C3B	0.0190 (7)	0.0179 (8)	0.0187 (7)	-0.0015 (6)	0.0054 (5)	0.0013 (5)
C4B	0.0217 (7)	0.0197 (8)	0.0197 (7)	0.0027 (6)	0.0068 (5)	-0.0009 (6)
C5B	0.0212 (7)	0.0182 (8)	0.0202 (7)	0.0034 (6)	0.0059 (5)	0.0004 (6)
C6B	0.0182 (7)	0.0150 (7)	0.0187 (6)	0.0006 (5)	0.0049 (5)	-0.0002 (5)
C7B	0.0175 (7)	0.0163 (7)	0.0203 (7)	-0.0020 (5)	0.0056 (5)	-0.0009 (5)
C8B	0.0220 (7)	0.0203 (8)	0.0215 (7)	0.0021 (6)	0.0044 (6)	-0.0007 (6)
C9B	0.0217 (8)	0.0230 (9)	0.0252 (8)	0.0004 (6)	0.0035 (6)	0.0007 (6)
C10B	0.0239 (8)	0.0230 (9)	0.0222 (7)	-0.0036 (6)	0.0036 (6)	0.0003 (6)
C11B	0.0228 (8)	0.0219 (8)	0.0226 (7)	-0.0016 (6)	0.0045 (6)	-0.0010 (6)
C12B	0.0199 (7)	0.0194 (8)	0.0206 (7)	0.0003 (6)	0.0038 (5)	-0.0036 (6)
C13B	0.0247 (8)	0.0241 (9)	0.0220 (7)	-0.0011 (7)	0.0015 (6)	-0.0041 (6)
C14B	0.0267 (8)	0.0275 (9)	0.0201 (7)	-0.0011 (7)	0.0022 (6)	-0.0012 (6)
C15B	0.0313 (9)	0.0242 (9)	0.0258 (8)	-0.0026 (7)	-0.0007 (7)	-0.0028 (7)
C16B	0.0277 (9)	0.0316 (10)	0.0243 (8)	0.0048 (7)	-0.0005 (7)	-0.0104 (7)
C17B	0.0301 (10)	0.0517 (15)	0.0371 (11)	-0.0113 (10)	0.0124 (8)	-0.0192 (10)
C18B	0.0352 (10)	0.0386 (12)	0.0375 (10)	-0.0153 (9)	0.0137 (8)	-0.0166 (9)
C19B	0.0285 (9)	0.0288 (10)	0.0238 (8)	-0.0024 (7)	-0.0002 (7)	0.0012 (7)
C20B	0.0416 (6)	0.0488 (7)	0.0299 (5)	-0.0111 (5)	-0.0030 (4)	0.0040 (5)
C21B	0.0416 (6)	0.0488 (7)	0.0299 (5)	-0.0111 (5)	-0.0030 (4)	0.0040 (5)
C22B	0.0196 (8)	0.0363 (11)	0.0250 (8)	0.0041 (7)	0.0041 (6)	0.0058 (7)

Geometric parameters (\AA , $^\circ$)

Cl1A—C16A	1.7412 (18)	Cl1B—C16B	1.7435 (19)
O1A—C3A	1.358 (2)	O1B—C3B	1.361 (2)
O1A—C22A	1.429 (2)	O1B—C22B	1.431 (2)
O2A—C9A	1.226 (2)	O2B—C9B	1.227 (2)
O3A—C19A	1.198 (2)	O3B—C19B	1.196 (3)

O4A—C19A	1.330 (2)	O4B—C19B	1.331 (2)
O4A—C20A	1.448 (2)	O4B—C20B	1.472 (3)
C1A—C6A	1.397 (2)	C1B—C2B	1.396 (2)
C1A—C2A	1.399 (2)	C1B—C6B	1.397 (2)
C1A—H1	0.9300	C1B—H6	0.9300
C2A—C3A	1.391 (2)	C2B—C3B	1.392 (2)
C2A—H2	0.9300	C2B—H7	0.9300
C3A—C4A	1.398 (2)	C3B—C4B	1.399 (2)
C4A—C5A	1.379 (2)	C4B—C5B	1.379 (2)
C4A—H3	0.9300	C4B—H8	0.9300
C5A—C6A	1.410 (2)	C5B—C6B	1.413 (2)
C5A—H4	0.9300	C5B—H9	0.9300
C6A—C7A	1.477 (2)	C6B—C7B	1.475 (2)
C7A—C8A	1.355 (2)	C7B—C8B	1.354 (2)
C7A—C12A	1.510 (2)	C7B—C12B	1.509 (2)
C8A—C9A	1.457 (2)	C8B—C9B	1.457 (2)
C8A—H5	0.9300	C8B—H10	0.9300
C9A—C10A	1.524 (2)	C9B—C10B	1.529 (3)
C10A—C19A	1.520 (3)	C10B—C19B	1.519 (3)
C10A—C11A	1.537 (3)	C10B—C11B	1.534 (3)
C10A—H10A	0.9800	C10B—H10B	0.9800
C11A—C13A	1.519 (2)	C11B—C13B	1.518 (3)
C11A—C12A	1.525 (2)	C11B—C12B	1.533 (2)
C11A—H11A	0.9800	C11B—H11B	0.9800
C12A—H12A	0.9700	C12B—H12C	0.9700
C12A—H12B	0.9700	C12B—H12D	0.9700
C13A—C14A	1.389 (3)	C13B—C14B	1.387 (3)
C13A—C18A	1.400 (3)	C13B—C18B	1.396 (3)
C14A—C15A	1.394 (2)	C14B—C15B	1.389 (3)
C14A—H14A	0.9300	C14B—H14B	0.9300
C15A—C16A	1.380 (2)	C15B—C16B	1.379 (3)
C15A—H15A	0.9300	C15B—H15B	0.9300
C16A—C17A	1.389 (3)	C16B—C17B	1.381 (3)
C17A—C18A	1.390 (3)	C17B—C18B	1.391 (3)
C17A—H17A	0.9300	C17B—H17B	0.9300
C18A—H18A	0.9300	C18B—H18B	0.9300
C20A—C21A	1.499 (3)	C20B—C21B	1.470 (3)
C20A—H20A	0.9700	C20B—H20C	0.9700
C20A—H20B	0.9700	C20B—H20D	0.9700
C21A—H21A	0.9600	C21B—H21D	0.9600
C21A—H21B	0.9600	C21B—H21E	0.9600
C21A—H21C	0.9600	C21B—H21F	0.9600
C22A—H22A	0.9600	C22B—H22D	0.9600
C22A—H22B	0.9600	C22B—H22E	0.9600
C22A—H22C	0.9600	C22B—H22F	0.9600
C3A—O1A—C22A	117.53 (14)	C3B—O1B—C22B	117.52 (14)
C19A—O4A—C20A	117.88 (16)	C19B—O4B—C20B	115.04 (17)
C6A—C1A—C2A	121.93 (15)	C2B—C1B—C6B	122.08 (15)
C6A—C1A—H1	119.0	C2B—C1B—H6	119.0

supplementary materials

C2A—C1A—H1	119.0	C6B—C1B—H6	119.0
C3A—C2A—C1A	119.54 (16)	C3B—C2B—C1B	119.46 (16)
C3A—C2A—H2	120.2	C3B—C2B—H7	120.3
C1A—C2A—H2	120.2	C1B—C2B—H7	120.3
O1A—C3A—C2A	125.56 (15)	O1B—C3B—C2B	125.33 (15)
O1A—C3A—C4A	114.96 (14)	O1B—C3B—C4B	115.09 (15)
C2A—C3A—C4A	119.47 (15)	C2B—C3B—C4B	119.58 (15)
C5A—C4A—C3A	120.34 (15)	C5B—C4B—C3B	120.34 (15)
C5A—C4A—H3	119.8	C5B—C4B—H8	119.8
C3A—C4A—H3	119.8	C3B—C4B—H8	119.8
C4A—C5A—C6A	121.60 (16)	C4B—C5B—C6B	121.43 (15)
C4A—C5A—H4	119.2	C4B—C5B—H9	119.3
C6A—C5A—H4	119.2	C6B—C5B—H9	119.3
C1A—C6A—C5A	117.10 (15)	C1B—C6B—C5B	117.08 (15)
C1A—C6A—C7A	121.23 (14)	C1B—C6B—C7B	120.96 (14)
C5A—C6A—C7A	121.62 (15)	C5B—C6B—C7B	121.92 (15)
C8A—C7A—C6A	121.44 (15)	C8B—C7B—C6B	122.04 (15)
C8A—C7A—C12A	120.09 (15)	C8B—C7B—C12B	119.66 (15)
C6A—C7A—C12A	118.42 (14)	C6B—C7B—C12B	118.25 (14)
C7A—C8A—C9A	123.66 (15)	C7B—C8B—C9B	123.75 (16)
C7A—C8A—H5	118.2	C7B—C8B—H10	118.1
C9A—C8A—H5	118.2	C9B—C8B—H10	118.1
O2A—C9A—C8A	121.92 (16)	O2B—C9B—C8B	121.78 (17)
O2A—C9A—C10A	121.42 (15)	O2B—C9B—C10B	120.58 (16)
C8A—C9A—C10A	116.49 (15)	C8B—C9B—C10B	117.54 (16)
C19A—C10A—C9A	108.61 (15)	C19B—C10B—C9B	107.99 (15)
C19A—C10A—C11A	111.23 (15)	C19B—C10B—C11B	110.80 (15)
C9A—C10A—C11A	109.63 (14)	C9B—C10B—C11B	109.50 (14)
C19A—C10A—H10A	109.1	C19B—C10B—H10B	109.5
C9A—C10A—H10A	109.1	C9B—C10B—H10B	109.5
C11A—C10A—H10A	109.1	C11B—C10B—H10B	109.5
C13A—C11A—C12A	113.23 (15)	C13B—C11B—C12B	111.51 (15)
C13A—C11A—C10A	112.21 (14)	C13B—C11B—C10B	112.82 (15)
C12A—C11A—C10A	109.22 (14)	C12B—C11B—C10B	109.90 (15)
C13A—C11A—H11A	107.3	C13B—C11B—H11B	107.5
C12A—C11A—H11A	107.3	C12B—C11B—H11B	107.5
C10A—C11A—H11A	107.3	C10B—C11B—H11B	107.5
C7A—C12A—C11A	112.34 (14)	C7B—C12B—C11B	112.52 (14)
C7A—C12A—H12A	109.1	C7B—C12B—H12C	109.1
C11A—C12A—H12A	109.1	C11B—C12B—H12C	109.1
C7A—C12A—H12B	109.1	C7B—C12B—H12D	109.1
C11A—C12A—H12B	109.1	C11B—C12B—H12D	109.1
H12A—C12A—H12B	107.9	H12C—C12B—H12D	107.8
C14A—C13A—C18A	118.50 (16)	C14B—C13B—C18B	118.69 (17)
C14A—C13A—C11A	118.87 (16)	C14B—C13B—C11B	119.00 (16)
C18A—C13A—C11A	122.63 (16)	C18B—C13B—C11B	122.30 (17)
C13A—C14A—C15A	121.00 (17)	C13B—C14B—C15B	120.91 (18)
C13A—C14A—H14A	119.5	C13B—C14B—H14B	119.5
C15A—C14A—H14A	119.5	C15B—C14B—H14B	119.5

C16A—C15A—C14A	119.22 (17)	C16B—C15B—C14B	119.28 (19)
C16A—C15A—H15A	120.4	C16B—C15B—H15B	120.4
C14A—C15A—H15A	120.4	C14B—C15B—H15B	120.4
C15A—C16A—C17A	121.35 (16)	C15B—C16B—C17B	121.28 (18)
C15A—C16A—Cl1A	119.08 (14)	C15B—C16B—Cl1B	119.37 (16)
C17A—C16A—Cl1A	119.57 (14)	C17B—C16B—Cl1B	119.35 (16)
C16A—C17A—C18A	118.69 (17)	C16B—C17B—C18B	119.0 (2)
C16A—C17A—H17A	120.7	C16B—C17B—H17B	120.5
C18A—C17A—H17A	120.7	C18B—C17B—H17B	120.5
C17A—C18A—C13A	121.22 (17)	C17B—C18B—C13B	120.9 (2)
C17A—C18A—H18A	119.4	C17B—C18B—H18B	119.6
C13A—C18A—H18A	119.4	C13B—C18B—H18B	119.6
O3A—C19A—O4A	125.08 (17)	O3B—C19B—O4B	123.55 (18)
O3A—C19A—C10A	124.77 (17)	O3B—C19B—C10B	124.74 (18)
O4A—C19A—C10A	110.14 (16)	O4B—C19B—C10B	111.70 (17)
O4A—C20A—C21A	108.05 (18)	C21B—C20B—O4B	107.91 (19)
O4A—C20A—H20A	110.1	C21B—C20B—H20C	110.1
C21A—C20A—H20A	110.1	O4B—C20B—H20C	110.1
O4A—C20A—H20B	110.1	C21B—C20B—H20D	110.1
C21A—C20A—H20B	110.1	O4B—C20B—H20D	110.1
H20A—C20A—H20B	108.4	H20C—C20B—H20D	108.4
C20A—C21A—H21A	109.5	C20B—C21B—H21D	109.5
C20A—C21A—H21B	109.5	C20B—C21B—H21E	109.5
H21A—C21A—H21B	109.5	H21D—C21B—H21E	109.5
C20A—C21A—H21C	109.5	C20B—C21B—H21F	109.5
H21A—C21A—H21C	109.5	H21D—C21B—H21F	109.5
H21B—C21A—H21C	109.5	H21E—C21B—H21F	109.5
O1A—C22A—H22A	109.5	O1B—C22B—H22D	109.5
O1A—C22A—H22B	109.5	O1B—C22B—H22E	109.5
H22A—C22A—H22B	109.5	H22D—C22B—H22E	109.5
O1A—C22A—H22C	109.5	O1B—C22B—H22F	109.5
H22A—C22A—H22C	109.5	H22D—C22B—H22F	109.5
H22B—C22A—H22C	109.5	H22E—C22B—H22F	109.5
C6A—C1A—C2A—C3A	-0.4 (3)	C6B—C1B—C2B—C3B	1.2 (3)
C22A—O1A—C3A—C2A	-5.5 (2)	C22B—O1B—C3B—C2B	8.7 (2)
C22A—O1A—C3A—C4A	174.43 (15)	C22B—O1B—C3B—C4B	-171.47 (15)
C1A—C2A—C3A—O1A	-178.75 (16)	C1B—C2B—C3B—O1B	178.15 (16)
C1A—C2A—C3A—C4A	1.3 (3)	C1B—C2B—C3B—C4B	-1.6 (3)
O1A—C3A—C4A—C5A	179.41 (16)	O1B—C3B—C4B—C5B	-179.46 (16)
C2A—C3A—C4A—C5A	-0.6 (3)	C2B—C3B—C4B—C5B	0.4 (3)
C3A—C4A—C5A—C6A	-0.9 (3)	C3B—C4B—C5B—C6B	1.4 (3)
C2A—C1A—C6A—C5A	-1.0 (3)	C2B—C1B—C6B—C5B	0.5 (3)
C2A—C1A—C6A—C7A	176.27 (16)	C2B—C1B—C6B—C7B	-177.18 (16)
C4A—C5A—C6A—C1A	1.7 (3)	C4B—C5B—C6B—C1B	-1.8 (3)
C4A—C5A—C6A—C7A	-175.58 (16)	C4B—C5B—C6B—C7B	175.83 (16)
C1A—C6A—C7A—C8A	-165.71 (16)	C1B—C6B—C7B—C8B	167.12 (17)
C5A—C6A—C7A—C8A	11.5 (2)	C5B—C6B—C7B—C8B	-10.5 (3)
C1A—C6A—C7A—C12A	11.8 (2)	C1B—C6B—C7B—C12B	-10.4 (2)
C5A—C6A—C7A—C12A	-171.04 (15)	C5B—C6B—C7B—C12B	172.03 (15)

supplementary materials


C6A—C7A—C8A—C9A	171.75 (16)	C6B—C7B—C8B—C9B	−171.91 (16)
C12A—C7A—C8A—C9A	−5.7 (3)	C12B—C7B—C8B—C9B	5.5 (3)
C7A—C8A—C9A—O2A	−177.50 (18)	C7B—C8B—C9B—O2B	176.40 (18)
C7A—C8A—C9A—C10A	−2.1 (3)	C7B—C8B—C9B—C10B	−0.2 (3)
O2A—C9A—C10A—C19A	−28.0 (2)	O2B—C9B—C10B—C19B	31.1 (2)
C8A—C9A—C10A—C19A	156.65 (16)	C8B—C9B—C10B—C19B	−152.28 (17)
O2A—C9A—C10A—C11A	−149.68 (18)	O2B—C9B—C10B—C11B	151.80 (18)
C8A—C9A—C10A—C11A	34.9 (2)	C8B—C9B—C10B—C11B	−31.5 (2)
C19A—C10A—C11A—C13A	53.8 (2)	C19B—C10B—C11B—C13B	−58.7 (2)
C9A—C10A—C11A—C13A	173.98 (15)	C9B—C10B—C11B—C13B	−177.67 (15)
C19A—C10A—C11A—C12A	−179.76 (14)	C19B—C10B—C11B—C12B	176.23 (14)
C9A—C10A—C11A—C12A	−59.61 (18)	C9B—C10B—C11B—C12B	57.21 (18)
C8A—C7A—C12A—C11A	−20.7 (2)	C8B—C7B—C12B—C11B	21.9 (2)
C6A—C7A—C12A—C11A	161.77 (14)	C6B—C7B—C12B—C11B	−160.50 (15)
C13A—C11A—C12A—C7A	178.76 (14)	C13B—C11B—C12B—C7B	−179.27 (15)
C10A—C11A—C12A—C7A	52.94 (18)	C10B—C11B—C12B—C7B	−53.39 (19)
C12A—C11A—C13A—C14A	124.12 (18)	C12B—C11B—C13B—C14B	−119.08 (18)
C10A—C11A—C13A—C14A	−111.67 (19)	C10B—C11B—C13B—C14B	116.67 (19)
C12A—C11A—C13A—C18A	−56.1 (2)	C12B—C11B—C13B—C18B	59.9 (2)
C10A—C11A—C13A—C18A	68.1 (2)	C10B—C11B—C13B—C18B	−64.3 (2)
C18A—C13A—C14A—C15A	−1.0 (3)	C18B—C13B—C14B—C15B	−0.2 (3)
C11A—C13A—C14A—C15A	178.80 (16)	C11B—C13B—C14B—C15B	178.82 (17)
C13A—C14A—C15A—C16A	−0.6 (3)	C13B—C14B—C15B—C16B	0.7 (3)
C14A—C15A—C16A—C17A	1.6 (3)	C14B—C15B—C16B—C17B	−1.0 (3)
C14A—C15A—C16A—C11A	−177.76 (14)	C14B—C15B—C16B—C11B	177.82 (15)
C15A—C16A—C17A—C18A	−1.1 (3)	C15B—C16B—C17B—C18B	0.9 (4)
C11A—C16A—C17A—C18A	178.31 (16)	C11B—C16B—C17B—C18B	−178.01 (19)
C16A—C17A—C18A—C13A	−0.5 (3)	C16B—C17B—C18B—C13B	−0.3 (4)
C14A—C13A—C18A—C17A	1.5 (3)	C14B—C13B—C18B—C17B	0.0 (3)
C11A—C13A—C18A—C17A	−178.23 (18)	C11B—C13B—C18B—C17B	−179.0 (2)
C20A—O4A—C19A—O3A	−7.5 (3)	C20B—O4B—C19B—O3B	2.9 (3)
C20A—O4A—C19A—C10A	171.64 (17)	C20B—O4B—C19B—C10B	−178.08 (19)
C9A—C10A—C19A—O3A	−65.2 (3)	C9B—C10B—C19B—O3B	68.3 (3)
C11A—C10A—C19A—O3A	55.6 (3)	C11B—C10B—C19B—O3B	−51.6 (3)
C9A—C10A—C19A—O4A	115.70 (17)	C9B—C10B—C19B—O4B	−110.75 (19)
C11A—C10A—C19A—O4A	−123.55 (17)	C11B—C10B—C19B—O4B	129.32 (18)
C19A—O4A—C20A—C21A	−105.0 (2)	C19B—O4B—C20B—C21B	177.3 (2)

Hydrogen-bond geometry (\AA , $^\circ$)

$D\cdots H\cdots A$	$D\cdots H$	$H\cdots A$	$D\cdots A$	$D\cdots H\cdots A$
C11A—H11A—O1B ⁱ	0.98	2.53	3.492 (2)	167
C11B—H11B—O1A ⁱⁱ	0.98	2.53	3.501 (2)	170
C12A—H12B—O2B	0.97	2.51	3.450 (2)	162
C12B—H12C—O2A ⁱⁱⁱ	0.97	2.56	3.441 (2)	151
C15B—H15B—O4B ^{iv}	0.93	2.59	3.485 (3)	163
C20B—H20D—C11A ^{iv}	0.97	2.83	3.585 (2)	136
C22A—H22A—Cg1 ⁱⁱ	0.97	2.83	3.666 (2)	146

Symmetry codes: (i) $-x+1, y-1/2, -z+3/2$; (ii) $-x+1, y+1/2, -z+3/2$; (iii) $x-1, y, z$; (iv) $x, y+1, z$.

Fig. 1



supplementary materials

Fig. 2

