

2,2-Bis(1*H*-indol-3-yl)indolin-3-one

Zhao-Hao Li,^{a,b} Jing Xu,^{a,b} Wen-Liang Wu^{a,b} and Wei-Ping Su^{b*}

^aGraduate School, The Chinese Academy of Sciences, Beijing 100039, People's Republic of China, and ^bThe State Key Laboratory of Structural Chemistry, Fujian Institute of Research on the Structure of Matter, The Chinese Academy of Sciences, Fuzhou, Fujian 350002, People's Republic of China
Correspondence e-mail: xujing@fjirsm.ac.cn


Received 22 April 2009; accepted 11 May 2009

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$; R factor = 0.048; wR factor = 0.118; data-to-parameter ratio = 16.8.

In the title molecule, $\text{C}_{24}\text{H}_{17}\text{N}_3\text{O}$, the mean plane of the indolone ring forms dihedral angles of $112.0(1)$ and $103.1(1)^\circ$ with the planes of the two indole rings. The dihedral angle between the mean planes of the two indole rings is $63.5(1)^\circ$. In the crystal structure, molecules are linked via intermolecular $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds, forming a two-dimensional network parallel to the ab plane.

Related literature

For the applications of indole derivatives, see: Ramesh *et al.* (2009). For the isolation of the title compound as a natural product, see: Ganachaud *et al.* (2008); Stull *et al.* (1995).



Experimental

Crystal data

$\text{C}_{24}\text{H}_{17}\text{N}_3\text{O}$

$M_r = 363.41$

Monoclinic, $P2_1/n$
 $a = 10.559(4)\text{ \AA}$
 $b = 8.931(3)\text{ \AA}$
 $c = 19.899(7)\text{ \AA}$
 $\beta = 98.480(6)^\circ$
 $V = 1856.1(11)\text{ \AA}^3$

$Z = 4$
Mo $K\alpha$ radiation
 $\mu = 0.08\text{ mm}^{-1}$
 $T = 293\text{ K}$
 $0.40 \times 0.35 \times 0.15\text{ mm}$

Data collection

Rigaku Mercury CCD diffractometer
Absorption correction: multi-scan (*CrystalClear*; Rigaku/MSC, 2005)
 $T_{\min} = 0.968$, $T_{\max} = 0.988$

14004 measured reflections
4245 independent reflections
3606 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.024$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.048$
 $wR(F^2) = 0.118$
 $S = 1.06$
4245 reflections

253 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.21\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.19\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N2—H2B \cdots O1 ⁱ	0.86	2.12	2.9412 (17)	159
N3—H3B \cdots O1 ⁱⁱ	0.86	2.18	2.9830 (16)	156

Symmetry codes: (i) $-x + \frac{3}{2}, y - \frac{1}{2}, -z + \frac{1}{2}$, (ii) $-x + \frac{5}{2}, y - \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: *CrystalClear* (Rigaku/MSC, 2005); cell refinement: *CrystalClear*; data reduction: *CrystalClear*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *DIAMOND* (Brandenburg, 1999); software used to prepare material for publication: *SHELXL97*.

This work was supported by the Natural Science Foundation of Fujian province (grant No. 2006 J0273) and the 'One Hundred Talent' Projects from the Chinese Academy of Sciences.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: LH2811).

References

- Brandenburg, K. (1999). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- Ganachaud, C., Garagnani, V., Tron, T. & Iacazio, G. (2008). *Tetrahedron Lett.* **49**, 2476–2478.
- Ramesh, P., Sundaresan, S. S., Lakshmi, N. V., Perumal, P. T. & Ponnuswamy, M. N. (2009). *Acta Cryst. E* **65**, o994.
- Rigaku/MSC (2005). *CrystalClear*. Rigaku/MSC, The Woodlands, Texas, USA.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Stull, T. L., Hyun, L., Sharetsky, C., Wooten, J., McCauley, J. P. & Smith, A. B. III (1995). *J. Biol. Chem.* **270**, 5–8.

supplementary materials

Acta Cryst. (2009). E65, o1353 [doi:10.1107/S1600536809017553]

2,2-Bis(1*H*-indol-3-yl)indolin-3-one

Z.-H. Li, J. Xu, W.-L. Wu and W.-P. Su

Comment

Indole derivatives are used as bioactive drugs and they exhibit anti-allergic, central nervous system depressant and muscle relaxant properties (Ramesh *et al.*, 2009). The title compound is a natural product that has been isolated from bacterial sources (Stull *et al.*, 1995; Ganachaud *et al.*, 2008). Recently, we found different indole derivatives could be formed by oxidation solely on the basis of the reaction solvent and temperature. As part of our studies, we report herein the synthesis and crystal structure of the title compound (*I*) (Fig. 1).

The title compound is a trimeric condensation product of indole through the formation of a quaternary carbon (C8) centre. The indolone ring forms dihedral angle of 112.0 (1) $^{\circ}$ and 103.1 (1) $^{\circ}$, respectively, with the two indole rings (C9/C10/C11/C12/C13/C14/C15/C16/N2) and (C17/C18/C19/C20/C21/C22/C23/C24/N3). The mean planes of the two indole form a dihedral angle of 63.5 (1) $^{\circ}$. In the crystal structure, the carbonyl atom O1 acts as a bifurcated acceptor for the N-H groups of atoms N2 and N3 to form a two-dimensional network parallel to the *ab* plane (Table 1, Fig. 2).

Experimental

The title compound was obtained from a mixture of 1*H*-indole (78 mg) with *t*BuCOOH (0.18 ml) and H₃PMo₁₂O₄₀ (9 mg) in toluene (1 ml) and CH₃COOH (1 ml) under a nitrogen atmosphere at room temperature for 24 h. The crude product was isolated and purified by silica gel column chromatography. Yellow prism-shaped crystals of (*I*) suitable for X-ray diffraction were grown by slow evaporation of a dichloromethane solution at room temperature.

Refinement

H atoms were positioned geometrically and allowed to ride on their parent atoms, with C—H = 0.93 Å; N—H = 0.86 Å and *U*_{iso}(H) = 1.2*U*_{eq}(C,N).

Figures




Fig. 1. The molecular structure of (*I*), showing 30% probability displacement ellipsoids. H atoms are shown as small spheres.

supplementary materials




Fig. 2. Part of the crystal structure of (I) showing the hydrogen bonded layers parallel to the *ab* plane. Intermolecular hydrogen bonds are shown as dashed lines and H atoms not involved in H-bonding have been omitted.

2,2-Bis(1*H*-indol-3-yl)indolin-3-one

Crystal data

C ₂₄ H ₁₇ N ₃ O	$F_{000} = 760$
$M_r = 363.41$	$D_x = 1.300 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/n$	Mo $K\alpha$ radiation
Hall symbol: -P 2yn	$\lambda = 0.71073 \text{ \AA}$
$a = 10.559 (4) \text{ \AA}$	Cell parameters from 4234 reflections
$b = 8.931 (3) \text{ \AA}$	$\theta = 3.0\text{--}27.5^\circ$
$c = 19.899 (7) \text{ \AA}$	$\mu = 0.08 \text{ mm}^{-1}$
$\beta = 98.480 (6)^\circ$	$T = 293 \text{ K}$
$V = 1856.1 (11) \text{ \AA}^3$	Prism, yellow
$Z = 4$	$0.40 \times 0.35 \times 0.15 \text{ mm}$

Data collection

Rigaku Mercury CCD diffractometer	4245 independent reflections
Radiation source: fine-focus sealed tube	3606 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.024$
Detector resolution: 14.6306 pixels mm ⁻¹	$\theta_{\text{max}} = 27.5^\circ$
$T = 293 \text{ K}$	$\theta_{\text{min}} = 2.1^\circ$
CCD_Profile_fitting scans	$h = -11\text{--}13$
Absorption correction: multi-scan (CrystalClear; Rigaku/MSC, 2005)	$k = -11\text{--}11$
$T_{\text{min}} = 0.968, T_{\text{max}} = 0.988$	$l = -22\text{--}25$
14004 measured reflections	

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.048$	H-atom parameters constrained
$wR(F^2) = 0.118$	$w = 1/[\sigma^2(F_o^2) + (0.0513P)^2 + 0.4603P]$
$S = 1.06$	where $P = (F_o^2 + 2F_c^2)/3$
4245 reflections	$(\Delta/\sigma)_{\text{max}} < 0.001$
253 parameters	$\Delta\rho_{\text{max}} = 0.21 \text{ e \AA}^{-3}$
	$\Delta\rho_{\text{min}} = -0.19 \text{ e \AA}^{-3}$

Primary atom site location: structure-invariant direct Extinction correction: none
methods

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.96565 (9)	0.43252 (10)	0.15911 (5)	0.0369 (2)
N1	0.95403 (12)	0.04726 (13)	0.11858 (6)	0.0406 (3)
H1A	0.9668	-0.0474	0.1239	0.049*
N2	0.75509 (12)	0.05077 (15)	0.28328 (7)	0.0490 (3)
H2B	0.6914	-0.0017	0.2923	0.059*
N3	1.31856 (12)	0.10180 (15)	0.26178 (6)	0.0471 (3)
H3B	1.3724	0.0659	0.2945	0.056*
C1	0.89773 (13)	0.11138 (16)	0.05986 (7)	0.0374 (3)
C2	0.85094 (14)	0.0420 (2)	-0.00189 (8)	0.0483 (4)
H2A	0.8532	-0.0615	-0.0065	0.058*
C3	0.80181 (17)	0.1314 (2)	-0.05518 (8)	0.0635 (5)
H3A	0.7706	0.0868	-0.0966	0.076*
C4	0.7967 (2)	0.2867 (3)	-0.04991 (9)	0.0746 (6)
H4A	0.7630	0.3437	-0.0874	0.090*
C5	0.84151 (19)	0.3558 (2)	0.01064 (8)	0.0604 (5)
H5A	0.8383	0.4595	0.0147	0.072*
C6	0.89185 (13)	0.26719 (17)	0.06585 (7)	0.0392 (3)
C7	0.94976 (12)	0.30732 (14)	0.13383 (6)	0.0315 (3)
C8	0.99057 (13)	0.15946 (14)	0.17220 (6)	0.0321 (3)
C9	0.81798 (14)	0.15783 (17)	0.32497 (7)	0.0417 (3)
C10	0.79721 (18)	0.2065 (2)	0.38885 (8)	0.0557 (4)
H10A	0.7324	0.1655	0.4100	0.067*
C11	0.8753 (2)	0.3166 (2)	0.41938 (8)	0.0616 (5)
H11A	0.8635	0.3509	0.4621	0.074*
C12	0.97253 (17)	0.3786 (2)	0.38756 (8)	0.0550 (4)
H12A	1.0236	0.4541	0.4093	0.066*
C13	0.99417 (15)	0.32982 (17)	0.32449 (7)	0.0424 (3)
H13A	1.0597	0.3712	0.3040	0.051*
C14	0.91605 (13)	0.21709 (15)	0.29179 (7)	0.0351 (3)
C15	0.90865 (13)	0.13904 (15)	0.22798 (7)	0.0342 (3)
C16	0.80919 (14)	0.04053 (17)	0.22537 (8)	0.0432 (3)

supplementary materials

H16A	0.7823	-0.0240	0.1894	0.052*
C17	1.34968 (13)	0.18478 (15)	0.20847 (7)	0.0380 (3)
C18	1.46884 (15)	0.23042 (18)	0.19416 (9)	0.0491 (4)
H18A	1.5436	0.2065	0.2230	0.059*
C19	1.47219 (15)	0.31203 (18)	0.13592 (9)	0.0511 (4)
H19A	1.5507	0.3436	0.1251	0.061*
C20	1.36003 (15)	0.34840 (18)	0.09265 (8)	0.0461 (4)
H20A	1.3651	0.4030	0.0533	0.055*
C21	1.24194 (14)	0.30469 (15)	0.10722 (7)	0.0380 (3)
H21A	1.1679	0.3304	0.0782	0.046*
C22	1.23417 (12)	0.22113 (14)	0.16619 (6)	0.0322 (3)
C23	1.13200 (12)	0.15690 (14)	0.19763 (6)	0.0326 (3)
C24	1.18882 (14)	0.08515 (16)	0.25473 (7)	0.0412 (3)
H24A	1.1454	0.0325	0.2846	0.049*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0351 (5)	0.0344 (5)	0.0407 (5)	0.0007 (4)	0.0040 (4)	-0.0052 (4)
N1	0.0497 (7)	0.0320 (6)	0.0391 (6)	-0.0014 (5)	0.0029 (5)	-0.0058 (5)
N2	0.0361 (7)	0.0527 (8)	0.0603 (8)	-0.0086 (6)	0.0138 (6)	0.0105 (6)
N3	0.0361 (7)	0.0560 (8)	0.0472 (7)	0.0108 (6)	-0.0002 (5)	0.0156 (6)
C1	0.0288 (7)	0.0464 (8)	0.0367 (7)	-0.0010 (6)	0.0041 (5)	-0.0075 (6)
C2	0.0377 (8)	0.0583 (9)	0.0474 (8)	-0.0003 (7)	0.0016 (6)	-0.0203 (7)
C3	0.0549 (11)	0.0890 (14)	0.0415 (9)	0.0140 (10)	-0.0100 (8)	-0.0235 (9)
C4	0.0916 (16)	0.0844 (14)	0.0395 (9)	0.0287 (12)	-0.0181 (9)	-0.0053 (9)
C5	0.0750 (12)	0.0580 (10)	0.0423 (8)	0.0195 (9)	-0.0110 (8)	-0.0010 (7)
C6	0.0353 (7)	0.0455 (8)	0.0347 (7)	0.0055 (6)	-0.0015 (6)	-0.0053 (6)
C7	0.0252 (6)	0.0364 (7)	0.0330 (6)	0.0018 (5)	0.0048 (5)	-0.0025 (5)
C8	0.0327 (7)	0.0322 (6)	0.0311 (6)	-0.0015 (5)	0.0034 (5)	-0.0032 (5)
C9	0.0367 (8)	0.0463 (8)	0.0437 (8)	0.0066 (6)	0.0114 (6)	0.0125 (6)
C10	0.0577 (10)	0.0660 (11)	0.0486 (9)	0.0120 (9)	0.0249 (8)	0.0170 (8)
C11	0.0735 (13)	0.0775 (13)	0.0365 (8)	0.0178 (10)	0.0174 (8)	0.0021 (8)
C12	0.0601 (11)	0.0623 (10)	0.0419 (8)	0.0047 (8)	0.0051 (7)	-0.0090 (7)
C13	0.0425 (8)	0.0485 (8)	0.0366 (7)	-0.0009 (6)	0.0067 (6)	-0.0014 (6)
C14	0.0327 (7)	0.0391 (7)	0.0338 (6)	0.0038 (5)	0.0058 (5)	0.0054 (5)
C15	0.0309 (7)	0.0353 (7)	0.0359 (7)	-0.0004 (5)	0.0039 (5)	0.0026 (5)
C16	0.0367 (8)	0.0436 (8)	0.0486 (8)	-0.0051 (6)	0.0044 (6)	0.0023 (6)
C17	0.0338 (7)	0.0366 (7)	0.0430 (7)	0.0063 (5)	0.0037 (6)	0.0013 (6)
C18	0.0297 (7)	0.0524 (9)	0.0640 (10)	0.0058 (6)	0.0030 (7)	0.0001 (8)
C19	0.0359 (8)	0.0515 (9)	0.0686 (11)	-0.0027 (7)	0.0172 (7)	-0.0014 (8)
C20	0.0472 (9)	0.0473 (8)	0.0463 (8)	-0.0034 (7)	0.0154 (7)	0.0031 (7)
C21	0.0371 (7)	0.0407 (7)	0.0359 (7)	0.0006 (6)	0.0049 (6)	0.0006 (6)
C22	0.0313 (7)	0.0308 (6)	0.0346 (6)	0.0036 (5)	0.0051 (5)	-0.0022 (5)
C23	0.0319 (7)	0.0327 (6)	0.0334 (6)	0.0029 (5)	0.0056 (5)	0.0001 (5)
C24	0.0376 (8)	0.0436 (8)	0.0428 (8)	0.0054 (6)	0.0070 (6)	0.0096 (6)

Geometric parameters (Å, °)

O1—C7	1.2274 (16)	C9—C14	1.411 (2)
N1—C1	1.3571 (18)	C10—C11	1.367 (3)
N1—C8	1.4731 (16)	C10—H10A	0.9300
N1—H1A	0.8600	C11—C12	1.398 (3)
N2—C16	1.363 (2)	C11—H11A	0.9300
N2—C9	1.371 (2)	C12—C13	1.379 (2)
N2—H2B	0.8600	C12—H12A	0.9300
N3—C24	1.3645 (19)	C13—C14	1.400 (2)
N3—C17	1.3731 (19)	C13—H13A	0.9300
N3—H3B	0.8600	C14—C15	1.4405 (19)
C1—C6	1.399 (2)	C15—C16	1.3652 (19)
C1—C2	1.3997 (19)	C16—H16A	0.9300
C2—C3	1.367 (3)	C17—C18	1.391 (2)
C2—H2A	0.9300	C17—C22	1.4136 (19)
C3—C4	1.392 (3)	C18—C19	1.374 (2)
C3—H3A	0.9300	C18—H18A	0.9300
C4—C5	1.374 (2)	C19—C20	1.396 (2)
C4—H4A	0.9300	C19—H19A	0.9300
C5—C6	1.394 (2)	C20—C21	1.378 (2)
C5—H5A	0.9300	C20—H20A	0.9300
C6—C7	1.4451 (18)	C21—C22	1.4031 (19)
C7—C8	1.5542 (18)	C21—H21A	0.9300
C8—C23	1.5046 (19)	C22—C23	1.4434 (18)
C8—C15	1.5155 (19)	C23—C24	1.3640 (19)
C9—C10	1.391 (2)	C24—H24A	0.9300
C1—N1—C8	111.79 (11)	C10—C11—C12	121.26 (16)
C1—N1—H1A	124.1	C10—C11—H11A	119.4
C8—N1—H1A	124.1	C12—C11—H11A	119.4
C16—N2—C9	109.42 (12)	C13—C12—C11	121.24 (17)
C16—N2—H2B	125.3	C13—C12—H12A	119.4
C9—N2—H2B	125.3	C11—C12—H12A	119.4
C24—N3—C17	109.29 (12)	C12—C13—C14	119.00 (15)
C24—N3—H3B	125.4	C12—C13—H13A	120.5
C17—N3—H3B	125.4	C14—C13—H13A	120.5
N1—C1—C6	111.45 (12)	C13—C14—C9	118.37 (13)
N1—C1—C2	128.48 (14)	C13—C14—C15	135.19 (13)
C6—C1—C2	120.06 (14)	C9—C14—C15	106.43 (13)
C3—C2—C1	117.83 (16)	C16—C15—C14	106.67 (12)
C3—C2—H2A	121.1	C16—C15—C8	124.66 (13)
C1—C2—H2A	121.1	C14—C15—C8	128.66 (12)
C2—C3—C4	122.53 (15)	N2—C16—C15	109.91 (14)
C2—C3—H3A	118.7	N2—C16—H16A	125.0
C4—C3—H3A	118.7	C15—C16—H16A	125.0
C5—C4—C3	120.11 (17)	N3—C17—C18	130.01 (13)
C5—C4—H4A	119.9	N3—C17—C22	107.46 (12)
C3—C4—H4A	119.9	C18—C17—C22	122.53 (14)

supplementary materials


C4—C5—C6	118.52 (17)	C19—C18—C17	117.69 (14)
C4—C5—H5A	120.7	C19—C18—H18A	121.2
C6—C5—H5A	120.7	C17—C18—H18A	121.2
C5—C6—C1	120.95 (14)	C18—C19—C20	121.24 (14)
C5—C6—C7	131.01 (14)	C18—C19—H19A	119.4
C1—C6—C7	107.97 (12)	C20—C19—H19A	119.4
O1—C7—C6	128.53 (13)	C21—C20—C19	121.07 (14)
O1—C7—C8	124.16 (12)	C21—C20—H20A	119.5
C6—C7—C8	107.31 (11)	C19—C20—H20A	119.5
N1—C8—C23	111.98 (11)	C20—C21—C22	119.54 (13)
N1—C8—C15	109.37 (11)	C20—C21—H21A	120.2
C23—C8—C15	113.41 (11)	C22—C21—H21A	120.2
N1—C8—C7	101.43 (10)	C21—C22—C17	117.92 (12)
C23—C8—C7	111.57 (10)	C21—C22—C23	135.52 (12)
C15—C8—C7	108.37 (11)	C17—C22—C23	106.56 (12)
N2—C9—C10	130.02 (15)	C24—C23—C22	106.43 (12)
N2—C9—C14	107.57 (13)	C24—C23—C8	125.45 (12)
C10—C9—C14	122.41 (15)	C22—C23—C8	128.05 (11)
C11—C10—C9	117.71 (15)	C23—C24—N3	110.25 (13)
C11—C10—H10A	121.1	C23—C24—H24A	124.9
C9—C10—H10A	121.1	N3—C24—H24A	124.9

Hydrogen-bond geometry (\AA , $^\circ$)

$D\text{—H}\cdots A$	$D\text{—H}$	$H\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
N2—H2B \cdots O1 ⁱ	0.86	2.12	2.9412 (17)	159
N3—H3B \cdots O1 ⁱⁱ	0.86	2.18	2.9830 (16)	156

Symmetry codes: (i) $-x+3/2, y-1/2, -z+1/2$; (ii) $-x+5/2, y-1/2, -z+1/2$.

Fig. 1



supplementary materials

Fig. 2

