

## N'-(2,4-Dimethoxybenzylidene)-3,4,5-trihydroxybenzohydrazide ethanol solvate

Abeer A. Alhadi,<sup>a</sup> Siti Munirah Saharin,<sup>a</sup> Hapipah Mohd Ali,<sup>a\*</sup> Ward T. Robinson<sup>a</sup> and Mahmood A. Abdulla<sup>b</sup>

<sup>a</sup>Department of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia,

and <sup>b</sup>Department of Molecular Medicine, University of Malaya, 50603 Kuala Lumpur, Malaysia

Correspondence e-mail: hapipah@um.edu.my


Received 30 April 2009; accepted 19 May 2009

Key indicators: single-crystal X-ray study;  $T = 100\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$ ;  $R$  factor = 0.044;  $wR$  factor = 0.121; data-to-parameter ratio = 20.5.

The title compound,  $\text{C}_{16}\text{H}_{16}\text{N}_2\text{O}_6\cdot\text{C}_2\text{H}_5\text{OH}$ , was synthesized from 3,4,5-trihydroxybenzoylhydrazide and 2,4-dimethoxybenzaldehyde in ethanol. The compound is not planar, with the two aromatic planes of the two aromatic rings twisted by  $15.6(1)^\circ$ . The hydroxy groups are involved in both intramolecular  $\text{O}-\text{H}\cdots\text{O}$  and intermolecular  $\text{O}-\text{H}\cdots\text{N}$  and  $\text{O}-\text{H}\cdots\text{O}$  hydrogen bonds and a  $\text{C}-\text{H}\cdots\text{O}$  interaction also occurs.

### Related literature

For related compounds, see Abdul Alhadi *et al.* (2009). For the parent *N'*-(2-hydroxybenzylidene)benzohydrazide, see Lyubchova *et al.* (1995).



### Experimental

#### Crystal data

$\text{C}_{16}\text{H}_{16}\text{N}_2\text{O}_6\cdot\text{C}_2\text{H}_6\text{O}$   
 $M_r = 378.38$   
Monoclinic,  $P2_1/n$

$a = 7.8347(1)\text{ \AA}$   
 $b = 17.5412(3)\text{ \AA}$   
 $c = 13.0230(2)\text{ \AA}$

$\beta = 93.936(1)^\circ$   
 $V = 1785.53(5)\text{ \AA}^3$   
 $Z = 4$   
Mo  $K\alpha$  radiation

$\mu = 0.11\text{ mm}^{-1}$   
 $T = 100\text{ K}$   
 $0.31 \times 0.16 \times 0.12\text{ mm}$

#### Data collection

Bruker APEXII CCD area-detector diffractometer  
Absorption correction: multi-scan (*SADABS*; Sheldrick, 1996)  
 $T_{\min} = 0.967$ ,  $T_{\max} = 0.987$

19580 measured reflections  
5188 independent reflections  
3586 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.041$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.044$   
 $wR(F^2) = 0.121$   
 $S = 1.02$   
5188 reflections

253 parameters  
H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 0.41\text{ e \AA}^{-3}$   
 $\Delta\rho_{\text{min}} = -0.27\text{ e \AA}^{-3}$

**Table 1**  
Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O2—H2 $\cdots$ O3	0.84	2.54	2.9325 (13)	109
O3—H3 $\cdots$ O2	0.84	2.54	2.9325 (13)	110
O4—H4 $\cdots$ O3	0.84	2.25	2.7009 (14)	114
N1—H1 $\cdots$ O7 <sup>i</sup>	0.88	2.04	2.8844 (15)	160
O2—H2 $\cdots$ O1 <sup>ii</sup>	0.84	1.86	2.6871 (13)	170
O2—H2 $\cdots$ N2 <sup>ii</sup>	0.84	2.57	2.9293 (15)	107
O3—H3 $\cdots$ O1 <sup>ii</sup>	0.84	1.88	2.7200 (13)	177
O4—H4 $\cdots$ O6 <sup>iii</sup>	0.84	2.14	2.7366 (14)	127
C14—H14 $\cdots$ O2 <sup>iv</sup>	0.95	2.42	3.3539 (17)	167

Symmetry codes: (i)  $x, y - 1, z$ ; (ii)  $x + \frac{1}{2}, -y + \frac{1}{2}, z + \frac{1}{2}$ ; (iii)  $x + 1, y, z + 1$ ; (iv)  $x - \frac{1}{2}, -y + \frac{1}{2}, z - \frac{1}{2}$ .

Data collection: *APEX2* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2008).

The authors thank the University of Malaya for funding this study (Science Fund Grant 12-0-03-2031, 12-02-03-2051 and PJP FS350-2008).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HG2510).

### References

- Abdul Alhadi, A. A., Ali, H. M. & Ng, S. W. (2009). *Acta Cryst. E65*, o909.
- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
- Bruker (2007). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Lyubchova, A., Cossé-Barbi, A., Doucet, J. P., Robert, F., Souron, J.-P. & Quarton, M. (1995). *Acta Cryst. C51*, 1893–1895.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Westrip, S. P. (2008). *publCIF*. In preparation.

## **supplementary materials**

*Acta Cryst.* (2009). E65, o1373 [doi:10.1107/S1600536809018947]

### ***N<sup>1</sup>-(2,4-Dimethoxybenzylidene)-3,4,5-trihydroxybenzohydrazide ethanol solvate***

**A. A. Alhadi, S. M. Saharin, H. Mohd Ali, W. T. Robinson and M. A. Abdulla**

#### **Experimental**

A mixture of 3,4,5-trihydroxybenzoylhydrazide and 2,4-dimethoxybenzaldehyde were heated in ethanol (50 ml) for 12 h. The yellow crystals were obtained by recrystallization from ethanol.

#### **Refinement**

All H atoms were placed at calculated positions (C—H 0.95 - 0.99, N—H 0.88, and O—H 0.84 Å) with  $U_{\text{iso}}(\text{H})$  set to 1.2 - 1.5 times  $U_{\text{eq}}(\text{C}, \text{N}, \text{O})$

#### **Figures**

---

Fig. 1. Thermal ellipsoid plot (Barbour, 2001) of  $\text{C}_{16}\text{H}_{16}\text{N}_2\text{O}_6 \cdot \text{C}_2\text{H}_5\text{OH}$  at 70% probability level. Hydrogen atoms are drawn as sphere of arbitrary radius.

### ***N<sup>1</sup>-(2,4-Dimethoxybenzylidene)-3,4,5-trihydroxybenzohydrazide ethanol solvate***

#### *Crystal data*

$\text{C}_{16}\text{H}_{16}\text{N}_2\text{O}_6 \cdot \text{C}_2\text{H}_6\text{O}$	$F_{000} = 800$
$M_r = 378.38$	$D_x = 1.408 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/n$	Mo $K\alpha$ radiation
Hall symbol: -P 2yn	$\lambda = 0.71073 \text{ \AA}$
$a = 7.83470 (10) \text{ \AA}$	Cell parameters from 3445 reflections
$b = 17.5412 (3) \text{ \AA}$	$\theta = 2.8\text{--}29.1^\circ$
$c = 13.0230 (2) \text{ \AA}$	$\mu = 0.11 \text{ mm}^{-1}$
$\beta = 93.9360 (10)^\circ$	$T = 100 \text{ K}$
$V = 1785.53 (5) \text{ \AA}^3$	Block, yellow
$Z = 4$	$0.31 \times 0.16 \times 0.12 \text{ mm}$

#### *Data collection*

Bruker APEXII CCD area-detector diffractometer	5188 independent reflections
Radiation source: fine-focus sealed tube	3586 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.041$
$T = 100 \text{ K}$	$\theta_{\text{max}} = 30.5^\circ$
$\omega$ scans	$\theta_{\text{min}} = 2.0^\circ$
Absorption correction: multi-scan	$h = -11 \rightarrow 10$

# supplementary materials

---

(SADABS; Sheldrick, 1996)

$T_{\min} = 0.967$ ,  $T_{\max} = 0.987$

19580 measured reflections

$k = -24 \rightarrow 24$

$l = -18 \rightarrow 18$

## Refinement

Refinement on  $F^2$

Secondary atom site location: difference Fourier map

Least-squares matrix: full

Hydrogen site location: inferred from neighbouring sites

$R[F^2 > 2\sigma(F^2)] = 0.044$

H-atom parameters constrained

$wR(F^2) = 0.121$

$$w = 1/[\sigma^2(F_o^2) + (0.0566P)^2 + 0.3538P]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$S = 1.02$

$$(\Delta/\sigma)_{\max} = 0.001$$

5188 reflections

$$\Delta\rho_{\max} = 0.41 \text{ e \AA}^{-3}$$

253 parameters

$$\Delta\rho_{\min} = -0.26 \text{ e \AA}^{-3}$$

Primary atom site location: structure-invariant direct methods

Extinction correction: none

## Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

## Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	1.15941 (17)	0.21466 (7)	0.76768 (10)	0.0156 (3)
H1	1.1458	0.2583	0.7248	0.019*
C2	1.25203 (17)	0.22016 (7)	0.86245 (10)	0.0153 (3)
C3	1.26837 (17)	0.15678 (8)	0.92716 (10)	0.0154 (3)
C4	1.19696 (18)	0.08744 (8)	0.89317 (10)	0.0170 (3)
C5	1.10756 (18)	0.08150 (8)	0.79830 (10)	0.0176 (3)
H5	1.0605	0.0339	0.7760	0.021*
C6	1.08672 (17)	0.14571 (7)	0.73529 (10)	0.0154 (3)
C7	0.98683 (17)	0.14395 (8)	0.63409 (10)	0.0159 (3)
C8	0.74521 (17)	0.00842 (8)	0.49550 (10)	0.0168 (3)
H8	0.7539	-0.0296	0.5476	0.023 (4)*
C9	0.64100 (17)	-0.00596 (8)	0.40063 (10)	0.0165 (3)
C10	0.57036 (17)	-0.07846 (8)	0.38154 (10)	0.0163 (3)
C11	0.47202 (17)	-0.09404 (8)	0.29061 (10)	0.0172 (3)
H11	0.4236	-0.1431	0.2785	0.021*

C12	0.44633 (18)	-0.03640 (8)	0.21828 (11)	0.0188 (3)
C13	0.51352 (19)	0.03637 (8)	0.23587 (11)	0.0209 (3)
H13	0.4939	0.0754	0.1859	0.025*
C14	0.60860 (18)	0.05093 (8)	0.32640 (11)	0.0190 (3)
H14	0.6533	0.1007	0.3389	0.023*
C15	0.2879 (2)	-0.11915 (8)	0.09897 (11)	0.0227 (3)
H15A	0.3824	-0.1557	0.0978	0.034*
H15B	0.2270	-0.1169	0.0307	0.034*
H15C	0.2087	-0.1354	0.1498	0.034*
C16	0.5321 (2)	-0.20516 (8)	0.44383 (12)	0.0247 (3)
H16A	0.4072	-0.2007	0.4358	0.037*
H16B	0.5647	-0.2364	0.5044	0.037*
H16C	0.5737	-0.2292	0.3824	0.037*
C17	0.7901 (2)	0.81507 (9)	0.72385 (14)	0.0313 (4)
H19A	0.6929	0.8370	0.6826	0.047*
H19B	0.7482	0.7866	0.7818	0.047*
H19C	0.8531	0.7806	0.6809	0.047*
C18	0.9073 (2)	0.87827 (9)	0.76416 (12)	0.0259 (3)
H18A	0.8458	0.9110	0.8113	0.031*
H18B	1.0074	0.8559	0.8038	0.031*
N1	0.92094 (15)	0.07647 (6)	0.60343 (9)	0.0168 (2)
H1'	0.9379	0.0357	0.6422	0.030 (5)*
N2	0.82556 (15)	0.07168 (6)	0.50974 (9)	0.0174 (2)
O1	0.96209 (13)	0.20261 (5)	0.58110 (7)	0.0219 (2)
O2	1.32670 (14)	0.28879 (5)	0.88620 (7)	0.0218 (2)
H2	1.3569	0.2901	0.9493	0.033*
O3	1.35316 (13)	0.15568 (6)	1.02261 (7)	0.0209 (2)
H3	1.3877	0.1998	1.0381	0.031*
O4	1.21299 (15)	0.02350 (6)	0.95174 (8)	0.0273 (3)
H4	1.2718	0.0332	1.0065	0.041*
O5	0.60584 (13)	-0.13118 (6)	0.45673 (7)	0.0224 (2)
O6	0.35424 (14)	-0.04532 (6)	0.12604 (8)	0.0264 (3)
O7	0.96498 (14)	0.92367 (6)	0.68270 (8)	0.0229 (2)
H7	1.0291	0.8975	0.6475	0.034*

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C1	0.0190 (6)	0.0138 (6)	0.0137 (6)	0.0025 (5)	-0.0015 (5)	0.0010 (5)
C2	0.0187 (6)	0.0123 (6)	0.0148 (6)	-0.0006 (5)	0.0000 (5)	-0.0015 (5)
C3	0.0175 (6)	0.0163 (6)	0.0119 (6)	0.0006 (5)	-0.0021 (5)	-0.0010 (5)
C4	0.0214 (7)	0.0143 (6)	0.0147 (6)	0.0000 (5)	-0.0029 (5)	0.0027 (5)
C5	0.0223 (7)	0.0140 (6)	0.0159 (7)	-0.0007 (5)	-0.0034 (5)	-0.0003 (5)
C6	0.0171 (6)	0.0156 (6)	0.0131 (6)	0.0011 (5)	-0.0020 (5)	-0.0006 (5)
C7	0.0175 (6)	0.0157 (6)	0.0141 (6)	0.0019 (5)	-0.0016 (5)	-0.0005 (5)
C8	0.0194 (7)	0.0158 (6)	0.0148 (6)	0.0008 (5)	-0.0021 (5)	0.0007 (5)
C9	0.0170 (6)	0.0168 (6)	0.0151 (7)	-0.0002 (5)	-0.0021 (5)	-0.0009 (5)
C10	0.0187 (7)	0.0162 (6)	0.0139 (6)	0.0010 (5)	-0.0010 (5)	0.0013 (5)

## supplementary materials

---

C11	0.0188 (7)	0.0162 (6)	0.0161 (6)	-0.0014 (5)	-0.0020 (5)	-0.0015 (5)
C12	0.0201 (7)	0.0192 (7)	0.0162 (7)	0.0000 (5)	-0.0059 (5)	-0.0012 (5)
C13	0.0274 (8)	0.0156 (6)	0.0186 (7)	-0.0001 (6)	-0.0065 (6)	0.0026 (5)
C14	0.0208 (7)	0.0141 (6)	0.0212 (7)	-0.0009 (5)	-0.0040 (5)	-0.0008 (5)
C15	0.0287 (8)	0.0186 (7)	0.0197 (7)	-0.0034 (6)	-0.0074 (6)	-0.0025 (6)
C16	0.0339 (8)	0.0158 (7)	0.0236 (8)	-0.0045 (6)	-0.0042 (6)	0.0033 (6)
C17	0.0369 (9)	0.0197 (7)	0.0381 (10)	-0.0044 (7)	0.0079 (7)	-0.0009 (7)
C18	0.0344 (8)	0.0217 (7)	0.0212 (7)	0.0002 (6)	-0.0020 (6)	0.0049 (6)
N1	0.0215 (6)	0.0151 (5)	0.0129 (5)	-0.0010 (4)	-0.0063 (4)	0.0013 (4)
N2	0.0199 (6)	0.0172 (6)	0.0142 (6)	0.0000 (5)	-0.0064 (4)	-0.0011 (4)
O1	0.0323 (6)	0.0142 (5)	0.0176 (5)	0.0012 (4)	-0.0101 (4)	0.0016 (4)
O2	0.0353 (6)	0.0145 (5)	0.0145 (5)	-0.0057 (4)	-0.0070 (4)	0.0012 (4)
O3	0.0310 (6)	0.0167 (5)	0.0137 (5)	-0.0028 (4)	-0.0079 (4)	0.0011 (4)
O4	0.0446 (7)	0.0167 (5)	0.0184 (5)	-0.0065 (5)	-0.0141 (5)	0.0056 (4)
O5	0.0320 (6)	0.0162 (5)	0.0177 (5)	-0.0042 (4)	-0.0076 (4)	0.0030 (4)
O6	0.0382 (6)	0.0176 (5)	0.0209 (5)	-0.0043 (4)	-0.0160 (5)	0.0010 (4)
O7	0.0295 (6)	0.0163 (5)	0.0232 (6)	0.0001 (4)	0.0034 (4)	-0.0001 (4)

*Geometric parameters ( $\text{\AA}$ ,  $^{\circ}$ )*

C1—C6	1.3900 (18)	C13—C14	1.3745 (19)
C1—C2	1.3914 (18)	C13—H13	0.9500
C1—H1	0.9500	C14—H14	0.9500
C2—O2	1.3647 (15)	C15—O6	1.4308 (16)
C2—C3	1.3958 (18)	C15—H15A	0.9800
C3—O3	1.3687 (15)	C15—H15B	0.9800
C3—C4	1.3979 (18)	C15—H15C	0.9800
C4—O4	1.3574 (16)	C16—O5	1.4258 (17)
C4—C5	1.3818 (18)	C16—H16A	0.9800
C5—C6	1.3965 (18)	C16—H16B	0.9800
C5—H5	0.9500	C16—H16C	0.9800
C6—C7	1.4862 (18)	C17—C18	1.511 (2)
C7—O1	1.2466 (16)	C17—H19A	0.9800
C7—N1	1.3412 (17)	C17—H19B	0.9800
C8—N2	1.2830 (17)	C17—H19C	0.9800
C8—C9	1.4554 (18)	C18—O7	1.4247 (18)
C8—H8	0.9500	C18—H18A	0.9900
C9—C14	1.4004 (19)	C18—H18B	0.9900
C9—C10	1.4023 (18)	N1—N2	1.3889 (15)
C10—O5	1.3615 (16)	N1—H1'	0.8800
C10—C11	1.3948 (18)	O2—H2	0.8400
C11—C12	1.3869 (19)	O3—H3	0.8400
C11—H11	0.9500	O4—H4	0.8400
C12—O6	1.3672 (16)	O7—H7	0.8400
C12—C13	1.3938 (19)		
C6—C1—C2	120.50 (12)	C12—C13—H13	120.4
C6—C1—H1	119.8	C13—C14—C9	121.44 (13)
C2—C1—H1	119.8	C13—C14—H14	119.3
O2—C2—C1	116.85 (12)	C9—C14—H14	119.3

O2—C2—C3	123.07 (11)	O6—C15—H15A	109.5
C1—C2—C3	120.06 (12)	O6—C15—H15B	109.5
O3—C3—C2	125.32 (12)	H15A—C15—H15B	109.5
O3—C3—C4	115.73 (11)	O6—C15—H15C	109.5
C2—C3—C4	118.93 (11)	H15A—C15—H15C	109.5
O4—C4—C5	117.51 (12)	H15B—C15—H15C	109.5
O4—C4—C3	121.43 (12)	O5—C16—H16A	109.5
C5—C4—C3	121.07 (12)	O5—C16—H16B	109.5
C4—C5—C6	119.75 (12)	H16A—C16—H16B	109.5
C4—C5—H5	120.1	O5—C16—H16C	109.5
C6—C5—H5	120.1	H16A—C16—H16C	109.5
C1—C6—C5	119.63 (12)	H16B—C16—H16C	109.5
C1—C6—C7	117.84 (11)	C18—C17—H19A	109.5
C5—C6—C7	122.52 (12)	C18—C17—H19B	109.5
O1—C7—N1	121.44 (12)	H19A—C17—H19B	109.5
O1—C7—C6	121.75 (12)	C18—C17—H19C	109.5
N1—C7—C6	116.79 (12)	H19A—C17—H19C	109.5
N2—C8—C9	120.91 (12)	H19B—C17—H19C	109.5
N2—C8—H8	119.5	O7—C18—C17	111.58 (13)
C9—C8—H8	119.5	O7—C18—H18A	109.3
C14—C9—C10	118.24 (12)	C17—C18—H18A	109.3
C14—C9—C8	121.81 (12)	O7—C18—H18B	109.3
C10—C9—C8	119.95 (12)	C17—C18—H18B	109.3
O5—C10—C11	123.45 (12)	H18A—C18—H18B	108.0
O5—C10—C9	115.45 (11)	C7—N1—N2	119.19 (11)
C11—C10—C9	121.09 (12)	C7—N1—H1'	120.4
C12—C11—C10	118.67 (12)	N2—N1—H1'	120.4
C12—C11—H11	120.7	C8—N2—N1	114.10 (11)
C10—C11—H11	120.7	C2—O2—H2	109.5
O6—C12—C11	123.82 (12)	C3—O3—H3	109.5
O6—C12—C13	114.82 (12)	C4—O4—H4	109.5
C11—C12—C13	121.36 (12)	C10—O5—C16	118.18 (11)
C14—C13—C12	119.17 (13)	C12—O6—C15	118.75 (11)
C14—C13—H13	120.4	C18—O7—H7	109.5

*Hydrogen-bond geometry (Å, °)*

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
O2—H2···O3	0.84	2.54	2.9325 (13)	109
O3—H3···O2	0.84	2.54	2.9325 (13)	110
O4—H4···O3	0.84	2.25	2.7009 (14)	114
N1—H1'···O7 <sup>i</sup>	0.88	2.04	2.8844 (15)	160
O2—H2···O1 <sup>ii</sup>	0.84	1.86	2.6871 (13)	170
O2—H2···N2 <sup>ii</sup>	0.84	2.57	2.9293 (15)	107
O3—H3···O1 <sup>ii</sup>	0.84	1.88	2.7200 (13)	177
O4—H4···O6 <sup>iii</sup>	0.84	2.14	2.7366 (14)	127
C14—H14···O2 <sup>iv</sup>	0.95	2.42	3.3539 (17)	167

Symmetry codes: (i)  $x, y-1, z$ ; (ii)  $x+1/2, -y+1/2, z+1/2$ ; (iii)  $x+1, y, z+1$ ; (iv)  $x-1/2, -y+1/2, z-1/2$ .

## supplementary materials

---

Fig. 1

