

Dichloridobis(2-methylquinolin-8-olato- $\kappa^2 N,O$)tin(IV)

Kong Mun Lo and Seik Weng Ng*

Department of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia
Correspondence e-mail: seikweng@um.edu.my


Received 27 May 2009; accepted 28 May 2009

Key indicators: single-crystal X-ray study; $T = 133$ K; mean $\sigma(C-C) = 0.004$ Å;
 R factor = 0.023; wR factor = 0.060; data-to-parameter ratio = 17.0.

The bis-chelated Sn atom in the title compound, $[Sn(C_{10}H_8NO)_2Cl_2]$, exists in a distorted *cis*-Cl₂,*cis*-N₂,*trans*-O₂ octahedral environment.

Related literature

For the crystal structure of dichloridobis(8-oxidoquinoline), see: Archer *et al.* (1987).



Experimental

Crystal data

$[Sn(C_{10}H_8NO)_2Cl_2]$	$\gamma = 109.236 (1)^\circ$
$M_r = 505.94$	$V = 926.29 (2) \text{ \AA}^3$
Triclinic, $P\bar{1}$	$Z = 2$
$a = 7.9651 (1) \text{ \AA}$	Mo $K\alpha$ radiation
$b = 9.6336 (1) \text{ \AA}$	$\mu = 1.69 \text{ mm}^{-1}$
$c = 12.8337 (2) \text{ \AA}$	$T = 133 \text{ K}$
$\alpha = 94.599 (1)^\circ$	$0.20 \times 0.10 \times 0.05 \text{ mm}$
$\beta = 90.262 (1)^\circ$	

Data collection

Bruker SMART APEX	7670 measured reflections
diffractometer	4189 independent reflections
Absorption correction: multi-scan	3905 reflections with $I > 2\sigma(I)$
(SADABS; Sheldrick, 1996)	
$T_{\min} = 0.729$, $T_{\max} = 0.920$	$R_{\text{int}} = 0.015$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.023$	246 parameters
$wR(F^2) = 0.060$	H-atom parameters constrained
$S = 1.03$	$\Delta\rho_{\max} = 0.79 \text{ e \AA}^{-3}$
4189 reflections	$\Delta\rho_{\min} = -0.34 \text{ e \AA}^{-3}$

Data collection: *APEX2* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2009).

We thank the University of Malaya (RG020/09AFR) for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: TK2467).

References

- Archer, S. J., Koch, K. R. & Schmidt, S. (1987). *Inorg. Chim. Acta*, **126**, 209–218.
- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
- Bruker (2007). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Westrip, S. P. (2009). *publCIF*. In preparation.

supplementary materials

Acta Cryst. (2009). E65, m719 [doi:10.1107/S1600536809020340]

Dichloridobis(2-methylquinolin-8-olato- κ^2N,O)tin(IV)

K. M. Lo and S. W. Ng

Experimental

Di(*p*-chlorobenzyl)tin dichloride (0.44 g, 1 mmol) and 8-hydroxyquinaldine (0.16 g, 1 mmol) were dissolved in chloroform (100 ml) and the solution was heated for 1 hour. Slow evaporation of the filtrate gave yellow crystals. The organic groups at tin were probably cleaved by the heterocycle in the reaction.

Refinement

Hydrogen atoms were placed at calculated positions (C–H 0.95–0.98 Å) and were treated as riding on their parent atoms, with $U(H)$ set to 1.2–1.5 times $U_{\text{eq}}(\text{C})$.

Figures




Fig. 1. Thermal ellipsoid plot (Barbour, 2001) of $\text{SnCl}_2(\text{C}_{10}\text{H}_8\text{NO})_2$ at the 70% probability level. Hydrogen atoms are drawn as spheres of arbitrary radius.

Dichloridobis(2-methylquinolin-8-olato- κ^2N,O)tin(IV)

Crystal data

[$\text{Sn}(\text{C}_{10}\text{H}_8\text{NO})_2\text{Cl}_2$]	$Z = 2$
$M_r = 505.94$	$F_{000} = 500$
Triclinic, $P\bar{1}$	$D_x = 1.814 \text{ Mg m}^{-3}$
Hall symbol: -P 1	Mo $K\alpha$ radiation
$a = 7.9651 (1) \text{ \AA}$	$\lambda = 0.71073 \text{ \AA}$
$b = 9.6336 (1) \text{ \AA}$	Cell parameters from 9941 reflections
$c = 12.8337 (2) \text{ \AA}$	$\theta = 2.4\text{--}28.2^\circ$
$\alpha = 94.599 (1)^\circ$	$\mu = 1.69 \text{ mm}^{-1}$
$\beta = 90.262 (1)^\circ$	$T = 133 \text{ K}$
$\gamma = 109.236 (1)^\circ$	Prism, pale yellow
$V = 926.29 (2) \text{ \AA}^3$	$0.20 \times 0.10 \times 0.05 \text{ mm}$

supplementary materials

Data collection

Bruker SMART APEX diffractometer	4189 independent reflections
Radiation source: fine-focus sealed tube	3905 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.015$
$T = 133 \text{ K}$	$\theta_{\text{max}} = 27.5^\circ$
ω scans	$\theta_{\text{min}} = 1.6^\circ$
Absorption correction: Multi-scan (SADABS; Sheldrick, 1996)	$h = -10 \rightarrow 10$
$T_{\text{min}} = 0.729, T_{\text{max}} = 0.920$	$k = -12 \rightarrow 12$
7670 measured reflections	$l = -16 \rightarrow 16$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.023$	H-atom parameters constrained
$wR(F^2) = 0.060$	$w = 1/[\sigma^2(F_o^2) + (0.0346P)^2 + 0.4719P]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.03$	$(\Delta/\sigma)_{\text{max}} = 0.001$
4189 reflections	$\Delta\rho_{\text{max}} = 0.79 \text{ e \AA}^{-3}$
246 parameters	$\Delta\rho_{\text{min}} = -0.34 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Sn1	0.276677 (18)	0.727420 (15)	0.279378 (11)	0.01746 (6)
Cl1	0.03285 (7)	0.77602 (6)	0.20001 (4)	0.02382 (12)
Cl2	0.24196 (8)	0.82902 (6)	0.44903 (4)	0.02532 (12)
N1	0.4783 (2)	0.6315 (2)	0.34153 (14)	0.0183 (4)
N2	0.3536 (2)	0.6733 (2)	0.11290 (15)	0.0200 (4)
O1	0.1236 (2)	0.51786 (17)	0.29867 (13)	0.0221 (3)
O2	0.4661 (2)	0.91742 (17)	0.25039 (13)	0.0240 (3)
C1	0.7366 (3)	0.8568 (3)	0.37939 (19)	0.0250 (5)
H1A	0.6522	0.9047	0.4057	0.038*
H1B	0.7746	0.8885	0.3101	0.038*
H1C	0.8406	0.8847	0.4275	0.038*
C2	0.6490 (3)	0.6930 (3)	0.37157 (17)	0.0200 (4)
C3	0.7488 (3)	0.6025 (3)	0.39744 (19)	0.0247 (5)
H3	0.8709	0.6470	0.4177	0.030*
C4	0.6739 (3)	0.4537 (3)	0.39396 (19)	0.0260 (5)
H4	0.7444	0.3948	0.4093	0.031*
C5	0.4912 (3)	0.3865 (3)	0.36755 (17)	0.0225 (5)

C6	0.3986 (4)	0.2331 (3)	0.36603 (19)	0.0283 (5)
H6	0.4603	0.1673	0.3807	0.034*
C7	0.2184 (4)	0.1802 (3)	0.3431 (2)	0.0311 (6)
H7	0.1567	0.0772	0.3419	0.037*
C8	0.1239 (3)	0.2753 (3)	0.32133 (19)	0.0272 (5)
H8	-0.0009	0.2360	0.3073	0.033*
C9	0.2103 (3)	0.4248 (2)	0.32009 (17)	0.0208 (4)
C10	0.3970 (3)	0.4819 (2)	0.34239 (17)	0.0192 (4)
C11	0.1273 (3)	0.4305 (3)	0.0696 (2)	0.0285 (5)
H11A	0.0424	0.4658	0.1094	0.043*
H11B	0.1706	0.3684	0.1120	0.043*
H11C	0.0685	0.3729	0.0053	0.043*
C12	0.2808 (3)	0.5597 (3)	0.04230 (18)	0.0228 (5)
C13	0.3471 (4)	0.5605 (3)	-0.0607 (2)	0.0302 (5)
H13	0.2915	0.4812	-0.1117	0.036*
C14	0.4896 (4)	0.6746 (3)	-0.0858 (2)	0.0313 (6)
H14	0.5352	0.6729	-0.1540	0.038*
C15	0.5713 (3)	0.7955 (3)	-0.01254 (19)	0.0269 (5)
C16	0.7162 (3)	0.9198 (3)	-0.0338 (2)	0.0348 (6)
H16	0.7712	0.9228	-0.0995	0.042*
C17	0.7769 (4)	1.0357 (3)	0.0405 (2)	0.0374 (7)
H17	0.8772	1.1177	0.0265	0.045*
C18	0.6947 (3)	1.0370 (3)	0.1376 (2)	0.0311 (6)
H18	0.7378	1.1204	0.1871	0.037*
C19	0.5515 (3)	0.9171 (3)	0.16080 (19)	0.0238 (5)
C20	0.4932 (3)	0.7921 (3)	0.08655 (18)	0.0212 (5)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Sn1	0.01921 (8)	0.01482 (8)	0.01839 (9)	0.00572 (6)	0.00134 (6)	0.00120 (6)
Cl1	0.0241 (3)	0.0261 (3)	0.0234 (3)	0.0112 (2)	0.0001 (2)	0.0025 (2)
Cl2	0.0310 (3)	0.0255 (3)	0.0213 (3)	0.0128 (2)	0.0006 (2)	-0.0018 (2)
N1	0.0213 (9)	0.0184 (9)	0.0158 (9)	0.0071 (7)	0.0017 (7)	0.0015 (7)
N2	0.0210 (9)	0.0214 (9)	0.0193 (9)	0.0091 (8)	0.0002 (7)	0.0034 (8)
O1	0.0194 (7)	0.0183 (8)	0.0275 (9)	0.0041 (6)	0.0000 (6)	0.0045 (7)
O2	0.0267 (8)	0.0169 (8)	0.0258 (9)	0.0037 (6)	0.0018 (7)	0.0016 (7)
C1	0.0209 (10)	0.0239 (12)	0.0271 (12)	0.0038 (9)	-0.0021 (9)	-0.0004 (10)
C2	0.0215 (10)	0.0230 (11)	0.0152 (10)	0.0067 (9)	0.0024 (8)	0.0019 (9)
C3	0.0206 (10)	0.0332 (13)	0.0220 (12)	0.0110 (10)	0.0017 (9)	0.0042 (10)
C4	0.0322 (12)	0.0309 (13)	0.0214 (12)	0.0187 (11)	0.0019 (9)	0.0048 (10)
C5	0.0310 (12)	0.0241 (11)	0.0150 (11)	0.0124 (10)	0.0016 (9)	0.0026 (9)
C6	0.0436 (14)	0.0219 (12)	0.0236 (12)	0.0161 (11)	-0.0004 (10)	0.0035 (10)
C7	0.0463 (15)	0.0151 (11)	0.0286 (13)	0.0058 (10)	-0.0042 (11)	0.0019 (10)
C8	0.0304 (12)	0.0218 (12)	0.0260 (13)	0.0036 (10)	-0.0031 (10)	0.0040 (10)
C9	0.0265 (11)	0.0172 (10)	0.0182 (11)	0.0064 (9)	0.0004 (9)	0.0020 (9)
C10	0.0262 (11)	0.0173 (10)	0.0144 (10)	0.0080 (9)	0.0014 (8)	0.0005 (8)
C11	0.0342 (13)	0.0235 (12)	0.0256 (13)	0.0074 (10)	-0.0039 (10)	-0.0018 (10)

supplementary materials

C12	0.0259 (11)	0.0239 (11)	0.0216 (12)	0.0123 (9)	-0.0006 (9)	0.0010 (9)
C13	0.0362 (13)	0.0364 (14)	0.0233 (12)	0.0198 (12)	-0.0007 (10)	-0.0004 (11)
C14	0.0365 (13)	0.0439 (15)	0.0199 (12)	0.0213 (12)	0.0050 (10)	0.0065 (11)
C15	0.0265 (12)	0.0349 (14)	0.0241 (12)	0.0150 (10)	0.0038 (10)	0.0088 (11)
C16	0.0294 (13)	0.0446 (16)	0.0329 (15)	0.0120 (12)	0.0106 (11)	0.0184 (13)
C17	0.0258 (12)	0.0406 (16)	0.0446 (17)	0.0049 (11)	0.0063 (11)	0.0231 (14)
C18	0.0285 (12)	0.0271 (13)	0.0349 (14)	0.0035 (10)	-0.0028 (11)	0.0110 (11)
C19	0.0219 (10)	0.0247 (12)	0.0257 (12)	0.0076 (9)	0.0000 (9)	0.0090 (10)
C20	0.0194 (10)	0.0242 (11)	0.0223 (12)	0.0091 (9)	0.0005 (8)	0.0073 (9)

Geometric parameters (\AA , $^\circ$)

Sn1—O2	2.0149 (16)	C6—H6	0.9500
Sn1—O1	2.0211 (16)	C7—C8	1.406 (3)
Sn1—N1	2.2703 (18)	C7—H7	0.9500
Sn1—N2	2.2925 (19)	C8—C9	1.379 (3)
Sn1—Cl2	2.3700 (6)	C8—H8	0.9500
Sn1—C11	2.3846 (5)	C9—C10	1.424 (3)
N1—C2	1.332 (3)	C11—C12	1.495 (3)
N1—C10	1.373 (3)	C11—H11A	0.9800
N2—C12	1.332 (3)	C11—H11B	0.9800
N2—C20	1.374 (3)	C11—H11C	0.9800
O1—C9	1.342 (3)	C12—C13	1.425 (3)
O2—C19	1.339 (3)	C13—C14	1.356 (4)
C1—C2	1.496 (3)	C13—H13	0.9500
C1—H1A	0.9800	C14—C15	1.408 (4)
C1—H1B	0.9800	C14—H14	0.9500
C1—H1C	0.9800	C15—C16	1.409 (4)
C2—C3	1.415 (3)	C15—C20	1.416 (3)
C3—C4	1.356 (4)	C16—C17	1.364 (4)
C3—H3	0.9500	C16—H16	0.9500
C4—C5	1.411 (3)	C17—C18	1.411 (4)
C4—H4	0.9500	C17—H17	0.9500
C5—C6	1.416 (3)	C18—C19	1.384 (3)
C5—C10	1.418 (3)	C18—H18	0.9500
C6—C7	1.377 (4)	C19—C20	1.420 (3)
O2—Sn1—O1	168.37 (6)	C6—C7—H7	119.3
O2—Sn1—N1	92.76 (7)	C8—C7—H7	119.3
O1—Sn1—N1	78.04 (6)	C9—C8—C7	120.7 (2)
O2—Sn1—N2	77.90 (7)	C9—C8—H8	119.7
O1—Sn1—N2	94.63 (7)	C7—C8—H8	119.7
N1—Sn1—N2	88.77 (6)	O1—C9—C8	121.9 (2)
O2—Sn1—Cl2	90.74 (5)	O1—C9—C10	119.2 (2)
O1—Sn1—Cl2	96.53 (5)	C8—C9—C10	118.9 (2)
N1—Sn1—Cl2	91.39 (5)	N1—C10—C5	122.3 (2)
N2—Sn1—Cl2	168.62 (5)	N1—C10—C9	117.31 (19)
O2—Sn1—Cl1	97.04 (5)	C5—C10—C9	120.4 (2)
O1—Sn1—Cl1	91.31 (5)	C12—C11—H11A	109.5
N1—Sn1—Cl1	167.95 (5)	C12—C11—H11B	109.5

N2—Sn1—Cl1	86.47 (5)	H11A—C11—H11B	109.5
Cl2—Sn1—Cl1	95.46 (2)	C12—C11—H11C	109.5
C2—N1—C10	119.95 (19)	H11A—C11—H11C	109.5
C2—N1—Sn1	132.09 (15)	H11B—C11—H11C	109.5
C10—N1—Sn1	107.93 (14)	N2—C12—C13	120.3 (2)
C12—N2—C20	120.0 (2)	N2—C12—C11	120.5 (2)
C12—N2—Sn1	132.03 (15)	C13—C12—C11	119.2 (2)
C20—N2—Sn1	107.76 (15)	C14—C13—C12	120.0 (3)
C9—O1—Sn1	116.18 (13)	C14—C13—H13	120.0
C19—O2—Sn1	116.82 (15)	C12—C13—H13	120.0
C2—C1—H1A	109.5	C13—C14—C15	121.2 (2)
C2—C1—H1B	109.5	C13—C14—H14	119.4
H1A—C1—H1B	109.5	C15—C14—H14	119.4
C2—C1—H1C	109.5	C16—C15—C20	119.3 (2)
H1A—C1—H1C	109.5	C16—C15—C14	124.4 (2)
H1B—C1—H1C	109.5	C20—C15—C14	116.2 (2)
N1—C2—C3	119.7 (2)	C17—C16—C15	119.5 (2)
N1—C2—C1	120.9 (2)	C17—C16—H16	120.2
C3—C2—C1	119.4 (2)	C15—C16—H16	120.2
C4—C3—C2	121.4 (2)	C16—C17—C18	121.7 (2)
C4—C3—H3	119.3	C16—C17—H17	119.1
C2—C3—H3	119.3	C18—C17—H17	119.1
C3—C4—C5	119.9 (2)	C19—C18—C17	120.2 (3)
C3—C4—H4	120.0	C19—C18—H18	119.9
C5—C4—H4	120.0	C17—C18—H18	119.9
C4—C5—C6	124.3 (2)	O2—C19—C18	121.6 (2)
C4—C5—C10	116.5 (2)	O2—C19—C20	119.7 (2)
C6—C5—C10	119.2 (2)	C18—C19—C20	118.7 (2)
C7—C6—C5	119.4 (2)	N2—C20—C15	122.2 (2)
C7—C6—H6	120.3	N2—C20—C19	117.4 (2)
C5—C6—H6	120.3	C15—C20—C19	120.3 (2)
C6—C7—C8	121.4 (2)		
O2—Sn1—N1—C2	14.7 (2)	Sn1—O1—C9—C10	8.9 (3)
O1—Sn1—N1—C2	-172.5 (2)	C7—C8—C9—O1	179.4 (2)
N2—Sn1—N1—C2	92.5 (2)	C7—C8—C9—C10	-0.7 (4)
Cl2—Sn1—N1—C2	-76.09 (19)	C2—N1—C10—C5	-4.4 (3)
Cl1—Sn1—N1—C2	159.21 (17)	Sn1—N1—C10—C5	173.84 (17)
O2—Sn1—N1—C10	-163.29 (14)	C2—N1—C10—C9	173.6 (2)
O1—Sn1—N1—C10	9.52 (14)	Sn1—N1—C10—C9	-8.2 (2)
N2—Sn1—N1—C10	-85.47 (14)	C4—C5—C10—N1	1.2 (3)
Cl2—Sn1—N1—C10	105.91 (13)	C6—C5—C10—N1	-179.5 (2)
Cl1—Sn1—N1—C10	-18.8 (3)	C4—C5—C10—C9	-176.7 (2)
O2—Sn1—N2—C12	-170.0 (2)	C6—C5—C10—C9	2.6 (3)
O1—Sn1—N2—C12	19.05 (19)	O1—C9—C10—N1	0.5 (3)
N1—Sn1—N2—C12	96.95 (19)	C8—C9—C10—N1	-179.3 (2)
Cl2—Sn1—N2—C12	-172.14 (17)	O1—C9—C10—C5	178.6 (2)
Cl1—Sn1—N2—C12	-71.98 (19)	C8—C9—C10—C5	-1.3 (3)
O2—Sn1—N2—C20	4.49 (13)	C20—N2—C12—C13	0.0 (3)
O1—Sn1—N2—C20	-166.49 (13)	Sn1—N2—C12—C13	173.93 (16)

supplementary materials

N1—Sn1—N2—C20	-88.60 (14)	C20—N2—C12—C11	-179.3 (2)
Cl2—Sn1—N2—C20	2.3 (3)	Sn1—N2—C12—C11	-5.4 (3)
Cl1—Sn1—N2—C20	102.47 (13)	N2—C12—C13—C14	2.7 (3)
O2—Sn1—O1—C9	28.5 (4)	C11—C12—C13—C14	-178.0 (2)
N1—Sn1—O1—C9	-9.83 (15)	C12—C13—C14—C15	-1.8 (4)
N2—Sn1—O1—C9	77.92 (16)	C13—C14—C15—C16	-178.1 (2)
Cl2—Sn1—O1—C9	-99.87 (15)	C13—C14—C15—C20	-1.6 (3)
Cl1—Sn1—O1—C9	164.49 (15)	C20—C15—C16—C17	-1.1 (4)
O1—Sn1—O2—C19	44.7 (4)	C14—C15—C16—C17	175.2 (2)
N1—Sn1—O2—C19	82.07 (16)	C15—C16—C17—C18	-2.1 (4)
N2—Sn1—O2—C19	-6.07 (15)	C16—C17—C18—C19	1.8 (4)
Cl2—Sn1—O2—C19	173.50 (15)	Sn1—O2—C19—C18	-173.95 (17)
Cl1—Sn1—O2—C19	-90.91 (15)	Sn1—O2—C19—C20	7.0 (3)
C10—N1—C2—C3	4.3 (3)	C17—C18—C19—O2	-177.2 (2)
Sn1—N1—C2—C3	-173.52 (15)	C17—C18—C19—C20	1.9 (3)
C10—N1—C2—C1	-175.1 (2)	C12—N2—C20—C15	-3.7 (3)
Sn1—N1—C2—C1	7.1 (3)	Sn1—N2—C20—C15	-178.95 (17)
N1—C2—C3—C4	-1.0 (3)	C12—N2—C20—C19	172.74 (19)
C1—C2—C3—C4	178.4 (2)	Sn1—N2—C20—C19	-2.5 (2)
C2—C3—C4—C5	-2.2 (4)	C16—C15—C20—N2	-178.9 (2)
C3—C4—C5—C6	-177.2 (2)	C14—C15—C20—N2	4.4 (3)
C3—C4—C5—C10	2.1 (3)	C16—C15—C20—C19	4.7 (3)
C4—C5—C6—C7	177.4 (2)	C14—C15—C20—C19	-171.9 (2)
C10—C5—C6—C7	-1.8 (3)	O2—C19—C20—N2	-2.5 (3)
C5—C6—C7—C8	-0.2 (4)	C18—C19—C20—N2	178.4 (2)
C6—C7—C8—C9	1.5 (4)	O2—C19—C20—C15	174.04 (19)
Sn1—O1—C9—C8	-171.26 (18)	C18—C19—C20—C15	-5.1 (3)

Fig. 1

