

Di- μ -chlorido-bis[chlorido(*N,N*-dimethylethylenediamine- κ^2 *N,N'*)zinc(II)]

Ming-Ming Yu, Qiu-Zhi Shi, Yu-Na Zhang and Zhan-Xian Li*

Department of Chemistry, Zhengzhou University, Zhengzhou 450001, People's Republic of China

Correspondence e-mail: lizx@zzu.edu.cn


Received 8 April 2009; accepted 22 May 2009

Key indicators: single-crystal X-ray study; $T = 295$ K; mean $\sigma(\text{C}-\text{C}) = 0.011$ Å; R factor = 0.049; wR factor = 0.132; data-to-parameter ratio = 19.8.

The centrosymmetric dinuclear title compound, $[\text{Zn}_2\text{Cl}_4(\text{C}_4\text{H}_{12}\text{N}_2)_2]$, is isostructural with its previously reported Cu^{II} analogue [Phelps, Goodman & Hodgson (1976). *Inorg. Chem.* **15**, 2266–2270]. In the title compound, each of the Zn^{II} ions is coordinated by two N atoms from a chelating *N,N*-dimethylethylenediamine ligand, two bridging Cl atoms and one terminal Cl atom. The coordination environment is distorted square-pyramidal. The Zn–Cl bond distances of the two bridging Cl atoms are distinctly different: the equatorial Cl atom exhibits a Zn–Cl distance of 2.318 (1) Å and the axial Cl atom exhibits a Zn–Cl distance of 2.747 (2) Å, which is significantly longer. The molecule can thus be seen as a dimer of two nearly square-planar monomeric units which are related to each other by an inversion center located in the middle of the dimer. Within one monomeric unit, the Zn atom, the two N atoms and the two Cl atoms are almost coplanar, with a mean deviation of only 0.05 (1) Å from the associated least-squares plane. The Zn \cdots Zn distance within the dimer is 3.472 (3) Å. N–H \cdots Cl and C–H \cdots Cl hydrogen-bond interactions connect neighboring molecules with each other.

Related literature

For the isostructural Cu^{II} complex, see: Phelps *et al.* (1976). For general background on the coordination behaviour of *N,N*-dimethylethylenediamine, see: Basak *et al.* (2007); Hlavinka & Hagadorn (2003); Knight *et al.* (2008). Allen (2002) describes the Cambridge Structural Database.



Experimental

Crystal data

$[\text{Zn}_2\text{Cl}_4(\text{C}_4\text{H}_{12}\text{N}_2)_2]$
 $M_r = 448.85$
 Orthorhombic, *Pbca*
 $a = 9.808$ (2) Å
 $b = 8.5109$ (17) Å
 $c = 20.851$ (4) Å

$V = 1740.5$ (6) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 3.36$ mm⁻¹
 $T = 295$ K
 $0.15 \times 0.12 \times 0.07$ mm

Data collection

Bruker SMART 1K CCD area-detector diffractometer
 Absorption correction: multi-scan (SADABS; Bruker, 2000)
 $T_{\min} = 0.633$, $T_{\max} = 0.799$

7050 measured reflections
 1620 independent reflections
 1300 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.042$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.049$
 $wR(F^2) = 0.132$
 $S = 1.10$
 1620 reflections

82 parameters
 H-atom parameters constrained
 $\Delta\rho_{\max} = 1.14$ e Å⁻³
 $\Delta\rho_{\min} = -0.42$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

<i>D</i> –H \cdots <i>A</i>	<i>D</i> –H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> –H \cdots <i>A</i>
N1–H1D \cdots Cl1 ⁱ	0.90	2.51	3.342 (2)	155
C4–H4C \cdots Cl2	0.96	2.78	3.350 (9)	119
N1–H1A \cdots Cl2 ⁱⁱ	0.90	2.90	3.697 (2)	149

Symmetry codes: (i) $-x + \frac{1}{2}, y - \frac{1}{2}, z$; (ii) $-x + 1, -y + 2, -z + 1$.

Data collection: *SMART* (Bruker, 2000); cell refinement: *SAINT* (Bruker, 2000); data reduction: *SAINT*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL*.

This work was supported by the Natural Science Foundation of China (grant No. 50873093).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: ZL2193).

References

- Allen, F. H. (2002). *Acta Cryst.* **B58**, 380–388.
- Basak, S., Sen, S., Banerjee, S., Mitra, S., Rosair, G. & Rodriguez, M. T. G. (2007). *Polyhedron*, **26**, 5104–5112.
- Bruker (2000). *SMART, SAINT and SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Hlavinka, M. L. & Hagadorn, J. R. (2003). *Chem. Commun.* pp. 2686–2687.
- Knight, P. D., White, J. P. & Williams, C. K. (2008). *Inorg. Chem.* **47**, 11711–11719.
- Phelps, D. W., Goodman, W. H. & Hodgson, D. J. (1976). *Inorg. Chem.* **15**, 2266–2270.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supplementary materials

Acta Cryst. (2009). E65, m744-m745 [doi:10.1107/S1600536809019473]

Di- μ -chlorido-bis[chlorido(*N,N*-dimethylethylenediamine- κ^2N,N')zinc(II)]

M.-M. Yu, Q.-Z. Shi, Y.-N. Zhang and Z.-X. Li

Comment

N,N-Dimethylethylenediamine has the potential to function as a bidentate nitrogen ligand by coordinating to metal ions in a chelating fashion (Hlavinka & Hagadorn, 2003; Knight *et al.*, 2008; Basak *et al.*, 2007). Here, we report the crystal structure of the title compound, an asymmetrically chloro-bridged dimeric zinc(II) complex.

In the centrosymmetric dinuclear title compound, $[\text{Zn}_2\text{Cl}_4(\text{C}_4\text{H}_{12}\text{N}_2)_2]$, each of the Zn^{II} ions is coordinated by two N atoms from a chelating *N,N*-dimethylethylenediamine ligand, two bridging Cl atoms and one terminal Cl atom. The coordination environment is distorted square-pyramidal. In the dimeric structure, two Zn^{II} ions are bridged through the Cl atoms, resulting in a planar Zn_2Cl_2 core. The Zn—Cl bond distances of the two bridging Cl atoms are distinctly different: The equatorial Cl atoms exhibit a Zn—Cl distance of 2.318 (1) Å, the Zn—Cl distances of the axial chlorides are with 2.747 (2) Å significantly longer. The title compound could thus be considered as a dimer of two nearly square planar monomeric units which are related to each other by an inversion center located in the middle of the molecule [symmetry code: 1 - *x*, 2 - *y*, 1 - *z*]. Within one monomeric unit the atoms Zn1, N1, N2, Cl1 and Cl2 are almost coplanar with a mean deviation of only 0.05 (1) Å from the associated least-squares plane.

The methyl substituted N atom N2 is located opposite of the bridging Cl atom Cl1, probably due to its larger steric demand when compared to the unsubstituted NH_2 group and due the ability to form an intramolecular N—H \cdots Cl hydrogen bond to the terminal Cl atom in the other half of the dimer (see Table 1 and below).

The Cambridge Structural Database (Allen, 2002) does not list any crystal structures with a Zn^{II} ion in a square-pyramidal environment with two bridging Cl atoms and one terminal Cl atom. This motif seems to be more typical for Cu^{II} complexes for which the CSD has 15 entries. The structure of the title complex is indeed isostructural to its copper(II) analogue $[\text{CuCl}_2(\text{C}_4\text{H}_{12}\text{N}_2)]_2$ (Phelps *et al.*, 1976). Both structures are very similar, as proved by the distance of M—Cl, the M \cdots M separation and the bridging M—Cl—M angle (Zn—Cl = 2.318 (1) Å, Zn—Cl' = 2.747 (2) Å, Zn \cdots Zn = 3.472 (3) Å, Zn—Cl—Zn = 86.11 (4) °; Cu—Cl = 2.309 (2) Å, Cu—Cl' = 2.734 (3) Å, Cu \cdots Cu = 3.458 (3) Å, Cu—Cl—Cu = 86.11 (8) °).

In the crystal structure, the dimer is strengthened by intramolecular hydrogen bond interactions involving the methyl and amino protons of the ligand and the terminal Cl atom [C4—H4C \cdots Cl2 and N1—H1A \cdots Cl2ⁱ, symmetry code: (i) 1 - *x*, 2 - *y*, 1 - *z*]. An intermolecular N1—H1D \cdots Cl1ⁱⁱ hydrogen bond interaction between the other amino H atom and one of the bridging Cl atoms leads to the formation of a one-dimensional supramolecular chain (Table 1, Fig. 2).

Experimental

Colourless crystals of the title complex were obtained by slow evaporation of a solution in ethanol (20 ml) and water (5 ml) of *N,N*-dimethylethylenediamine (0.044 g, 0.5 mmol) and ZnCl_2 (0.068 g, 0.5 mmol). Yield, 85%. Selected IR data (cm^{-1} , KBr

supplementary materials

pellet): 3342, 3285 (*m*), 3161 (*w*), 3048 (*w*), 1465 (*m*), 1332 (*w*), 1292 (*w*), 1248 (*w*), 1189 (*w*), 1007 (*m*), 937 (*w*), 896 (*w*), 789(*w*), 631 (*m*). Anal. Calcd for $C_8H_{24}Cl_4N_4Zn_2$ requires C, 21.4; H, 5.39; N, 12.48. Found: C, 21.1; H, 5.41; N, 12.23%.

Refinement

The H atoms bound to C and N atoms were placed in calculated positions with C—H = 0.97 Å (CH₂), C—H = 0.96 Å (CH₃) and with $U_{iso}(H) = 1.2U_{eq}(C)$, and with N—H distances of 0.90 Å and with $U_{iso}(H) = 1.2U_{eq}(N)$.

Figures



Fig. 1. A view of title complex, showing 30% probability displacement ellipsoids and the atom-numbering scheme.




Fig. 2. N—H...Cl and C—H...Cl interactions (dashed lines) in the title compound. [Symmetry codes: (i) 1 - *x*, 2 - *y*, 1 - *z*; (ii) 0.5 - *x*, 0.5 + *y*, *z*; (iii) 0.5 + *x*, 1.5 - *y*, 1 - *z*]

Di- μ -chlorido-bis[chlorido(*N,N*-dimethylethylenediamine- κ^2N,N')zinc(II)]

Crystal data

[Zn₂Cl₄(C₄H₁₂N₂)₂]

$M_r = 448.85$

Orthorhombic, *Pbca*

Hall symbol: -P 2ac 2ab

$a = 9.808$ (2) Å

$b = 8.5109$ (17) Å

$c = 20.851$ (4) Å

$V = 1740.5$ (6) Å³

$Z = 4$

$F_{000} = 912$

$D_x = 1.713$ Mg m⁻³

Mo $K\alpha$ radiation

$\lambda = 0.71073$ Å

Cell parameters from 1620 reflections

$\theta = 2.0$ – 25.5°

$\mu = 3.36$ mm⁻¹

$T = 295$ K

Block, colourless

0.15 × 0.12 × 0.07 mm

Data collection

Bruker SMART 1K CCD area-detector diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 295$ K

φ and ω scans

1620 independent reflections

1300 reflections with $I > 2\sigma(I)$

$R_{int} = 0.042$

$\theta_{max} = 25.5^\circ$

$\theta_{min} = 2.0^\circ$

Absorption correction: multi-scan
(SADABS; Bruker, 2000) $h = -11 \rightarrow 11$
 $T_{\min} = 0.633$, $T_{\max} = 0.799$ $k = -6 \rightarrow 10$
 7050 measured reflections $l = -25 \rightarrow 20$

Refinement

Refinement on F^2 Secondary atom site location: difference Fourier map
 Least-squares matrix: full Hydrogen site location: inferred from neighbouring sites
 $R[F^2 > 2\sigma(F^2)] = 0.049$ H-atom parameters constrained
 $wR(F^2) = 0.132$ $w = 1/[\sigma^2(F_o^2) + (0.0632P)^2 + 2.354P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $S = 1.10$ $(\Delta/\sigma)_{\max} < 0.001$
 1620 reflections $\Delta\rho_{\max} = 1.14 \text{ e } \text{\AA}^{-3}$
 82 parameters $\Delta\rho_{\min} = -0.41 \text{ e } \text{\AA}^{-3}$
 Primary atom site location: structure-invariant direct methods Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Zn1	0.43590 (6)	0.94329 (7)	0.42593 (3)	0.0433 (3)
Cl1	0.31460 (12)	1.00877 (18)	0.51751 (7)	0.0526 (4)
Cl2	0.40919 (18)	1.18349 (18)	0.38224 (8)	0.0700 (5)
N1	0.4424 (4)	0.7235 (5)	0.4593 (2)	0.0490 (11)
H1A	0.5032	0.7173	0.4915	0.059*
H1D	0.3600	0.6964	0.4748	0.059*
N2	0.5142 (5)	0.8454 (6)	0.3424 (2)	0.0566 (12)
C3	0.4065 (7)	0.8464 (9)	0.2933 (3)	0.077 (2)
H3A	0.4408	0.8009	0.2544	0.116*
H3B	0.3784	0.9527	0.2853	0.116*
H3C	0.3299	0.7864	0.3082	0.116*
C1	0.4819 (7)	0.6142 (7)	0.4073 (4)	0.078 (2)
H1B	0.5299	0.5250	0.4254	0.093*
H1C	0.4005	0.5752	0.3861	0.093*

supplementary materials

C4	0.6311 (8)	0.9331 (11)	0.3154 (4)	0.099 (3)
H4A	0.6624	0.8815	0.2772	0.148*
H4B	0.7036	0.9364	0.3463	0.148*
H4C	0.6031	1.0382	0.3051	0.148*
C2	0.5679 (9)	0.6908 (9)	0.3608 (4)	0.095 (3)
H2A	0.5753	0.6251	0.3229	0.114*
H2B	0.6586	0.7035	0.3787	0.114*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Zn1	0.0489 (4)	0.0326 (4)	0.0485 (4)	0.0042 (2)	-0.0006 (3)	0.0006 (2)
Cl1	0.0422 (6)	0.0581 (8)	0.0575 (8)	0.0054 (6)	-0.0002 (6)	-0.0108 (6)
Cl2	0.0875 (11)	0.0378 (8)	0.0847 (11)	0.0070 (7)	-0.0056 (9)	0.0152 (7)
N1	0.056 (3)	0.035 (2)	0.056 (3)	-0.0057 (19)	0.000 (2)	0.0071 (19)
N2	0.074 (3)	0.057 (3)	0.039 (2)	0.014 (2)	0.000 (2)	-0.002 (2)
C3	0.089 (5)	0.086 (5)	0.058 (4)	-0.007 (4)	-0.015 (3)	-0.011 (4)
C1	0.074 (4)	0.026 (3)	0.133 (6)	-0.002 (3)	0.026 (4)	-0.004 (3)
C4	0.071 (5)	0.146 (9)	0.079 (5)	-0.003 (5)	0.024 (4)	-0.005 (5)
C2	0.133 (7)	0.071 (5)	0.080 (5)	0.037 (5)	0.015 (5)	-0.016 (4)

Geometric parameters (\AA , $^\circ$)

Zn1—N1	1.997 (4)	C3—H3A	0.9600
Zn1—N2	2.078 (4)	C3—H3B	0.9600
Zn1—Cl2	2.2533 (16)	C3—H3C	0.9600
Zn1—Cl1	2.3179 (14)	C1—C2	1.441 (5)
Zn1—Cl1 ⁱ	2.7468 (15)	C1—H1B	0.9700
Cl1—Zn1 ⁱ	2.7468 (15)	C1—H1C	0.9700
N1—C1	1.481 (8)	C4—H4A	0.9600
N1—H1A	0.9000	C4—H4B	0.9600
N1—H1D	0.9000	C4—H4C	0.9600
N2—C2	1.468 (9)	C2—H2A	0.9700
N2—C3	1.471 (8)	C2—H2B	0.9700
N2—C4	1.480 (9)		
N1—Zn1—N2	84.52 (19)	N2—C3—H3B	109.5
N1—Zn1—Cl2	173.95 (13)	H3A—C3—H3B	109.5
N2—Zn1—Cl2	93.89 (14)	N2—C3—H3C	109.5
N1—Zn1—Cl1	87.40 (14)	H3A—C3—H3C	109.5
N2—Zn1—Cl1	167.59 (16)	H3B—C3—H3C	109.5
Cl2—Zn1—Cl1	93.17 (6)	C2—C1—N1	111.2 (5)
N1—Zn1—Cl1 ⁱ	87.78 (13)	C2—C1—H1B	109.4
N2—Zn1—Cl1 ⁱ	95.19 (14)	N1—C1—H1B	109.4
Cl2—Zn1—Cl1 ⁱ	98.19 (6)	C2—C1—H1C	109.4
Cl1—Zn1—Cl1 ⁱ	93.89 (4)	N1—C1—H1C	109.4
Zn1—Cl1—Zn1 ⁱ	86.11 (4)	H1B—C1—H1C	108.0
C1—N1—Zn1	110.0 (4)	N2—C4—H4A	109.5

C1—N1—H1A	109.7	N2—C4—H4B	109.5
Zn1—N1—H1A	109.7	H4A—C4—H4B	109.5
C1—N1—H1D	109.7	N2—C4—H4C	109.5
Zn1—N1—H1D	109.7	H4A—C4—H4C	109.5
H1A—N1—H1D	108.2	H4B—C4—H4C	109.5
C2—N2—C3	116.5 (6)	C1—C2—N2	111.8 (6)
C2—N2—C4	105.9 (6)	C1—C2—H2A	109.3
C3—N2—C4	106.8 (5)	N2—C2—H2A	109.3
C2—N2—Zn1	105.8 (4)	C1—C2—H2B	109.3
C3—N2—Zn1	108.4 (4)	N2—C2—H2B	109.3
C4—N2—Zn1	113.8 (4)	H2A—C2—H2B	107.9
N2—C3—H3A	109.5		
N1—Zn1—Cl1—Zn1 ⁱ	87.59 (12)	Cl2—Zn1—N2—C3	-65.5 (4)
N2—Zn1—Cl1—Zn1 ⁱ	136.9 (6)	Cl1—Zn1—N2—C3	59.0 (8)
Cl2—Zn1—Cl1—Zn1 ⁱ	-98.44 (6)	Cl1 ⁱ —Zn1—N2—C3	-164.1 (4)
Cl1 ⁱ —Zn1—Cl1—Zn1 ⁱ	0.0	N1—Zn1—N2—C4	-132.8 (5)
N2—Zn1—N1—C1	-6.1 (4)	Cl2—Zn1—N2—C4	53.0 (5)
Cl1—Zn1—N1—C1	164.4 (4)	Cl1—Zn1—N2—C4	177.6 (5)
Cl1 ⁱ —Zn1—N1—C1	-101.6 (4)	Cl1 ⁱ —Zn1—N2—C4	-45.6 (5)
N1—Zn1—N2—C2	-17.0 (5)	Zn1—N1—C1—C2	29.5 (7)
Cl2—Zn1—N2—C2	168.8 (5)	N1—C1—C2—N2	-46.2 (9)
Cl1—Zn1—N2—C2	-66.6 (8)	C3—N2—C2—C1	-82.0 (8)
Cl1 ⁱ —Zn1—N2—C2	70.2 (5)	C4—N2—C2—C1	159.5 (7)
N1—Zn1—N2—C3	108.6 (4)	Zn1—N2—C2—C1	38.5 (8)

Symmetry codes: (i) $-x+1, -y+2, -z+1$.

Hydrogen-bond geometry ($\text{\AA}, ^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
N1—H1D \cdots Cl1 ⁱⁱ	0.90	2.51	3.342 (2)	155
C4—H4C \cdots Cl2	0.96	2.78	3.350 (9)	119
N1—H1A \cdots Cl2 ⁱ	0.90	2.90	3.697 (2)	149

Symmetry codes: (ii) $-x+1/2, y-1/2, z$; (i) $-x+1, -y+2, -z+1$.

Fig. 1

