

{6,6'-Dimethoxy-2,2'-[cyclohexane-1,2-diylbis(nitrilomethylidyne)]diphenolato- $\kappa^4 O^1, N, N', O^1$ }iron(II) monohydrate

Peng-Fei Yan, Yan Bao, Hong-Feng Li and Guang-Ming Li*

School of Chemistry and Materials Science, Heilongjiang University, Harbin 150080, People's Republic of China

Correspondence e-mail: gmli@hlju.edu.cn


Received 23 May 2009; accepted 11 June 2009

 Key indicators: single-crystal X-ray study; $T = 291$ K; mean $\sigma(C-C) = 0.005$ Å; R factor = 0.042; wR factor = 0.107; data-to-parameter ratio = 16.8.

In the title complex, $[Fe(C_{22}H_{24}N_2O_4)] \cdot H_2O$, the Fe^{II} center is four-coordinated by two O and two N atoms from 2,2'-[6,6'-dimethoxycyclohexane-1,2-diylbis(nitrilomethylidyne)]-diphenolate (*L*) ligands in a distorted square-planar geometry. Uncoordinated water and FeL molecules are paired *via* intermolecular water–methoxy $O-H \cdots O$ hydrogen bonds.

Related literature

For a manganese complex of a similar Schiff base ligand, see: Watkinson *et al.* (1999). For the isotopic Co^{II} compound, see: Bao *et al.* (2009).



Experimental

Crystal data

 $[Fe(C_{22}H_{24}N_2O_4)] \cdot H_2O$
 $M_r = 454.30$

 Monoclinic, $P2_1/n$
 $a = 11.243$ (5) Å
 $b = 10.617$ (3) Å
 $c = 17.863$ (7) Å
 $\beta = 107.042$ (14)°
 $V = 2038.5$ (13) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.78$ mm⁻¹
 $T = 291$ K
 $0.22 \times 0.21 \times 0.18$ mm

Data collection

 Rigaku R-Axis RAPID
 diffractometer
 Absorption correction: multi-scan
 (*ABSCOR*; Higashi, 1995)
 $T_{min} = 0.846$, $T_{max} = 0.873$

 18680 measured reflections
 4584 independent reflections
 3446 reflections with $I > 2\sigma(I)$
 $R_{int} = 0.054$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.042$
 $wR(F^2) = 0.107$
 $S = 1.03$
 4584 reflections

 273 parameters
 H-atom parameters constrained
 $\Delta\rho_{max} = 0.30$ e Å⁻³
 $\Delta\rho_{min} = -0.35$ e Å⁻³
Table 1

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
$O5-H2 \cdots O2$	0.85	2.15	2.915 (4)	149
$O5-H1 \cdots O4$	0.85	2.06	2.898 (4)	169

Data collection: *RAPID-AUTO* (Rigaku, 1998); cell refinement: *RAPID-AUTO*; data reduction: *CrystalStructure* (Rigaku/MS, 2002); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXL97*.

The authors gratefully acknowledge financial support from the National Natural Science Foundation of China (grant Nos. 20572018 and 20672032), Heilongjiang Province (grant Nos. 1055HZ001, ZJG0504 and JC200605) and Heilongjiang University.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CV2567).

References

- Bao, Y., Li, H.-F., Yan, P.-F., Li, G.-M. & Hou, G.-F. (2009). *Acta Cryst.* **E65**, m770.
 Higashi, T. (1995). *ABSCOR*. Rigaku Corporation, Tokyo, Japan.
 Rigaku (1998). *RAPID-AUTO*. Rigaku Corporation, Tokyo, Japan.
 Rigaku/MS (2002). *CrystalStructure*. Rigaku/MS Inc., The Woodlands, Texas, USA.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Watkinson, M., Fondo, M., Bermejo, M. R., Sousa, A., McAuliffe, C. A., Pritchard, R. G., Jaiboon, N., Aurangzeb, N. & Naem, M. (1999). *J. Chem. Soc. Dalton Trans.* pp. 31–41.

supplementary materials

Acta Cryst. (2009). E65, m832 [doi:10.1107/S1600536809022235]

{6,6'-Dimethoxy-2,2'-[cyclohexane-1,2-diylbis(nitrilomethylidyne)]diphenolato- $\kappa^4 O^1, N, N', O^1$ }iron(II) monohydrate

P.-F. Yan, Y. Bao, H.-F. Li and G.-M. Li

Comment

In the title compound (Fig. 1), the Fe^{II} ion is four-coordinated by the tetradentate Schiff base ligand in a square planar environment in a manner observed earlier for a manganese complex (Watkinson *et al.*, 1999). Uncoordinated water molecule is paired with the main molecule by the O—H···O hydrogen bonds (Table 1, Fig. 1).

Experimental

The title complex was obtained by the treatment of anhydrous ferrous chloride with the Schiff base in methanol/acetone (2:3). The yellow clear mixture turned to black precipitation immediately, stirred for 4 h; diethyl ether was allowed to diffuse slowly into the solution of the filtrate. Black single crystals were obtained after several days. Analysis calculated for C₂₂H₂₆FeN₂O₅: C, 58.16; H, 5.77; N, 6.17; Fe, 12.29; found: C, 57.56; H, 5.23; N, 6.77; F, 12.79%.

Refinement

H atoms bound to C atoms were placed in calculated positions and treated as riding on their parent atoms, with C—H = 0.93 Å (aromatic C), C—H = 0.97 Å (methylene C), C—H = 0.98 Å (methine C), and with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ or C—H = 0.96 Å (methly C) and with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$. Water H atoms were initially located in a difference Fourier map, but they were treated as riding on their parent atoms with O—H = 0.85 Å and with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{O})$.

Figures




Fig. 1. View of the title compound showing the atomic numbering and 30% probability displacement ellipsoids. Dashed lines indicate the hydrogen-bonding interactions.

{6,6'-Dimethoxy-2,2'-[cyclohexane-1,2-diylbis(nitrilomethylidyne)]diphenolato- $\kappa^4 O^1, N, N', O^1$ }iron(II) monohydrate

Crystal data

[Fe(C₂₂H₂₄N₂O₄)]·H₂O

$F_{000} = 952$

supplementary materials

$M_r = 454.30$	$D_x = 1.480 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/n$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2yn	Cell parameters from 13436 reflections
$a = 11.243 (5) \text{ \AA}$	$\theta = 3.1\text{--}27.5^\circ$
$b = 10.617 (3) \text{ \AA}$	$\mu = 0.78 \text{ mm}^{-1}$
$c = 17.863 (7) \text{ \AA}$	$T = 291 \text{ K}$
$\beta = 107.042 (14)^\circ$	Block, black
$V = 2038.5 (13) \text{ \AA}^3$	$0.22 \times 0.21 \times 0.18 \text{ mm}$
$Z = 4$	

Data collection

Rigaku R-Axis RAPID diffractometer	4584 independent reflections
Radiation source: fine-focus sealed tube	3446 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.054$
$T = 291 \text{ K}$	$\theta_{\text{max}} = 27.5^\circ$
ω scans	$\theta_{\text{min}} = 3.1^\circ$
Absorption correction: multi-scan (ABSCOR; Higashi, 1995)	$h = -14 \rightarrow 14$
$T_{\text{min}} = 0.846$, $T_{\text{max}} = 0.873$	$k = -13 \rightarrow 13$
18680 measured reflections	$l = -23 \rightarrow 23$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.042$	H-atom parameters constrained
$wR(F^2) = 0.107$	$w = 1/[\sigma^2(F_o^2) + (0.0436P)^2 + 1.092P]$
$S = 1.03$	where $P = (F_o^2 + 2F_c^2)/3$
4584 reflections	$(\Delta/\sigma)_{\text{max}} = 0.001$
273 parameters	$\Delta\rho_{\text{max}} = 0.30 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	$\Delta\rho_{\text{min}} = -0.35 \text{ e \AA}^{-3}$
	Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -

factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. *R*-factors based on F^2 are statistically about twice as large as those based on *F*, and *R*- factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Fe1	0.65626 (3)	0.46981 (3)	0.162776 (18)	0.02766 (11)
N1	0.7304 (2)	0.52432 (19)	0.08782 (12)	0.0356 (5)
O1	0.55829 (18)	0.34802 (16)	0.09933 (10)	0.0414 (4)
O3	0.57113 (19)	0.43064 (17)	0.23484 (11)	0.0447 (5)
N2	0.7650 (2)	0.5782 (2)	0.23003 (12)	0.0397 (5)
C7	0.6991 (3)	0.4904 (2)	0.01541 (15)	0.0385 (6)
H7	0.7339	0.5350	-0.0178	0.046*
C1	0.5517 (2)	0.3230 (2)	0.02601 (15)	0.0370 (6)
C2	0.4762 (3)	0.2205 (3)	-0.01171 (17)	0.0439 (6)
C20	0.5684 (3)	0.5024 (2)	0.29420 (15)	0.0401 (6)
O4	0.3917 (2)	0.3866 (2)	0.29798 (13)	0.0597 (6)
C19	0.4729 (3)	0.4826 (3)	0.33048 (17)	0.0481 (7)
O2	0.4206 (2)	0.1564 (2)	0.03582 (13)	0.0636 (6)
C8	0.8176 (3)	0.6325 (2)	0.11520 (15)	0.0394 (6)
H8	0.7676	0.7097	0.1075	0.047*
C14	0.7552 (3)	0.6262 (2)	0.29455 (15)	0.0414 (6)
H14	0.8169	0.6814	0.3220	0.050*
C6	0.6156 (2)	0.3903 (2)	-0.01837 (15)	0.0372 (6)
C4	0.5227 (3)	0.2627 (3)	-0.13222 (17)	0.0517 (7)
H4	0.5117	0.2439	-0.1847	0.062*
C15	0.6547 (3)	0.5989 (3)	0.32603 (15)	0.0421 (6)
C5	0.5980 (3)	0.3603 (3)	-0.09773 (16)	0.0464 (7)
H5	0.6381	0.4075	-0.1269	0.056*
C13	0.8726 (3)	0.6129 (2)	0.20281 (16)	0.0411 (6)
H13	0.9091	0.6918	0.2277	0.049*
C12	0.9700 (3)	0.5089 (3)	0.22201 (18)	0.0502 (7)
H12A	0.9306	0.4287	0.2040	0.060*
H12B	1.0067	0.5038	0.2783	0.060*
C3	0.4622 (3)	0.1911 (3)	-0.08882 (17)	0.0495 (7)
H3	0.4125	0.1234	-0.1121	0.059*
C9	0.9182 (3)	0.6509 (3)	0.07480 (17)	0.0486 (7)
H9A	0.8798	0.6520	0.0186	0.058*
H9B	0.9578	0.7319	0.0899	0.058*
C10	1.0165 (3)	0.5485 (3)	0.09490 (19)	0.0594 (8)
H10A	1.0822	0.5690	0.0719	0.071*
H10B	0.9797	0.4692	0.0727	0.071*
C11	1.0719 (3)	0.5339 (4)	0.1833 (2)	0.0648 (9)
H11A	1.1307	0.4646	0.1943	0.078*
H11B	1.1165	0.6101	0.2049	0.078*
C18	0.4656 (3)	0.5554 (3)	0.3928 (2)	0.0633 (9)
H18	0.4018	0.5412	0.4152	0.076*
C21	0.2942 (3)	0.3601 (4)	0.3316 (2)	0.0694 (10)

supplementary materials

H21A	0.3291	0.3303	0.3843	0.104*
H21B	0.2405	0.2967	0.3012	0.104*
H21C	0.2472	0.4354	0.3322	0.104*
C17	0.5525 (4)	0.6499 (3)	0.4229 (2)	0.0674 (10)
H17	0.5468	0.6982	0.4651	0.081*
C16	0.6452 (3)	0.6712 (3)	0.39035 (18)	0.0551 (8)
H16	0.7032	0.7342	0.4106	0.066*
O5	0.3096 (3)	0.3010 (3)	0.13694 (18)	0.1061 (11)
H1	0.3424	0.3299	0.1827	0.159*
H2	0.3649	0.2594	0.1238	0.159*
C22	0.3307 (3)	0.0621 (3)	0.0015 (2)	0.0666 (10)
H22A	0.2609	0.1004	-0.0360	0.100*
H22B	0.3032	0.0219	0.0416	0.100*
H22C	0.3674	0.0005	-0.0243	0.100*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Fe1	0.02801 (19)	0.03099 (18)	0.02628 (18)	-0.00228 (14)	0.01153 (13)	-0.00086 (13)
N1	0.0334 (12)	0.0387 (11)	0.0353 (11)	-0.0007 (9)	0.0108 (9)	0.0018 (9)
O1	0.0425 (11)	0.0466 (10)	0.0377 (10)	-0.0046 (8)	0.0157 (8)	-0.0025 (7)
O3	0.0515 (12)	0.0491 (10)	0.0391 (10)	-0.0053 (9)	0.0221 (9)	-0.0030 (8)
N2	0.0406 (13)	0.0432 (12)	0.0382 (12)	-0.0011 (10)	0.0163 (10)	-0.0013 (9)
C7	0.0384 (15)	0.0438 (14)	0.0351 (14)	0.0024 (11)	0.0136 (12)	0.0046 (10)
C1	0.0351 (15)	0.0369 (13)	0.0386 (14)	0.0047 (10)	0.0101 (12)	-0.0025 (10)
C2	0.0398 (16)	0.0464 (15)	0.0470 (16)	0.0015 (12)	0.0148 (13)	-0.0043 (12)
C20	0.0409 (16)	0.0498 (15)	0.0318 (13)	0.0055 (11)	0.0141 (12)	0.0041 (10)
O4	0.0513 (14)	0.0833 (15)	0.0531 (13)	-0.0119 (11)	0.0289 (11)	0.0000 (11)
C19	0.0444 (18)	0.0620 (18)	0.0408 (16)	0.0010 (14)	0.0169 (13)	0.0047 (13)
O2	0.0712 (16)	0.0618 (13)	0.0629 (14)	-0.0281 (11)	0.0277 (12)	-0.0130 (10)
C8	0.0363 (15)	0.0393 (14)	0.0432 (15)	-0.0015 (11)	0.0129 (12)	0.0038 (11)
C14	0.0414 (16)	0.0463 (15)	0.0359 (14)	0.0002 (12)	0.0101 (12)	-0.0035 (11)
C6	0.0342 (14)	0.0418 (14)	0.0349 (13)	0.0054 (11)	0.0086 (11)	-0.0004 (10)
C4	0.0539 (19)	0.0608 (18)	0.0390 (16)	0.0104 (14)	0.0112 (14)	-0.0096 (13)
C15	0.0470 (17)	0.0469 (15)	0.0340 (14)	0.0011 (12)	0.0143 (12)	-0.0002 (11)
C5	0.0456 (18)	0.0559 (17)	0.0383 (15)	0.0046 (13)	0.0134 (13)	-0.0003 (12)
C13	0.0387 (16)	0.0445 (14)	0.0417 (15)	-0.0055 (11)	0.0143 (12)	-0.0025 (11)
C12	0.0397 (17)	0.0653 (19)	0.0439 (16)	0.0075 (13)	0.0098 (13)	0.0074 (13)
C3	0.0473 (18)	0.0480 (16)	0.0498 (17)	0.0019 (13)	0.0086 (14)	-0.0111 (13)
C9	0.0451 (18)	0.0587 (18)	0.0440 (16)	-0.0118 (13)	0.0162 (14)	0.0028 (12)
C10	0.0436 (19)	0.085 (2)	0.056 (2)	0.0016 (16)	0.0244 (16)	0.0000 (16)
C11	0.0369 (18)	0.098 (3)	0.061 (2)	0.0095 (17)	0.0161 (15)	0.0101 (18)
C18	0.057 (2)	0.093 (3)	0.0510 (19)	0.0029 (18)	0.0322 (17)	-0.0024 (16)
C21	0.046 (2)	0.106 (3)	0.065 (2)	-0.0016 (19)	0.0290 (17)	0.0186 (19)
C17	0.077 (3)	0.084 (2)	0.051 (2)	-0.001 (2)	0.0337 (19)	-0.0176 (17)
C16	0.062 (2)	0.0628 (19)	0.0436 (17)	0.0018 (15)	0.0206 (15)	-0.0097 (13)
O5	0.080 (2)	0.167 (3)	0.080 (2)	-0.021 (2)	0.0366 (17)	-0.0236 (19)
C22	0.055 (2)	0.0543 (19)	0.085 (3)	-0.0180 (15)	0.0124 (19)	-0.0009 (16)

Geometric parameters (Å, °)

Fe1—N2	1.844 (2)	C4—H4	0.9300
Fe1—O1	1.8541 (18)	C15—C16	1.412 (4)
Fe1—O3	1.8623 (19)	C5—H5	0.9300
Fe1—N1	1.864 (2)	C13—C12	1.523 (4)
N1—C7	1.288 (3)	C13—H13	0.9800
N1—C8	1.496 (3)	C12—C11	1.525 (4)
O1—C1	1.317 (3)	C12—H12A	0.9700
O3—C20	1.313 (3)	C12—H12B	0.9700
N2—C14	1.294 (3)	C3—H3	0.9300
N2—C13	1.476 (3)	C9—C10	1.517 (4)
C7—C6	1.429 (4)	C9—H9A	0.9700
C7—H7	0.9300	C9—H9B	0.9700
C1—C6	1.410 (4)	C10—C11	1.525 (5)
C1—C2	1.423 (4)	C10—H10A	0.9700
C2—O2	1.373 (3)	C10—H10B	0.9700
C2—C3	1.376 (4)	C11—H11A	0.9700
C20—C15	1.411 (4)	C11—H11B	0.9700
C20—C19	1.424 (4)	C18—C17	1.394 (5)
O4—C19	1.378 (4)	C18—H18	0.9300
O4—C21	1.425 (4)	C21—H21A	0.9600
C19—C18	1.377 (4)	C21—H21B	0.9600
O2—C22	1.428 (4)	C21—H21C	0.9600
C8—C13	1.519 (4)	C17—C16	1.352 (5)
C8—C9	1.522 (4)	C17—H17	0.9300
C8—H8	0.9800	C16—H16	0.9300
C14—C15	1.431 (4)	O5—H1	0.8500
C14—H14	0.9300	O5—H2	0.8500
C6—C5	1.409 (4)	C22—H22A	0.9600
C4—C5	1.365 (4)	C22—H22B	0.9600
C4—C3	1.397 (4)	C22—H22C	0.9600
N2—Fe1—O1	174.23 (9)	C8—C13—C12	112.4 (2)
N2—Fe1—O3	93.78 (9)	N2—C13—H13	109.8
O1—Fe1—O3	86.17 (8)	C8—C13—H13	109.8
N2—Fe1—N1	85.62 (10)	C12—C13—H13	109.8
O1—Fe1—N1	95.05 (9)	C13—C12—C11	111.0 (2)
O3—Fe1—N1	173.78 (9)	C13—C12—H12A	109.4
C7—N1—C8	120.2 (2)	C11—C12—H12A	109.4
C7—N1—Fe1	125.91 (19)	C13—C12—H12B	109.4
C8—N1—Fe1	113.05 (16)	C11—C12—H12B	109.4
C1—O1—Fe1	126.75 (17)	H12A—C12—H12B	108.0
C20—O3—Fe1	124.53 (17)	C2—C3—C4	120.0 (3)
C14—N2—C13	119.1 (2)	C2—C3—H3	120.0
C14—N2—Fe1	127.8 (2)	C4—C3—H3	120.0
C13—N2—Fe1	113.07 (16)	C10—C9—C8	112.8 (2)
N1—C7—C6	125.7 (2)	C10—C9—H9A	109.0
N1—C7—H7	117.1	C8—C9—H9A	109.0

supplementary materials

C6—C7—H7	117.1	C10—C9—H9B	109.0
O1—C1—C6	124.6 (2)	C8—C9—H9B	109.0
O1—C1—C2	118.5 (2)	H9A—C9—H9B	107.8
C6—C1—C2	116.9 (2)	C9—C10—C11	111.5 (3)
O2—C2—C3	124.5 (3)	C9—C10—H10A	109.3
O2—C2—C1	113.8 (2)	C11—C10—H10A	109.3
C3—C2—C1	121.7 (3)	C9—C10—H10B	109.3
O3—C20—C15	124.8 (3)	C11—C10—H10B	109.3
O3—C20—C19	118.8 (3)	H10A—C10—H10B	108.0
C15—C20—C19	116.4 (2)	C12—C11—C10	110.7 (3)
C19—O4—C21	117.7 (3)	C12—C11—H11A	109.5
C18—C19—O4	124.5 (3)	C10—C11—H11A	109.5
C18—C19—C20	121.2 (3)	C12—C11—H11B	109.5
O4—C19—C20	114.3 (2)	C10—C11—H11B	109.5
C2—O2—C22	118.4 (3)	H11A—C11—H11B	108.1
N1—C8—C13	105.1 (2)	C19—C18—C17	120.9 (3)
N1—C8—C9	116.7 (2)	C19—C18—H18	119.6
C13—C8—C9	111.8 (2)	C17—C18—H18	119.6
N1—C8—H8	107.6	O4—C21—H21A	109.5
C13—C8—H8	107.6	O4—C21—H21B	109.5
C9—C8—H8	107.6	H21A—C21—H21B	109.5
N2—C14—C15	123.6 (3)	O4—C21—H21C	109.5
N2—C14—H14	118.2	H21A—C21—H21C	109.5
C15—C14—H14	118.2	H21B—C21—H21C	109.5
C5—C6—C1	120.4 (2)	C16—C17—C18	119.8 (3)
C5—C6—C7	118.3 (2)	C16—C17—H17	120.1
C1—C6—C7	121.3 (2)	C18—C17—H17	120.1
C5—C4—C3	120.1 (3)	C17—C16—C15	120.7 (3)
C5—C4—H4	120.0	C17—C16—H16	119.6
C3—C4—H4	120.0	C15—C16—H16	119.6
C20—C15—C16	121.0 (3)	H1—O5—H2	107.8
C20—C15—C14	121.1 (2)	O2—C22—H22A	109.5
C16—C15—C14	117.8 (3)	O2—C22—H22B	109.5
C4—C5—C6	120.8 (3)	H22A—C22—H22B	109.5
C4—C5—H5	119.6	O2—C22—H22C	109.5
C6—C5—H5	119.6	H22A—C22—H22C	109.5
N2—C13—C8	104.4 (2)	H22B—C22—H22C	109.5
N2—C13—C12	110.5 (2)		

Hydrogen-bond geometry (\AA , $^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O5—H2 \cdots O1	0.85	2.52	3.102 (4)	126
O5—H2 \cdots O2	0.85	2.15	2.915 (4)	149
O5—H1 \cdots O3	0.85	2.69	3.254 (4)	125
O5—H1 \cdots O4	0.85	2.06	2.898 (4)	169

Fig. 1

