

**(R)-2-[(R)-2,2-Dimethyl-1,3-dioxolan-4-yl]-1,3-oxathiolan-5-one**Qin-Pei Wu,<sup>a\*</sup> Da-Xin Shi,<sup>a</sup> Hao Wang<sup>b</sup> and Qing-Shan Zhang<sup>a</sup><sup>a</sup>School of Chemical Engineering and Environment, Beijing Institute of Technology, Beijing 100081, People's Republic of China, and <sup>b</sup>School of Life Science, Beijing Institute of Technology, Beijing 100081, People's Republic of China  
Correspondence e-mail: qpwu@bit.edu.cn


Received 14 April 2009; accepted 28 May 2009

Key indicators: single-crystal X-ray study;  $T = 293$  K; mean  $\sigma(\text{C}-\text{C}) = 0.005$  Å;  $R$  factor = 0.042;  $wR$  factor = 0.129; data-to-parameter ratio = 14.3.

In the title compound,  $\text{C}_8\text{H}_{12}\text{O}_4\text{S}$ , the two five-membered rings both adopt envelope conformations. In the crystal, weak  $\text{C}-\text{H}\cdots\text{O}$  interactions link neighbouring molecules.

**Related literature**

The title compound is a precursor for the preparation of an important nucleoside drug. For applications of nucleosides in the fields of biology, drugs and chemistry, see: Goodyear *et al.* (2005); Simons (2001); Vittori *et al.* (2006).

**Experimental***Crystal data*

$\text{C}_8\text{H}_{12}\text{O}_4\text{S}$   
 $M_r = 204.24$   
Monoclinic,  $P2_1$   
 $a = 6.5528$  (13) Å  
 $b = 9.4029$  (19) Å  
 $c = 7.9240$  (16) Å  
 $\beta = 106.60$  (3)°  
 $V = 467.89$  (16) Å<sup>3</sup>  
 $Z = 2$   
Mo  $K\alpha$  radiation  
 $\mu = 0.33$  mm<sup>-1</sup>  
 $T = 293$  K  
0.50 × 0.20 × 0.15 mm

*Data collection*

Rigaku Saturn CCD area-detector diffractometer  
Absorption correction: multi-scan (*CrystalClear*; Rigaku/MS, 2005)  
 $T_{\min} = 0.859$ ,  $T_{\max} = 0.952$   
1941 measured reflections  
1705 independent reflections  
1275 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.056$

*Refinement*

$R[F^2 > 2\sigma(F^2)] = 0.042$   
 $wR(F^2) = 0.129$   
 $S = 1.01$   
1705 reflections  
119 parameters  
1 restraint  
H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 0.24$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.20$  e Å<sup>-3</sup>  
Absolute structure: Flack (1983), 593 Friedel pairs  
Flack parameter:  $-0.01$  (13)

**Table 1**

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{C1}-\text{H1A}\cdots\text{O3}^{\text{i}}$	0.97	2.58	3.428 (4)	146
$\text{C1}-\text{H1B}\cdots\text{O2}^{\text{ii}}$	0.97	2.41	3.306 (6)	153
$\text{C3}-\text{H3}\cdots\text{O2}^{\text{iii}}$	0.98	2.55	3.265 (4)	129

Symmetry codes: (i)  $x, y, z - 1$ ; (ii)  $-x + 2, y + \frac{1}{2}, -z + 1$ ; (iii)  $x - 1, y, z$ .

Data collection: *RAPID-AUTO* (Rigaku/MS, 2005); cell refinement: *RAPID-AUTO*; data reduction: *CrystalStructure* (Rigaku/MS, 2005); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXL97*.

The authors thank the National Science Foundation of China (30340070) and the Ministry of Science and Technology of China (2006AA100216) for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: WN2324).

**References**

- Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.  
Goodyear, M. D., Hill, M. L., West, J. P. & Whitehead, A. J. (2005). *Tetrahedron Lett.* **46**, 8535–8538.  
Rigaku/MS (2005). *CrystalClear*, *CrystalStructure* and *RAPID-AUTO*. Rigaku/MS Inc., The Woodlands, Texas, USA.  
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
Simons, C. (2001). *Nucleoside Mimetics, Their Chemistry and Biological Properties*. Amsterdam: Gordon and Breach Science Publisher.  
Vittori, S., Dal Ben, D., Lambertucci, C., Marucci, G., Volpini, R. & Cristalli, G. (2006). *Curr. Med. Chem.* **13**, 3529–3552.

**supplementary materials**

*Acta Cryst.* (2009). E65, o1483 [ doi:10.1107/S160053680902039X ]

**(*R*)-2-[(*R*)-2,2-Dimethyl-1,3-dioxolan-4-yl]-1,3-oxathiolan-5-one**

**Q.-P. Wu, D.-X. Shi, H. Wang and Q.-S. Zhang**

**Comment**

Nucleosides are a very important series of compounds in the fields of biology, drugs and chemistry (Simons, 2001); as an example, lamivudine is used as a drug for HIV and HBV diseases (Goodyear *et al.*, 2005; Vittori *et al.*, 2006). Studies of the synthesis of nucleoside mimetics are essential.

The purpose of this structure determination was to establish the molecular conformation of the title compound obtained by coupling (*R*)-(+)-2,2-dimethyl-1,3-dioxolane-4-carboxaldehyde with 2-mercaptoacetic acid. The chirality at the 2-position (C3) is *R*; this satisfies our requirements for the preparation of corresponding *L*-nucleosides. All bond lengths and bond angles have expected values. The two 5-membered rings both adopt envelope conformations with atoms C3 and C6 at the flap. Three intermolecular C—H $\cdots$ O interactions link neighbouring molecules.

**Experimental**

A solution of (*R*)-(+)-2,2-dimethyl-1,3-dioxolane-4-carboxaldehyde (6.51 g, 50.0 mmol) and 2-mercaptoacetic acid (4.20 ml, 60.0 mmol) in toluene (200 ml) was heated under reflux for 1.5 h. After the reaction mixture was cooled to room temperature, a saturated aqueous solution of NaHCO<sub>3</sub> (30 ml) was added and these two layers were separated. The organic layer was washed with brine, dried (MgSO<sub>4</sub>) and concentrated under reduced pressure. The residue was isolated through short column chromatography on silica gel, which was eluted with EtOAc-petroleum to give the target compound (4.96 g, 48%). m.p. 75–77°C.

50 mg of the final product was dissolved in petroleum ether (5 ml) and the solution was kept at room temperature for 2 days to give colorless single crystals.

**Refinement**

H atoms were included in the riding model approximation, with C—H distances 0.96–0.98 Å, and with  $U_{\text{iso}}(\text{H}) = kU_{\text{eq}}(\text{C})$ , where  $k = 1.5$  for methyl H and 1.2 for all other H atoms.

**Figures**




Fig. 1. Molecular structure of the title compound with displacement ellipsoids drawn at the 30% probability level. Hydrogen atoms are drawn as spheres of arbitrary radius.




Fig. 2. The crystal packing of the title compound, viewed along the *a* axis. Hydrogen bonds are shown as dashed lines.

## (*R*)-2-[(*R*)-2,2-Dimethyl-1,3-dioxolan-4-yl]-1,3-oxathiolan-5-one

### Crystal data

$C_8H_{12}O_4S$

$M_r = 204.24$

Monoclinic,  $P2_1$

$a = 6.5528$  (13) Å

$b = 9.4029$  (19) Å

$c = 7.9240$  (16) Å

$\beta = 106.60$  (3)°

$V = 467.89$  (16) Å<sup>3</sup>

$Z = 2$

$F_{000} = 216$

$D_x = 1.450$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation

$\lambda = 0.71073$  Å

Cell parameters from 1941 reflections

$\theta = 2.7$ – $27.5$ °

$\mu = 0.33$  mm<sup>-1</sup>

$T = 293$  K

Block, colourless

$0.50 \times 0.20 \times 0.15$  mm

### Data collection

Rigaku Saturn CCD area-detector diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

Detector resolution: 10.00 pixels mm<sup>-1</sup>

$T = 293$  K

$\Omega$  scans

Absorption correction: multi-scan (CrystalClear; Rigaku/MSO, 2005)

$T_{\min} = 0.859$ ,  $T_{\max} = 0.952$

1941 measured reflections

1705 independent reflections

1275 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.056$

$\theta_{\text{max}} = 27.5$ °

$\theta_{\text{min}} = 2.7$ °

$h = -8 \rightarrow 8$

$k = -12 \rightarrow 11$

$l = -10 \rightarrow 10$

### Refinement

Refinement on  $F^2$

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.042$

$wR(F^2) = 0.129$

$S = 1.01$

1705 reflections

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.088P)^2]$$

where  $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\text{max}} < 0.001$

$\Delta\rho_{\text{max}} = 0.24$  e Å<sup>-3</sup>

$\Delta\rho_{\text{min}} = -0.20$  e Å<sup>-3</sup>

119 parameters Extinction correction: SHELXL97 (Sheldrick, 2008),  
 $F_c^* = kFc[1+0.001xFc^2\lambda^3/\sin(2\theta)]^{-1/4}$   
 1 restraint Extinction coefficient: 0.102 (15)  
 Primary atom site location: structure-invariant direct methods Absolute structure: Flack (1983), 593 Friedel pairs  
 Secondary atom site location: difference Fourier map Flack parameter: -0.01 (13)

*Special details*

**Experimental.** <sup>1</sup>H NMR(CDC1<sub>3</sub>,P.P.M.): 1.41 (d, 6 H), 3.58(d, 4 H), 3.77 (d, 1 H), 3.92 (dd, 1 H), 4.12 (dd, 1 H), 4.35 (m, 1 H), 5.45(d, 1 H).

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ - factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{iso}^*/U_{eq}$
S1	0.57296 (17)	0.19608 (12)	0.31212 (10)	0.0733 (4)
O1	0.7766 (4)	0.0255 (3)	0.5605 (3)	0.0514 (6)
O2	1.1103 (4)	0.0457 (3)	0.5630 (4)	0.0734 (8)
O3	0.8026 (4)	0.1607 (2)	0.9331 (3)	0.0560 (6)
O4	0.7352 (4)	0.2936 (2)	0.6920 (3)	0.0482 (5)
C1	0.8546 (6)	0.1980 (5)	0.3737 (4)	0.0612 (9)
H1A	0.9049	0.1803	0.2718	0.073*
H1B	0.9075	0.2899	0.4225	0.073*
C2	0.9307 (6)	0.0859 (4)	0.5061 (4)	0.0502 (8)
C3	0.5800 (5)	0.1011 (4)	0.5101 (4)	0.0509 (8)
H3	0.4622	0.0328	0.4851	0.061*
C4	0.5669 (5)	0.1947 (4)	0.6584 (4)	0.0486 (7)
H4	0.4302	0.2451	0.6269	0.058*
C5	0.5987 (5)	0.1192 (5)	0.8312 (4)	0.0567 (9)
H5A	0.4914	0.1477	0.8871	0.068*
H5B	0.5915	0.0169	0.8144	0.068*
C6	0.8418 (5)	0.2954 (4)	0.8756 (4)	0.0486 (8)
C7	1.0731 (6)	0.3117 (5)	0.9004 (6)	0.0742 (11)
H7A	1.1010	0.4044	0.8614	0.111*
H7B	1.1201	0.2405	0.8330	0.111*
H7C	1.1483	0.3006	1.0228	0.111*
C8	0.7487 (7)	0.4089 (5)	0.9608 (5)	0.0743 (12)
H8A	0.7778	0.5000	0.9182	0.111*
H8B	0.8106	0.4046	1.0861	0.111*

# supplementary materials

H8C                    0.5975                    0.3954                    0.9332                    0.111\*

## Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
S1	0.0745 (6)	0.0908 (9)	0.0459 (4)	-0.0108 (6)	0.0034 (4)	0.0026 (5)
O1	0.0541 (13)	0.0404 (12)	0.0659 (14)	-0.0020 (10)	0.0269 (11)	-0.0011 (11)
O2	0.0606 (17)	0.0628 (18)	0.107 (2)	0.0052 (14)	0.0402 (16)	-0.0045 (16)
O3	0.0649 (15)	0.0446 (14)	0.0563 (12)	0.0047 (11)	0.0139 (10)	0.0108 (11)
O4	0.0566 (12)	0.0465 (12)	0.0411 (10)	-0.0113 (11)	0.0132 (9)	0.0000 (9)
C1	0.082 (2)	0.053 (2)	0.0589 (18)	-0.012 (2)	0.0367 (17)	-0.0009 (17)
C2	0.059 (2)	0.0397 (17)	0.0597 (18)	-0.0016 (15)	0.0293 (16)	-0.0084 (14)
C3	0.0457 (16)	0.0508 (19)	0.0557 (17)	-0.0117 (15)	0.0134 (14)	-0.0050 (16)
C4	0.0389 (14)	0.0522 (19)	0.0563 (15)	-0.0025 (15)	0.0163 (12)	0.0032 (17)
C5	0.0558 (19)	0.060 (2)	0.062 (2)	-0.0076 (17)	0.0287 (17)	0.0027 (16)
C6	0.0548 (19)	0.0429 (18)	0.0453 (16)	0.0041 (15)	0.0100 (14)	0.0022 (13)
C7	0.058 (2)	0.065 (3)	0.088 (3)	-0.0082 (19)	0.004 (2)	-0.004 (2)
C8	0.106 (3)	0.057 (2)	0.063 (2)	0.015 (2)	0.027 (2)	-0.0093 (18)

## Geometric parameters ( $\text{\AA}$ , $^\circ$ )

S1—C1	1.769 (4)	C3—H3	0.9800
S1—C3	1.795 (4)	C4—C5	1.504 (5)
O1—C2	1.333 (4)	C4—H4	0.9800
O1—C3	1.425 (4)	C5—H5A	0.9700
O2—C2	1.195 (4)	C5—H5B	0.9700
O3—C6	1.395 (4)	C6—C8	1.483 (5)
O3—C5	1.405 (4)	C6—C7	1.480 (5)
O4—C4	1.408 (4)	C7—H7A	0.9600
O4—C6	1.423 (4)	C7—H7B	0.9600
C1—C2	1.470 (5)	C7—H7C	0.9600
C1—H1A	0.9700	C8—H8A	0.9600
C1—H1B	0.9700	C8—H8B	0.9600
C3—C4	1.490 (5)	C8—H8C	0.9600
C1—S1—C3	89.97 (16)	O3—C5—C4	104.7 (3)
C2—O1—C3	113.9 (3)	O3—C5—H5A	110.8
C6—O3—C5	107.4 (3)	C4—C5—H5A	110.8
C4—O4—C6	109.3 (2)	O3—C5—H5B	110.8
C2—C1—S1	107.7 (3)	C4—C5—H5B	110.8
C2—C1—H1A	110.2	H5A—C5—H5B	108.9
S1—C1—H1A	110.2	O3—C6—O4	103.9 (3)
C2—C1—H1B	110.2	O3—C6—C8	111.5 (3)
S1—C1—H1B	110.2	O4—C6—C8	109.2 (3)
H1A—C1—H1B	108.5	O3—C6—C7	109.1 (3)
O2—C2—O1	119.9 (3)	O4—C6—C7	108.8 (3)
O2—C2—C1	126.4 (3)	C8—C6—C7	113.8 (4)
O1—C2—C1	113.7 (3)	C6—C7—H7A	109.5
O1—C3—C4	109.0 (3)	C6—C7—H7B	109.5

O1—C3—S1	106.8 (2)	H7A—C7—H7B	109.5
C4—C3—S1	113.7 (3)	C6—C7—H7C	109.5
O1—C3—H3	109.1	H7A—C7—H7C	109.5
C4—C3—H3	109.1	H7B—C7—H7C	109.5
S1—C3—H3	109.1	C6—C8—H8A	109.5
O4—C4—C3	108.7 (2)	C6—C8—H8B	109.5
O4—C4—C5	104.0 (3)	H8A—C8—H8B	109.5
C3—C4—C5	114.5 (3)	C6—C8—H8C	109.5
O4—C4—H4	109.8	H8A—C8—H8C	109.5
C3—C4—H4	109.8	H8B—C8—H8C	109.5
C5—C4—H4	109.8		
C3—S1—C1—C2	19.2 (3)	S1—C3—C4—O4	-57.3 (3)
C3—O1—C2—O2	168.2 (3)	O1—C3—C4—C5	-54.1 (3)
C3—O1—C2—C1	-13.0 (4)	S1—C3—C4—C5	-173.1 (2)
S1—C1—C2—O2	171.1 (3)	C6—O3—C5—C4	28.4 (3)
S1—C1—C2—O1	-7.6 (4)	O4—C4—C5—O3	-11.8 (4)
C2—O1—C3—C4	-96.1 (3)	C3—C4—C5—O3	106.7 (3)
C2—O1—C3—S1	27.2 (3)	C5—O3—C6—O4	-33.7 (3)
C1—S1—C3—O1	-26.0 (3)	C5—O3—C6—C8	83.8 (3)
C1—S1—C3—C4	94.2 (3)	C5—O3—C6—C7	-149.6 (3)
C6—O4—C4—C3	-130.9 (3)	C4—O4—C6—O3	25.9 (3)
C6—O4—C4—C5	-8.5 (3)	C4—O4—C6—C8	-93.2 (4)
O1—C3—C4—O4	61.7 (3)	C4—O4—C6—C7	142.0 (3)

*Hydrogen-bond geometry* (Å, °)

<i>D</i> —H $\cdots$ <i>A</i>	<i>D</i> —H	H $\cdots$ <i>A</i>	<i>D</i> $\cdots$ <i>A</i>	<i>D</i> —H $\cdots$ <i>A</i>
C1—H1A $\cdots$ O3 <sup>i</sup>	0.97	2.58	3.428 (4)	146
C1—H1B $\cdots$ O2 <sup>ii</sup>	0.97	2.41	3.306 (6)	153
C3—H3 $\cdots$ O2 <sup>iii</sup>	0.98	2.55	3.265 (4)	129

Symmetry codes: (i)  $x, y, z-1$ ; (ii)  $-x+2, y+1/2, -z+1$ ; (iii)  $x-1, y, z$ .

Fig. 1






Fig. 2

