

2,3-Diaminopyridinium 4-hydroxybenzoate

Hoong-Kun Fun*‡ and Kasthuri Balasubramani

X-ray Crystallography Unit, School of Physics, Universiti Sains Malaysia, 11800 Universiti Sains Malaysia, Penang, Malaysia
Correspondence e-mail: hkfun@usm.my


Received 22 May 2009; accepted 2 June 2009

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.001$ Å; R factor = 0.049; wR factor = 0.138; data-to-parameter ratio = 24.6.

In the title compound, $\text{C}_5\text{H}_8\text{N}_3^+\cdot\text{C}_7\text{H}_5\text{O}_3^-$, the pyridine N atom is protonated. In the 4-hydroxybenzoate anion, the carboxylate group is twisted slightly out of the benzene ring plane by an angle of 3.77 (5)°. The protonated N atom and one of the two amino groups are hydrogen-bonded to the 4-hydroxybenzoate anion through a pair of $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds, forming an $R_2^2(8)$ ring motif. The crystal structure is further stabilized by $\text{O}-\text{H}\cdots\text{O}$ and $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds and $\pi-\pi$ interactions involving the pyridinium rings [centroid-centroid distance of 3.6277 (5) Å], leading to the formation of a three-dimensional network.

Related literature

For general background to substituted pyridines, see: Pozharski *et al.* (1997); Katritzky *et al.* (1996). For details of hydrogen bonding, see: Jeffrey & Saenger (1991); Jeffrey (1997); Scheiner (1997). For hydrogen-bond motifs, see: Bernstein *et al.* (1995). For bond-length data, see: Allen *et al.* (1987). For the stability of the temperature controller used in the data collection, see: Cosier & Glazer (1986).



‡ Thomson Reuters ResearcherID: A-3561-2009.

Experimental

Crystal data

$\text{C}_5\text{H}_8\text{N}_3^+\cdot\text{C}_7\text{H}_5\text{O}_3^-$
 $M_r = 247.25$
Monoclinic, $P2_1/c$
 $a = 10.2915$ (2) Å
 $b = 11.4946$ (2) Å
 $c = 11.0921$ (2) Å
 $\beta = 112.644$ (1)°

$V = 1211.01$ (4) Å³
 $Z = 4$
Mo $K\alpha$ radiation
 $\mu = 0.10$ mm⁻¹
 $T = 100$ K
 $0.51 \times 0.39 \times 0.14$ mm

Data collection

Bruker SMART APEXII CCD
area-detector diffractometer
Absorption correction: multi-scan
(*SADABS*; Bruker, 2005)
 $T_{\min} = 0.950$, $T_{\max} = 0.986$

25821 measured reflections
5296 independent reflections
4257 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.034$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.049$
 $wR(F^2) = 0.138$
 $S = 1.04$
5296 reflections
215 parameters

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.65$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.19$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N1}-\text{H1N1}\cdots\text{O2}$	0.89 (2)	1.903 (15)	2.7874 (9)	173 (1)
$\text{N2}-\text{H2N2}\cdots\text{O3}$	0.89 (1)	1.898 (14)	2.7843 (9)	176 (1)
$\text{N2}-\text{H1N2}\cdots\text{O2}^i$	0.87 (1)	2.014 (15)	2.8689 (9)	168 (1)
$\text{N3}-\text{H1N3}\cdots\text{O2}^i$	0.91 (2)	2.071 (16)	2.9790 (10)	174 (1)
$\text{N3}-\text{H2N3}\cdots\text{O3}^{ii}$	0.89 (2)	2.057 (15)	2.9285 (10)	166 (1)
$\text{O1}-\text{H1O1}\cdots\text{O3}^{iii}$	0.90 (2)	1.775 (19)	2.6595 (8)	168 (2)
$\text{C2}-\text{H2}\cdots\text{O3}^{iii}$	1.00 (1)	2.500 (14)	3.2104 (10)	128 (1)

Symmetry codes: (i) $x, -y + \frac{3}{2}, z - \frac{1}{2}$; (ii) $-x + 2, y + \frac{1}{2}, -z + \frac{1}{2}$; (iii) $-x + 1, y - \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: *APEX2* (Bruker, 2005); cell refinement: *SAINT* (Bruker, 2005); data reduction: *SAINT*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL* and *PLATON* (Spek, 2009).

HKF and KB thank the Malaysian Government and Universiti Sains Malaysia for the Science Fund grant No. 305/PFIZIK/613312. KB thanks Universiti Sains Malaysia for a post-doctoral research fellowship. HKF also thanks Universiti Sains Malaysia for the Research University Golden Goose grant No. 1001/PFIZIK/811012.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CI2813).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–S19.
Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
Bruker (2005). *APEX2*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
Cosier, J. & Glazer, A. M. (1986). *J. Appl. Cryst.* **19**, 105–107.

- Jeffrey, G. A. (1997). *An Introduction to Hydrogen Bonding*. Oxford University Press.
- Jeffrey, G. A. & Saenger, W. (1991). *Hydrogen Bonding in Biological Structures*. Berlin: Springer.
- Katritzky, A. R., Rees, C. W. & Scriven, E. F. V. (1996). *Comprehensive Heterocyclic Chemistry II*. Oxford: Pergamon Press.
- Pozharski, A. F., Soldatenkov, A. T. & Katritzky, A. R. (1997). *Heterocycles in Life and Society*. New York: Wiley.
- Scheiner, S. (1997). *Hydrogen Bonding, A Theoretical Perspective*. Oxford University Press.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

supplementary materials

Acta Cryst. (2009). E65, o1496-o1497 [doi:10.1107/S1600536809020832]

2,3-Diaminopyridinium 4-hydroxybenzoate

H.-K. Fun and K. Balasubramani

Comment

Pyridine and its derivatives play an important role in heterocyclic chemistry (Pozharski *et al.*, 1997; Katritzky *et al.*, 1996). Pyridine and its substituted derivatives are often involved in hydrogen-bond interactions (Jeffrey & Saenger, 1991; Jeffrey, 1997; Scheiner, 1997). Since our aim is to study some interesting hydrogen-bonding interactions, the crystal structure of the title compound is presented here.

The asymmetric unit of the title compound (Fig 1), contains one 2,3-diaminopyridinium cation and one 4-hydroxybenzoate anion. The bond lengths (Allen *et al.*, 1987) and angles are normal. The 2,3-diaminopyridinium cation is planar to within ± 0.015 (1) Å. The protonation of N1 atom resulted in a slight increase in the C8—N1—C12 angle [123.47 (7)°]. In the anion, the carboxylate group is twisted slightly away from the attached ring; the dihedral angle between C1-C6 and O2/O3/C7/C6 planes is 3.77 (5)°.

In the crystal packing (Fig. 2), the protonated N1 atom and the 2-amino group (N2) is hydrogen-bonded to the carboxylate oxygen atoms (O2 and O3) *via* a pair of N—H \cdots O hydrogen bonds forming a ring motif $R_2^2(8)$ (Bernstein *et al.*, 1995). The amino groups (N2 and N3) are involved in N—H \cdots O hydrogen bonding interactions to form a $R_1^2(7)$ ring motif. The hydroxyl group hydrogen atom is also hydrogen-bonded to the carboxylate oxygen atom through O—H \cdots O hydrogen bonds. Moreover O—H \cdots O and C—H \cdots O hydrogen bonds form a $R_1^2(6)$ ring motif (Table 1 and Fig 2). The crystal structure is further stabilized by π - π stacking interactions between the pyridinium rings (C8—C12/N1) at (x, y, z) and (2-x, 2-y, 1-z), with a centroid-to-centroid distance of 3.6277 (5) Å.

Experimental

Hot methanol solutions (20 ml) of 2,3-diaminopyridine (27 mg, Aldrich) and 4-hydroxybenzoic acid (35 mg, Merck) were mixed and warmed over a heating magnetic stirrer for 5 minutes. The resulting solution was allowed to cool slowly at room temperature. Crystals of the title compound appeared from the mother liquor after a few days.

Refinement

All the H atoms were located in a difference Fourier map and allowed to refine freely [N—H = 0.86–0.91 Å, C—H = 0.95–1.01 (15)Å and O—H = 0.89 (18) Å]

Figures




Fig. 1. The molecular structure of the title compound, showing 50% probability displacement ellipsoids and the atom numbering scheme. Dashed lines indicate hydrogen bonds.





Fig. 2. Crystal packing of the title compound, showing a part of the three-dimensional network. Hydrogen bonds are shown as dashed lines.

(I)

Crystal data



$$M_r = 247.25$$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

$$a = 10.2915 (2) \text{ \AA}$$

$$b = 11.4946 (2) \text{ \AA}$$

$$c = 11.0921 (2) \text{ \AA}$$

$$\beta = 112.644 (1)^\circ$$

$$V = 1211.01 (4) \text{ \AA}^3$$

$$Z = 4$$

$$F_{000} = 520$$

$$D_x = 1.356 \text{ Mg m}^{-3}$$

Mo $K\alpha$ radiation

$$\lambda = 0.71073 \text{ \AA}$$

Cell parameters from 9979 reflections

$$\theta = 2.7\text{--}37.8^\circ$$

$$\mu = 0.10 \text{ mm}^{-1}$$

$$T = 100 \text{ K}$$

Plate, brown

$$0.51 \times 0.39 \times 0.14 \text{ mm}$$

Data collection

Bruker SMART APEXII CCD area-detector diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$$T = 100 \text{ K}$$

φ and ω scans

Absorption correction: multi-scan (SADABS; Bruker, 2005)

$$T_{\min} = 0.950, T_{\max} = 0.986$$

25821 measured reflections

5296 independent reflections

4257 reflections with $I > 2\sigma(I)$

$$R_{\text{int}} = 0.034$$

$$\theta_{\max} = 35.0^\circ$$

$$\theta_{\min} = 2.1^\circ$$

$$h = -16 \rightarrow 16$$

$$k = -16 \rightarrow 18$$

$$l = -17 \rightarrow 17$$

Refinement

Refinement on F^2

Least-squares matrix: full

$$R[F^2 > 2\sigma(F^2)] = 0.049$$

$$wR(F^2) = 0.138$$

$$S = 1.04$$

5296 reflections

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent and constrained refinement

$$w = 1/[\sigma^2(F_o^2) + (0.0817P)^2 + 0.1673P]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} = 0.001$$

$$\Delta\rho_{\max} = 0.65 \text{ e \AA}^{-3}$$

215 parameters

$$\Delta\rho_{\min} = -0.18 \text{ e } \text{\AA}^{-3}$$

Primary atom site location: structure-invariant direct methods

Extinction correction: none

Special details

Experimental. The crystal was placed in the cold stream of an Oxford Cyrosystems Cobra open-flow nitrogen cryostat (Cosier & Glazer, 1986) operating at 100.0 (1) K.

Geometry. All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R- factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.45467 (8)	0.16900 (6)	0.43336 (6)	0.02563 (15)
O2	0.84786 (7)	0.61461 (5)	0.52034 (5)	0.02074 (13)
O3	0.73136 (6)	0.62156 (5)	0.30541 (5)	0.01784 (12)
N1	1.01845 (8)	0.79427 (6)	0.49674 (6)	0.01857 (14)
N2	0.87767 (8)	0.81703 (7)	0.27844 (7)	0.02195 (15)
N3	1.05767 (9)	0.99458 (8)	0.25977 (7)	0.02666 (17)
C1	0.57827 (9)	0.42070 (7)	0.30431 (7)	0.01757 (14)
C2	0.50323 (9)	0.32084 (7)	0.30618 (7)	0.01879 (15)
C3	0.52248 (9)	0.26810 (7)	0.42538 (7)	0.01899 (15)
C4	0.61617 (11)	0.31656 (8)	0.54174 (8)	0.02518 (18)
C5	0.69083 (10)	0.41624 (8)	0.53881 (7)	0.02324 (17)
C6	0.67313 (8)	0.46970 (7)	0.41995 (7)	0.01604 (14)
C7	0.75521 (8)	0.57533 (6)	0.41592 (7)	0.01540 (14)
C8	0.99181 (9)	0.84838 (7)	0.38181 (7)	0.01678 (14)
C9	1.08674 (9)	0.93716 (7)	0.37590 (7)	0.01803 (15)
C10	1.20049 (9)	0.96332 (8)	0.48910 (8)	0.02092 (16)
C11	1.22331 (9)	0.90438 (8)	0.60700 (8)	0.02314 (17)
C12	1.13117 (10)	0.82008 (8)	0.60867 (8)	0.02224 (17)
H1	0.5647 (15)	0.4565 (11)	0.2226 (13)	0.031 (3)*
H2	0.4376 (14)	0.2854 (12)	0.2229 (13)	0.029 (3)*
H4	0.6280 (18)	0.2759 (15)	0.6232 (17)	0.051 (4)*
H5	0.7615 (17)	0.4479 (13)	0.6225 (15)	0.039 (4)*
H10	1.2669 (14)	1.0227 (12)	0.4848 (12)	0.028 (3)*
H11	1.3091 (16)	0.9246 (13)	0.6883 (14)	0.039 (4)*
H12	1.1341 (17)	0.7745 (14)	0.6809 (15)	0.043 (4)*
H1N1	0.9588 (15)	0.7391 (12)	0.4987 (13)	0.031 (3)*
H1N2	0.8601 (17)	0.8454 (12)	0.2011 (15)	0.033 (3)*

supplementary materials

H2N2	0.8275 (15)	0.7565 (12)	0.2854 (13)	0.027 (3)*
H1N3	0.9924 (17)	0.9660 (13)	0.1840 (15)	0.040 (4)*
H2N3	1.1284 (16)	1.0365 (12)	0.2548 (14)	0.034 (3)*
H1O1	0.3995 (19)	0.1445 (15)	0.3531 (18)	0.051 (5)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0299 (3)	0.0248 (3)	0.0172 (3)	-0.0123 (2)	0.0034 (2)	0.0013 (2)
O2	0.0238 (3)	0.0217 (3)	0.0141 (2)	-0.0059 (2)	0.0044 (2)	-0.00153 (19)
O3	0.0188 (3)	0.0178 (3)	0.0149 (2)	0.0002 (2)	0.00434 (19)	0.00308 (19)
N1	0.0200 (3)	0.0194 (3)	0.0148 (3)	-0.0020 (2)	0.0051 (2)	0.0018 (2)
N2	0.0232 (3)	0.0234 (3)	0.0154 (3)	-0.0072 (3)	0.0031 (2)	0.0021 (2)
N3	0.0284 (4)	0.0324 (4)	0.0171 (3)	-0.0118 (3)	0.0065 (3)	0.0037 (3)
C1	0.0190 (3)	0.0187 (3)	0.0131 (3)	-0.0014 (3)	0.0041 (2)	0.0005 (2)
C2	0.0198 (4)	0.0205 (3)	0.0131 (3)	-0.0037 (3)	0.0031 (2)	-0.0007 (2)
C3	0.0202 (4)	0.0199 (3)	0.0150 (3)	-0.0046 (3)	0.0047 (3)	-0.0001 (2)
C4	0.0316 (5)	0.0273 (4)	0.0130 (3)	-0.0114 (3)	0.0046 (3)	0.0008 (3)
C5	0.0286 (4)	0.0251 (4)	0.0131 (3)	-0.0099 (3)	0.0049 (3)	-0.0015 (3)
C6	0.0173 (3)	0.0167 (3)	0.0131 (3)	-0.0016 (2)	0.0048 (2)	-0.0007 (2)
C7	0.0167 (3)	0.0152 (3)	0.0140 (3)	0.0013 (2)	0.0055 (2)	-0.0002 (2)
C8	0.0177 (3)	0.0174 (3)	0.0143 (3)	-0.0003 (3)	0.0052 (2)	0.0001 (2)
C9	0.0183 (3)	0.0204 (3)	0.0155 (3)	-0.0023 (3)	0.0066 (2)	0.0000 (2)
C10	0.0192 (4)	0.0249 (4)	0.0178 (3)	-0.0044 (3)	0.0062 (3)	-0.0010 (3)
C11	0.0200 (4)	0.0294 (4)	0.0166 (3)	-0.0029 (3)	0.0032 (3)	-0.0007 (3)
C12	0.0230 (4)	0.0262 (4)	0.0145 (3)	-0.0014 (3)	0.0038 (3)	0.0023 (3)

Geometric parameters (\AA , $^\circ$)

O1—C3	1.3565 (10)	C2—C3	1.3976 (11)
O1—H1O1	0.897 (18)	C2—H2	0.997 (13)
O2—C7	1.2666 (9)	C3—C4	1.3956 (11)
O3—C7	1.2702 (9)	C4—C5	1.3869 (12)
N1—C8	1.3484 (10)	C4—H4	0.982 (17)
N1—C12	1.3661 (11)	C5—C6	1.4014 (11)
N1—H1N1	0.888 (15)	C5—H5	1.001 (15)
N2—C8	1.3369 (10)	C6—C7	1.4896 (11)
N2—H1N2	0.869 (15)	C8—C9	1.4316 (11)
N2—H2N2	0.887 (14)	C9—C10	1.3805 (11)
N3—C9	1.3739 (10)	C10—C11	1.4100 (12)
N3—H1N3	0.911 (16)	C10—H10	0.981 (14)
N3—H2N3	0.892 (15)	C11—C12	1.3608 (13)
C1—C2	1.3881 (11)	C11—H11	1.017 (15)
C1—C6	1.3969 (10)	C12—H12	0.948 (15)
C1—H1	0.955 (13)		
C3—O1—H1O1	110.1 (11)	C4—C5—H5	119.3 (8)
C8—N1—C12	123.47 (7)	C6—C5—H5	119.9 (9)
C8—N1—H1N1	117.5 (9)	C1—C6—C5	118.61 (7)

C12—N1—H1N1	119.1 (9)	C1—C6—C7	120.33 (6)
C8—N2—H1N2	121.7 (10)	C5—C6—C7	121.05 (7)
C8—N2—H2N2	119.1 (9)	O2—C7—O3	122.08 (7)
H1N2—N2—H2N2	118.2 (13)	O2—C7—C6	119.94 (6)
C9—N3—H1N3	120.6 (10)	O3—C7—C6	117.96 (6)
C9—N3—H2N3	115.4 (9)	N2—C8—N1	118.57 (7)
H1N3—N3—H2N3	117.8 (13)	N2—C8—C9	122.86 (7)
C2—C1—C6	121.05 (7)	N1—C8—C9	118.57 (7)
C2—C1—H1	119.4 (8)	N3—C9—C10	123.40 (8)
C6—C1—H1	119.5 (8)	N3—C9—C8	118.64 (7)
C1—C2—C3	119.72 (7)	C10—C9—C8	117.92 (7)
C1—C2—H2	120.4 (8)	C9—C10—C11	121.27 (8)
C3—C2—H2	119.9 (8)	C9—C10—H10	118.1 (8)
O1—C3—C4	117.67 (7)	C11—C10—H10	120.6 (8)
O1—C3—C2	122.47 (7)	C12—C11—C10	119.12 (8)
C4—C3—C2	119.85 (7)	C12—C11—H11	121.5 (8)
C5—C4—C3	119.99 (7)	C10—C11—H11	119.4 (8)
C5—C4—H4	122.8 (10)	C11—C12—N1	119.65 (7)
C3—C4—H4	117.2 (10)	C11—C12—H12	127.5 (10)
C4—C5—C6	120.78 (7)	N1—C12—H12	112.8 (10)
C6—C1—C2—C3	-0.01 (13)	C5—C6—C7—O3	-178.02 (8)
C1—C2—C3—O1	-178.47 (8)	C12—N1—C8—N2	-179.40 (8)
C1—C2—C3—C4	0.46 (14)	C12—N1—C8—C9	0.41 (12)
O1—C3—C4—C5	178.40 (9)	N2—C8—C9—N3	1.55 (13)
C2—C3—C4—C5	-0.57 (15)	N1—C8—C9—N3	-178.25 (8)
C3—C4—C5—C6	0.24 (15)	N2—C8—C9—C10	179.31 (8)
C2—C1—C6—C5	-0.32 (13)	N1—C8—C9—C10	-0.49 (12)
C2—C1—C6—C7	178.59 (7)	N3—C9—C10—C11	177.91 (9)
C4—C5—C6—C1	0.21 (14)	C8—C9—C10—C11	0.26 (13)
C4—C5—C6—C7	-178.70 (9)	C9—C10—C11—C12	0.06 (14)
C1—C6—C7—O2	-175.21 (8)	C10—C11—C12—N1	-0.16 (14)
C5—C6—C7—O2	3.67 (12)	C8—N1—C12—C11	-0.08 (13)
C1—C6—C7—O3	3.09 (11)		

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N1—H1N1 \cdots O2	0.89 (2)	1.903 (15)	2.7874 (9)	173 (1)
N2—H2N2 \cdots O3	0.89 (1)	1.898 (14)	2.7843 (9)	176 (1)
N2—H1N2 \cdots O2 ⁱ	0.87 (1)	2.014 (15)	2.8689 (9)	168 (1)
N3—H1N3 \cdots O2 ⁱ	0.91 (2)	2.071 (16)	2.9790 (10)	174 (1)
N3—H2N3 \cdots O3 ⁱⁱ	0.89 (2)	2.057 (15)	2.9285 (10)	166 (1)
O1—H1O1 \cdots O3 ⁱⁱⁱ	0.90 (2)	1.775 (19)	2.6595 (8)	168 (2)
C2—H2 \cdots O3 ⁱⁱⁱ	1.00 (1)	2.500 (14)	3.2104 (10)	128 (1)

Symmetry codes: (i) $x, -y+3/2, z-1/2$; (ii) $-x+2, y+1/2, -z+1/2$; (iii) $-x+1, y-1/2, -z+1/2$.

Fig. 1




Fig. 2

