

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

***N'*-(5-Chloro-2-hydroxybenzylidene)-nicotinohydrazide**

Chong-Gui Ren

Department of Chemistry and Chemical Engineering, Zaozhuang University,
Zaozhuang Shandong 277160, People's Republic of China
Correspondence e-mail: renchonggui@163.com


Received 25 May 2009; accepted 2 June 2009

Key indicators: single-crystal X-ray study; $T = 298$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å;
R factor = 0.049; wR factor = 0.115; data-to-parameter ratio = 15.2.

There are two independent Schiff base molecules in the asymmetric unit of the title compound, $\text{C}_{13}\text{H}_{10}\text{ClN}_3\text{O}_2$. The dihedral angles between the benzene and pyridine rings are 12.8 (2) and 1.9 (2)° in the two molecules. Intramolecular $\text{O}-\text{H}\cdots\text{N}$ hydrogen bonds are observed. Molecules are linked into centrosymmetric $R_4^2(26)$ motifs by $\text{N}-\text{H}\cdots\text{O}$ and $\text{N}-\text{H}\cdots\text{N}$ interactions.

Related literature

For the biological properties of Schiff base compounds, see: Jeewoth *et al.* (1999); Ren *et al.* (2002); Eltayeb *et al.* (2008); Sinha *et al.* (2008). For metal complexes of Schiff base compounds, see: Shivakumar *et al.* (2008); Prabhakaran *et al.* (2006); Dhar *et al.* (2005). For related structures, see: Cui *et al.* (2007); Jing *et al.* (2007); Ma *et al.* (2008); Salhin *et al.* (2007); Lin *et al.* (2007); Alhadi *et al.* (2008); Xue *et al.* (2008); Wang *et al.* (2008); Lu (2008); Diao *et al.* (2008); Qiu (2009); Mohd Lair *et al.* (2009*a,b*). For reference structural data, see: Allen *et al.* (1987). For hydrogen-bond motifs, see: Bernstein *et al.* (1995).



Experimental

Crystal data

$\text{C}_{13}\text{H}_{10}\text{ClN}_3\text{O}_2$
 $M_r = 275.69$
Monoclinic, $P2_1/n$
 $a = 9.792$ (2) Å
 $b = 23.358$ (3) Å
 $c = 10.926$ (2) Å
 $\beta = 96.848$ (2)°

$V = 2481.2$ (8) Å³
 $Z = 8$
Mo $K\alpha$ radiation
 $\mu = 0.31$ mm⁻¹
 $T = 298$ K
 $0.30 \times 0.30 \times 0.27$ mm

Data collection

Bruker SMART CCD area-detector diffractometer
Absorption correction: multi-scan (*SADABS*; Sheldrick, 1996)
 $T_{\min} = 0.913$, $T_{\max} = 0.921$
14387 measured reflections
5342 independent reflections
3193 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.042$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.049$
 $wR(F^2) = 0.115$
 $S = 1.01$
5342 reflections
351 parameters
2 restraints

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.22$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.24$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{O1}-\text{H1}\cdots\text{N1}$	0.82	1.88	2.594 (2)	145
$\text{O3}-\text{H3}\cdots\text{N4}$	0.82	1.82	2.538 (2)	146
$\text{N5}-\text{H5}\cdots\text{N3}^i$	0.891 (10)	2.109 (11)	2.991 (3)	171 (2)
$\text{N2}-\text{H2}\cdots\text{O3}$	0.892 (10)	2.097 (10)	2.984 (2)	173 (2)

Symmetry code: (i) $-x + 1, -y, -z + 2$.

Data collection: *SMART* (Bruker, 2002); cell refinement: *SAINTE* (Bruker, 2002); data reduction: *SAINTE*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXL97*.

The author acknowledges Zaozhuang University for funding this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BX2215).

References

- Alhadi, A. A., Ali, H. M., Puvaneswary, S., Robinson, W. T. & Ng, S. W. (2008). *Acta Cryst.* **E64**, o1584.
Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
Bernstein, J., Davis, R. E., Shimon, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
Bruker (2002). *SAINTE* and *SMART*. Bruker AXS Inc., Madison, Wisconsin, USA.
Cui, J.-C., Pan, Q.-X., Yin, H.-D. & Qiao, Y.-L. (2007). *Acta Cryst.* **E63**, o2633.
Dhar, S., Nethaji, M. & Chakravarty, A. R. (2005). *Inorg. Chim. Acta*, **358**, 2437–2444.
Diao, Y.-P., Huang, S.-S., Zhang, H.-L. & Kang, T.-G. (2008). *Z. Kristallogr. New Cryst. Struct.* **223**, 165–166.
Eltayeb, N. E., Teoh, S. G., Chantrapromma, S., Fun, H.-K. & Adnan, R. (2008). *Acta Cryst.* **E64**, o576–o577.
Jeewoth, T., Bhowon, M. G. & Wah, H. L. K. (1999). *Transition Met. Chem.* **24**, 445–448.
Jing, Z.-L., Yu, M. & Chen, X. (2007). *Acta Cryst.* **E63**, o4902.
Lin, X.-C., Yin, H. & Lin, Y. (2007). *Acta Cryst.* **E63**, o2864.
Lu, J.-F. (2008). *Acta Cryst.* **E64**, o2032.
Ma, H.-B., Huang, S.-S. & Diao, Y.-P. (2008). *Acta Cryst.* **E64**, o210.
Mohd Lair, N., Mohd Ali, H. & Ng, S. W. (2009*a*). *Acta Cryst.* **E65**, o189.
Mohd Lair, N., Mohd Ali, H. & Ng, S. W. (2009*b*). *Acta Cryst.* **E65**, o190.
Prabhakaran, R., Huang, R. & Natarajan, K. (2006). *Inorg. Chim. Acta*, **359**, 3359–3362.
Qiu, X.-Y. (2009). *Z. Kristallogr. New Cryst. Struct.* **224**, 109–110.

- Ren, S., Wang, R., Komatsu, K., Bonaz-Krause, P., Zyrianov, Y., McKenna, C. E., Csipke, C., Tokes, Z. A. & Lien, E. J. (2002). *J. Med. Chem.* **45**, 410–419.
- Salhin, A., Tameem, A. A., Saad, B., Ng, S.-L. & Fun, H.-K. (2007). *Acta Cryst. E* **63**, o2880.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Shivakumar, K., Shashidhar, T., Reddy, P. V. & Halli, M. B. (2008). *J. Coord. Chem.* **61**, 2274–2287.
- Sinha, D., Tiwari, A. K., Singh, S., Shukla, G., Mishra, P., Chandra, H. & Mishra, A. K. (2008). *Eur. J. Med. Chem.* **43**, 160–165.
- Wang, X.-Y., Cao, G.-B. & Yang, T. (2008). *Acta Cryst. E* **64**, o2022.
- Xue, L.-W., Han, Y.-J., Hao, C.-J., Zhao, G.-Q. & Liu, Q.-R. (2008). *Acta Cryst. E* **64**, o1938.

supplementary materials

Acta Cryst. (2009). E65, o1505-o1506 [doi:10.1107/S1600536809020819]

N'-(5-Chloro-2-hydroxybenzylidene)nicotinothiazide

C.-G. Ren

Comment

The Schiff base compounds show excellent biological properties (Jeewoth *et al.*, 1999; Ren *et al.*, 2002; Eltayeb *et al.*, 2008; Sinha *et al.*, 2008). Moreover, the Schiff base compounds have been widely used as versatile ligands in coordination chemistry (Shivakumar *et al.*, 2008; Prabhakaran *et al.*, 2006; Dhar *et al.*, 2005). We report here the crystal structure of the title compound. In the title compound, Fig. 1, there are two independent molecules in the asymmetric unit. The dihedral angles between the benzene and pyridine rings are 12.8 (2) and 1.9 (2)°, respectively. All the bond lengths are within normal values (Allen *et al.*, 1987) and comparable to those in other similar compounds (Cui *et al.*, 2007; Jing *et al.*, 2007; Ma *et al.*, 2008; Salhin *et al.*, 2007; Lin *et al.*, 2007; Alhadi *et al.*, 2008; Xue *et al.*, 2008; Wang *et al.*, 2008; Lu, 2008; Diao *et al.*, 2008; Qiu, 2009; Lair *et al.*, 2009a,b). The molecules of the title compound are linked into centrosymmetric $R^4_4(26)$ motifs by N–H···O and N–H···N interactions (Table 1, Fig. 2) (Bernstein *et al.*, 1995).

Experimental

All the starting materials were obtained with AR grade from Lancaster. 5-Chloro-2-hydroxybenzaldehyde (1.0 mmol, 157.1 mg) and nicotinothiazide (1.0 mmol, 137.1 mg) were refluxed in a 30 ml methanol solution for 30 min to give a clear yellow solution. Yellow block-shaped single crystals of the compound were obtained by slow evaporation of the solution for five days at room temperature.

Refinement

H2 and H5 were located from a difference Fourier map and refined isotropically, with the N–H distance restrained to 0.90 (1) Å, and with U_{iso} restrained to 0.08 Å². Other H atoms were constrained to ideal geometries, with $d(\text{C–H}) = 0.93$ Å, $d(\text{O–H}) = 0.82$ Å, and with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ and $1.5U_{\text{eq}}(\text{O})$.

Figures




Fig. 1. The molecular structure of the compound with 30% probability ellipsoids. The intramolecular O–H···N hydrogen bonds are shown as dashed lines.




Fig. 2. Molecular packing of the compound with hydrogen bonds drawn as dashed lines. Hydrogen atoms not involved in hydrogen bonding have been omitted.

(I)

Crystal data

$C_{13}H_{10}ClN_3O_2$

$M_r = 275.69$

Monoclinic, $P2_1/n$

Hall symbol: $-P\ 2_1/n$

$a = 9.792\ (2)\ \text{\AA}$

$b = 23.358\ (3)\ \text{\AA}$

$c = 10.926\ (2)\ \text{\AA}$

$\beta = 96.848\ (2)^\circ$

$V = 2481.2\ (8)\ \text{\AA}^3$

$Z = 8$

$F_{000} = 1136$

$D_x = 1.476\ \text{Mg m}^{-3}$

Mo $K\alpha$ radiation

$\lambda = 0.71073\ \text{\AA}$

Cell parameters from 2236 reflections

$\theta = 2.6\text{--}24.5^\circ$

$\mu = 0.31\ \text{mm}^{-1}$

$T = 298\ \text{K}$

Block, yellow

$0.30 \times 0.30 \times 0.27\ \text{mm}$

Data collection

Bruker SMART CCD area-detector diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 298\ \text{K}$

ω scans

Absorption correction: multi-scan (SADABS; Sheldrick, 1996)

$T_{\min} = 0.913$, $T_{\max} = 0.921$

14387 measured reflections

5342 independent reflections

3193 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.042$

$\theta_{\max} = 27.0^\circ$

$\theta_{\min} = 1.7^\circ$

$h = -12 \rightarrow 12$

$k = -29 \rightarrow 27$

$l = -12 \rightarrow 13$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.049$

$wR(F^2) = 0.115$

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent and constrained refinement

$$w = 1/[\sigma^2(F_o^2) + (0.0466P)^2]$$

where $P = (F_o^2 + 2F_c^2)/3$

$S = 1.01$ $(\Delta/\sigma)_{\max} < 0.001$
 5342 reflections $\Delta\rho_{\max} = 0.22 \text{ e } \text{\AA}^{-3}$
 351 parameters $\Delta\rho_{\min} = -0.24 \text{ e } \text{\AA}^{-3}$
 2 restraints Extinction correction: none
 Primary atom site location: structure-invariant direct methods

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C11	-0.31666 (7)	0.15824 (3)	0.49570 (7)	0.0648 (2)
C12	0.67723 (7)	-0.10226 (3)	0.57393 (6)	0.0648 (2)
N1	0.28860 (19)	0.22587 (8)	0.70899 (16)	0.0414 (5)
N2	0.4128 (2)	0.20261 (8)	0.75424 (18)	0.0421 (5)
N3	0.7948 (2)	0.13722 (8)	0.92250 (17)	0.0438 (5)
N4	0.29718 (18)	0.00154 (8)	0.94498 (16)	0.0387 (4)
N5	0.2156 (2)	-0.01091 (8)	1.03441 (16)	0.0402 (5)
N6	-0.0434 (2)	-0.04219 (10)	1.3111 (2)	0.0659 (6)
O1	0.13564 (17)	0.31528 (6)	0.65672 (16)	0.0520 (4)
H1	0.2075	0.2981	0.6784	0.078*
O2	0.50725 (19)	0.29034 (7)	0.76057 (19)	0.0755 (6)
O3	0.39503 (17)	0.07872 (6)	0.81707 (15)	0.0486 (4)
H3	0.3484	0.0658	0.8682	0.073*
O4	0.17570 (18)	0.08396 (7)	1.05029 (15)	0.0598 (5)
C1	0.0545 (2)	0.21766 (9)	0.63088 (19)	0.0368 (5)
C2	0.0344 (2)	0.27712 (9)	0.62187 (19)	0.0394 (5)
C3	-0.0937 (3)	0.29868 (10)	0.5765 (2)	0.0496 (6)
H3A	-0.1072	0.3381	0.5715	0.059*
C4	-0.2008 (3)	0.26227 (11)	0.5390 (2)	0.0504 (6)
H4	-0.2865	0.2770	0.5088	0.060*
C5	-0.1810 (2)	0.20380 (10)	0.5461 (2)	0.0438 (6)
C6	-0.0555 (2)	0.18151 (10)	0.59222 (19)	0.0415 (6)
H6	-0.0439	0.1420	0.5977	0.050*
C7	0.1862 (2)	0.19285 (10)	0.6792 (2)	0.0411 (6)
H7	0.1957	0.1534	0.6882	0.049*
C8	0.5195 (2)	0.23961 (10)	0.7816 (2)	0.0444 (6)

supplementary materials

C9	0.6530 (2)	0.21501 (9)	0.83757 (19)	0.0359 (5)
C10	0.7653 (3)	0.25086 (10)	0.8548 (2)	0.0483 (6)
H10	0.7567	0.2890	0.8304	0.058*
C11	0.8900 (3)	0.23037 (11)	0.9080 (2)	0.0585 (7)
H11	0.9658	0.2545	0.9224	0.070*
C12	0.9001 (3)	0.17355 (11)	0.9395 (2)	0.0509 (6)
H12	0.9849	0.1597	0.9746	0.061*
C13	0.6739 (2)	0.15881 (9)	0.8743 (2)	0.0425 (6)
H13	0.5985	0.1343	0.8649	0.051*
C14	0.4402 (2)	-0.02227 (9)	0.79497 (18)	0.0357 (5)
C15	0.4588 (2)	0.03497 (9)	0.7643 (2)	0.0380 (5)
C16	0.5456 (2)	0.04896 (10)	0.6773 (2)	0.0476 (6)
H16	0.5579	0.0872	0.6572	0.057*
C17	0.6134 (2)	0.00709 (10)	0.6205 (2)	0.0464 (6)
H17	0.6724	0.0169	0.5632	0.056*
C18	0.5936 (2)	-0.04957 (10)	0.6489 (2)	0.0446 (6)
C19	0.5080 (2)	-0.06441 (9)	0.7344 (2)	0.0414 (6)
H19	0.4951	-0.1028	0.7523	0.050*
C20	0.3537 (2)	-0.03840 (9)	0.88893 (19)	0.0395 (5)
H20	0.3400	-0.0767	0.9076	0.047*
C21	0.1567 (2)	0.03509 (10)	1.0846 (2)	0.0419 (6)
C22	0.0660 (2)	0.02299 (9)	1.18211 (19)	0.0401 (5)
C23	0.0067 (3)	0.06862 (11)	1.2362 (2)	0.0538 (7)
H23	0.0236	0.1059	1.2124	0.065*
C24	-0.0774 (3)	0.05834 (13)	1.3256 (2)	0.0651 (8)
H24	-0.1193	0.0885	1.3624	0.078*
C25	-0.0986 (3)	0.00294 (14)	1.3595 (2)	0.0657 (8)
H25	-0.1554	-0.0034	1.4206	0.079*
C26	0.0382 (3)	-0.03068 (10)	1.2238 (2)	0.0542 (7)
H26	0.0790	-0.0616	1.1887	0.065*
H5	0.206 (3)	-0.0477 (5)	1.053 (2)	0.080*
H2	0.411 (3)	0.1663 (5)	0.779 (2)	0.080*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C11	0.0444 (4)	0.0649 (5)	0.0827 (5)	-0.0128 (3)	-0.0025 (4)	0.0037 (4)
C12	0.0697 (5)	0.0567 (4)	0.0709 (5)	0.0172 (4)	0.0197 (4)	-0.0037 (3)
N1	0.0369 (12)	0.0387 (11)	0.0479 (11)	0.0041 (9)	0.0020 (9)	0.0047 (9)
N2	0.0351 (12)	0.0342 (11)	0.0552 (12)	0.0012 (9)	-0.0016 (10)	0.0047 (9)
N3	0.0362 (12)	0.0371 (11)	0.0574 (13)	0.0001 (9)	0.0023 (10)	0.0029 (9)
N4	0.0374 (12)	0.0354 (11)	0.0432 (11)	-0.0026 (9)	0.0048 (9)	0.0047 (9)
N5	0.0422 (12)	0.0333 (11)	0.0465 (11)	0.0001 (9)	0.0116 (10)	0.0040 (9)
N6	0.0726 (17)	0.0628 (15)	0.0674 (15)	0.0088 (13)	0.0299 (13)	0.0101 (12)
O1	0.0514 (11)	0.0317 (9)	0.0697 (11)	-0.0002 (8)	-0.0060 (10)	0.0021 (8)
O2	0.0604 (13)	0.0346 (11)	0.1240 (16)	-0.0032 (9)	-0.0194 (12)	0.0196 (10)
O3	0.0549 (11)	0.0292 (9)	0.0649 (11)	-0.0015 (8)	0.0206 (9)	0.0034 (7)
O4	0.0713 (13)	0.0321 (10)	0.0788 (12)	0.0011 (9)	0.0205 (10)	0.0045 (9)

C1	0.0369 (14)	0.0353 (13)	0.0390 (13)	0.0020 (10)	0.0073 (10)	0.0041 (10)
C2	0.0432 (15)	0.0366 (13)	0.0387 (13)	-0.0007 (11)	0.0057 (11)	0.0002 (10)
C3	0.0506 (17)	0.0368 (14)	0.0600 (16)	0.0088 (12)	0.0013 (13)	0.0051 (11)
C4	0.0419 (16)	0.0554 (17)	0.0527 (16)	0.0080 (13)	0.0008 (12)	0.0060 (12)
C5	0.0358 (15)	0.0476 (15)	0.0481 (14)	-0.0051 (11)	0.0056 (11)	0.0018 (11)
C6	0.0430 (15)	0.0354 (13)	0.0467 (14)	-0.0024 (11)	0.0079 (11)	0.0033 (10)
C7	0.0426 (15)	0.0326 (13)	0.0482 (14)	0.0022 (11)	0.0052 (12)	0.0025 (10)
C8	0.0445 (16)	0.0353 (14)	0.0523 (15)	0.0002 (11)	0.0012 (12)	0.0038 (11)
C9	0.0376 (14)	0.0289 (12)	0.0410 (12)	-0.0028 (10)	0.0042 (10)	-0.0003 (9)
C10	0.0498 (17)	0.0308 (13)	0.0635 (16)	-0.0067 (11)	0.0035 (13)	0.0037 (11)
C11	0.0433 (16)	0.0477 (16)	0.0815 (19)	-0.0131 (13)	-0.0043 (14)	0.0003 (14)
C12	0.0367 (15)	0.0520 (17)	0.0625 (16)	0.0016 (12)	0.0000 (12)	0.0000 (12)
C13	0.0338 (14)	0.0358 (13)	0.0571 (15)	-0.0048 (11)	0.0027 (12)	0.0019 (11)
C14	0.0332 (13)	0.0330 (12)	0.0399 (13)	-0.0017 (10)	-0.0006 (10)	0.0050 (10)
C15	0.0335 (13)	0.0355 (13)	0.0444 (13)	-0.0027 (10)	0.0020 (11)	0.0005 (10)
C16	0.0530 (16)	0.0365 (14)	0.0542 (15)	-0.0072 (12)	0.0097 (13)	0.0051 (11)
C17	0.0454 (16)	0.0496 (15)	0.0457 (14)	-0.0044 (12)	0.0121 (12)	0.0040 (12)
C18	0.0449 (15)	0.0427 (14)	0.0455 (14)	0.0065 (11)	0.0029 (12)	0.0007 (11)
C19	0.0445 (15)	0.0330 (13)	0.0459 (13)	0.0020 (11)	0.0025 (11)	0.0062 (10)
C20	0.0417 (14)	0.0281 (12)	0.0482 (14)	-0.0016 (10)	0.0037 (11)	0.0049 (10)
C21	0.0393 (14)	0.0365 (14)	0.0484 (14)	-0.0008 (11)	-0.0004 (11)	-0.0002 (11)
C22	0.0367 (14)	0.0385 (14)	0.0439 (13)	0.0042 (10)	0.0005 (11)	-0.0027 (10)
C23	0.0542 (17)	0.0453 (15)	0.0622 (17)	0.0041 (13)	0.0073 (14)	-0.0108 (12)
C24	0.0600 (19)	0.068 (2)	0.0687 (19)	0.0135 (15)	0.0139 (15)	-0.0213 (15)
C25	0.0514 (18)	0.094 (2)	0.0538 (17)	0.0100 (17)	0.0166 (14)	-0.0020 (16)
C26	0.0606 (18)	0.0439 (16)	0.0619 (16)	0.0083 (13)	0.0230 (14)	-0.0006 (12)

Geometric parameters (Å, °)

C11—C5	1.740 (2)	C7—H7	0.9300
C12—C18	1.736 (2)	C8—C9	1.491 (3)
N1—C7	1.276 (3)	C9—C10	1.376 (3)
N1—N2	1.369 (2)	C9—C13	1.381 (3)
N2—C8	1.361 (3)	C10—C11	1.375 (3)
N2—H2	0.892 (10)	C10—H10	0.9300
N3—C12	1.332 (3)	C11—C12	1.372 (3)
N3—C13	1.336 (3)	C11—H11	0.9300
N4—C20	1.278 (3)	C12—H12	0.9300
N4—N5	1.365 (2)	C13—H13	0.9300
N5—C21	1.365 (3)	C14—C15	1.396 (3)
N5—H5	0.891 (10)	C14—C19	1.397 (3)
N6—C25	1.324 (3)	C14—C20	1.456 (3)
N6—C26	1.342 (3)	C15—C16	1.388 (3)
O1—C2	1.354 (3)	C16—C17	1.371 (3)
O1—H1	0.8200	C16—H16	0.9300
O2—C8	1.210 (2)	C17—C18	1.378 (3)
O3—C15	1.361 (2)	C17—H17	0.9300
O3—H3	0.8200	C18—C19	1.373 (3)
O4—C21	1.223 (2)	C19—H19	0.9300

supplementary materials

C1—C6	1.394 (3)	C20—H20	0.9300
C1—C2	1.405 (3)	C21—C22	1.493 (3)
C1—C7	1.455 (3)	C22—C26	1.372 (3)
C2—C3	1.387 (3)	C22—C23	1.380 (3)
C3—C4	1.375 (3)	C23—C24	1.372 (3)
C3—H3A	0.9300	C23—H23	0.9300
C4—C5	1.380 (3)	C24—C25	1.369 (4)
C4—H4	0.9300	C24—H24	0.9300
C5—C6	1.374 (3)	C25—H25	0.9300
C6—H6	0.9300	C26—H26	0.9300
C7—N1—N2	119.29 (19)	N3—C12—C11	123.3 (2)
C8—N2—N1	116.92 (19)	N3—C12—H12	118.3
C8—N2—H2	125.2 (17)	C11—C12—H12	118.3
N1—N2—H2	116.2 (18)	N3—C13—C9	124.5 (2)
C12—N3—C13	116.7 (2)	N3—C13—H13	117.7
C20—N4—N5	120.75 (18)	C9—C13—H13	117.7
C21—N5—N4	115.52 (18)	C15—C14—C19	118.6 (2)
C21—N5—H5	127.1 (17)	C15—C14—C20	121.4 (2)
N4—N5—H5	117.4 (17)	C19—C14—C20	120.05 (19)
C25—N6—C26	115.5 (2)	O3—C15—C16	117.57 (19)
C2—O1—H1	109.5	O3—C15—C14	122.55 (19)
C15—O3—H3	109.5	C16—C15—C14	119.9 (2)
C6—C1—C2	118.8 (2)	C17—C16—C15	120.7 (2)
C6—C1—C7	119.2 (2)	C17—C16—H16	119.6
C2—C1—C7	122.0 (2)	C15—C16—H16	119.6
O1—C2—C3	117.5 (2)	C16—C17—C18	119.7 (2)
O1—C2—C1	122.7 (2)	C16—C17—H17	120.2
C3—C2—C1	119.8 (2)	C18—C17—H17	120.2
C4—C3—C2	120.5 (2)	C19—C18—C17	120.6 (2)
C4—C3—H3A	119.8	C19—C18—C12	120.18 (18)
C2—C3—H3A	119.8	C17—C18—C12	119.23 (18)
C3—C4—C5	119.9 (2)	C18—C19—C14	120.5 (2)
C3—C4—H4	120.1	C18—C19—H19	119.7
C5—C4—H4	120.1	C14—C19—H19	119.7
C6—C5—C4	120.6 (2)	N4—C20—C14	118.06 (19)
C6—C5—C11	120.01 (19)	N4—C20—H20	121.0
C4—C5—C11	119.37 (19)	C14—C20—H20	121.0
C5—C6—C1	120.4 (2)	O4—C21—N5	121.7 (2)
C5—C6—H6	119.8	O4—C21—C22	121.3 (2)
C1—C6—H6	119.8	N5—C21—C22	116.94 (19)
N1—C7—C1	119.2 (2)	C26—C22—C23	116.9 (2)
N1—C7—H7	120.4	C26—C22—C21	124.6 (2)
C1—C7—H7	120.4	C23—C22—C21	118.4 (2)
O2—C8—N2	121.6 (2)	C24—C23—C22	119.3 (2)
O2—C8—C9	121.2 (2)	C24—C23—H23	120.4
N2—C8—C9	117.3 (2)	C22—C23—H23	120.4
C10—C9—C13	116.8 (2)	C25—C24—C23	118.8 (2)
C10—C9—C8	117.9 (2)	C25—C24—H24	120.6
C13—C9—C8	125.3 (2)	C23—C24—H24	120.6

C11—C10—C9	120.0 (2)	N6—C25—C24	124.2 (3)
C11—C10—H10	120.0	N6—C25—H25	117.9
C9—C10—H10	120.0	C24—C25—H25	117.9
C12—C11—C10	118.5 (2)	N6—C26—C22	125.3 (2)
C12—C11—H11	120.8	N6—C26—H26	117.4
C10—C11—H11	120.8	C22—C26—H26	117.4
C7—N1—N2—C8	178.3 (2)	C8—C9—C13—N3	178.8 (2)
C20—N4—N5—C21	-177.7 (2)	C19—C14—C15—O3	178.9 (2)
C6—C1—C2—O1	-179.7 (2)	C20—C14—C15—O3	-2.0 (3)
C7—C1—C2—O1	0.3 (3)	C19—C14—C15—C16	-1.3 (3)
C6—C1—C2—C3	0.8 (3)	C20—C14—C15—C16	177.9 (2)
C7—C1—C2—C3	-179.2 (2)	O3—C15—C16—C17	179.9 (2)
O1—C2—C3—C4	179.7 (2)	C14—C15—C16—C17	0.1 (3)
C1—C2—C3—C4	-0.7 (3)	C15—C16—C17—C18	1.0 (4)
C2—C3—C4—C5	-0.2 (4)	C16—C17—C18—C19	-0.8 (4)
C3—C4—C5—C6	1.0 (3)	C16—C17—C18—C12	178.62 (18)
C3—C4—C5—C11	-178.88 (18)	C17—C18—C19—C14	-0.5 (3)
C4—C5—C6—C1	-1.0 (3)	C12—C18—C19—C14	-179.87 (17)
C11—C5—C6—C1	178.92 (16)	C15—C14—C19—C18	1.5 (3)
C2—C1—C6—C5	0.1 (3)	C20—C14—C19—C18	-177.7 (2)
C7—C1—C6—C5	-179.9 (2)	N5—N4—C20—C14	-179.66 (18)
N2—N1—C7—C1	-179.88 (18)	C15—C14—C20—N4	-1.5 (3)
C6—C1—C7—N1	176.7 (2)	C19—C14—C20—N4	177.7 (2)
C2—C1—C7—N1	-3.3 (3)	N4—N5—C21—O4	0.5 (3)
N1—N2—C8—O2	-3.9 (3)	N4—N5—C21—C22	179.61 (18)
N1—N2—C8—C9	176.60 (18)	O4—C21—C22—C26	178.9 (2)
O2—C8—C9—C10	-5.6 (3)	N5—C21—C22—C26	-0.2 (3)
N2—C8—C9—C10	173.9 (2)	O4—C21—C22—C23	-2.0 (3)
O2—C8—C9—C13	174.2 (2)	N5—C21—C22—C23	178.9 (2)
N2—C8—C9—C13	-6.3 (3)	C26—C22—C23—C24	-1.1 (4)
C13—C9—C10—C11	-0.9 (3)	C21—C22—C23—C24	179.7 (2)
C8—C9—C10—C11	178.9 (2)	C22—C23—C24—C25	0.8 (4)
C9—C10—C11—C12	2.0 (4)	C26—N6—C25—C24	0.2 (4)
C13—N3—C12—C11	-1.3 (4)	C23—C24—C25—N6	-0.4 (4)
C10—C11—C12—N3	-0.8 (4)	C25—N6—C26—C22	-0.6 (4)
C12—N3—C13—C9	2.5 (3)	C23—C22—C26—N6	1.1 (4)
C10—C9—C13—N3	-1.3 (3)	C21—C22—C26—N6	-179.8 (2)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O1—H1 \cdots N1	0.82	1.88	2.594 (2)	145
O3—H3 \cdots N4	0.82	1.82	2.538 (2)	146
N5—H5 \cdots N3 ⁱ	0.891 (10)	2.109 (11)	2.991 (3)	171 (2)
N2—H2 \cdots O3	0.892 (10)	2.097 (10)	2.984 (2)	173 (2)

Symmetry codes: (i) $-x+1, -y, -z+2$.

Fig. 1




Fig. 2

