

N,N'-(Biphenyl-2,2'-diyl)bis(furan-2-carboxamide)

Chin Hui Kee, Noel F. Thomas, Azhar Ariffin, Khalijah Awang and Seik Weng Ng*

Department of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia
Correspondence e-mail: seikweng@um.edu.my


Received 6 June 2009; accepted 9 June 2009

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.037; wR factor = 0.111; data-to-parameter ratio = 14.7.

The title molecule, $\text{C}_{22}\text{H}_{16}\text{N}_2\text{O}_4$, is a 2,2'-disubstituted biphenyl whose phenylene rings are rotated by $66.5(1)^\circ$ so as to avoid repulsion by the substituents. Only one of the two amide $-\text{NH}-$ fragments engages in hydrogen bonding, and this interacts with the amido $-\text{C}(=\text{O})-$ acceptor of an inversion-related molecule to generate a hydrogen-bonded dimer.

Related literature

The Heck reaction produces the desired stilbene along with a symmetrical biaryl owing to homocoupling of the aryl halide reactant. For the synthesis of stilbene carboxamides synthesized by using radical cations in a modified Heck reaction; see: Thomas *et al.* (2008).



Experimental

Crystal data

$\text{C}_{22}\text{H}_{16}\text{N}_2\text{O}_4$
 $M_r = 372.37$
Triclinic, $P\bar{1}$
 $a = 8.1784(2)$ Å
 $b = 10.1399(2)$ Å
 $c = 11.1475(2)$ Å
 $\alpha = 99.938(1)^\circ$
 $\beta = 107.521(1)^\circ$
 $\gamma = 92.439(1)^\circ$
 $V = 863.92(3)$ Å³
 $Z = 2$
Mo $K\alpha$ radiation
 $\mu = 0.10$ mm⁻¹
 $T = 100$ K
 $0.30 \times 0.25 \times 0.20$ mm

Data collection

Bruker SMART APEX diffractometer
Absorption correction: none
6080 measured reflections
3836 independent reflections
3349 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.015$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.037$
 $wR(F^2) = 0.111$
 $S = 1.03$
3836 reflections
261 parameters
2 restraints
H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.33$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.27$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N2}-\text{H2}\cdots\text{O2}^i$	0.88 (1)	2.12 (1)	2.970 (1)	165 (1)

Symmetry code: (i) $-x + 1, -y + 1, -z + 1$.

Data collection: *APEX2* (Bruker, 2008); cell refinement: *SAINT* (Bruker, 2008); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2009).

We thank the Malaysia Toray Science Foundation for supporting this study. CHK thanks MOSTI for an NSF scholarship.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: TK2474).

References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
Bruker (2008). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Thomas, N. F., Kee, C.-H., Ariffin, A., Awang, K., Weber, J.-F. F., Lim, C.-G., Mukhtar, M. R. & Abdul Hadi, A. H. (2008). *Heterocycles*, **75**, 1097–1108.
Westrip, S. P. (2009). *publCIF*. In preparation.

supplementary materials

Acta Cryst. (2009). E65, o1556 [doi:10.1107/S1600536809021874]

N,N'-(Biphenyl-2,2'-diyl)bis(furan-2-carboxamide)

C. H. Kee, N. F. Thomas, A. Ariffin, K. Awang and S. W. Ng

Experimental

N-(2-Iodophenyl)furan-2-carboxamide (1.51 g, 4.81 mmol) was dissolved in DMF (30 ml) under a nitrogen atmosphere. The solution was heated to 393 K. Palladium acetate (0.01 g, 0.05 mmol) was added followed by triethylamine (2.4 ml, 0.2 mol) and 4-methoxystyrene (0.83 g, 6.16 mmol). The mixture was heated for 48 h. The solution was cooled and then mixed with saturated sodium chloride. The organic compound was extracted with ethylacetate. The ethylacetate solution was dried with sodium sulfate. The solvent was evaporated and the product purified by column chromatography (0.12 g, 10% yield). Single crystals were obtained by recrystallization from hexane/chloroform.

Refinement

Carbon-bound H atoms were placed in calculated positions (C—H 0.95 Å) and were included in the refinement in the riding model approximation with $U(\text{H})$ set to $1.2U_{\text{eq}}(\text{C})$. The nitrogen-bound H atoms were located in a difference Fourier map, and were refined with a restraint of N—H 0.88±0.01 Å

Figures




Fig. 1. Thermal ellipsoid plot (Barbour, 2001) of $\text{C}_{22}\text{H}_{16}\text{N}_2\text{O}_4$ at the 70% probability level. Hydrogen atoms are drawn as spheres of arbitrary radii.




Fig. 2. Hydrogen-bonded dimer in the crystal structure of (I). Red dashed lines represent N—H...O hydrogen bonds.

N,N'-(Biphenyl-2,2'-diyl)bis(furan-2-carboxamide)

Crystal data

$\text{C}_{22}\text{H}_{16}\text{N}_2\text{O}_4$

$M_r = 372.37$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$Z = 2$

$F_{000} = 388$

$D_x = 1.431 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation

supplementary materials

$a = 8.1784$ (2) Å	$\lambda = 0.71073$ Å
$b = 10.1399$ (2) Å	Cell parameters from 3508 reflections
$c = 11.1475$ (2) Å	$\theta = 2.6\text{--}28.3^\circ$
$\alpha = 99.938$ (1) $^\circ$	$\mu = 0.10$ mm $^{-1}$
$\beta = 107.521$ (1) $^\circ$	$T = 100$ K
$\gamma = 92.439$ (1) $^\circ$	Block, colorless
$V = 863.92$ (3) Å 3	$0.30 \times 0.25 \times 0.20$ mm

Data collection

Bruker SMART APEX diffractometer	3349 reflections with $I > 2\sigma(I)$
Radiation source: fine-focus sealed tube	$R_{\text{int}} = 0.015$
Monochromator: graphite	$\theta_{\text{max}} = 27.5^\circ$
$T = 100$ K	$\theta_{\text{min}} = 2.0^\circ$
ω scans	$h = -10 \rightarrow 10$
Absorption correction: None	$k = -12 \rightarrow 13$
6080 measured reflections	$l = -14 \rightarrow 14$
3836 independent reflections	

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.037$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.111$	$w = 1/[\sigma^2(F_o^2) + (0.0627P)^2 + 0.3539P]$
$S = 1.03$	where $P = (F_o^2 + 2F_c^2)/3$
3836 reflections	$(\Delta/\sigma)_{\text{max}} = 0.001$
261 parameters	$\Delta\rho_{\text{max}} = 0.33$ e Å $^{-3}$
2 restraints	$\Delta\rho_{\text{min}} = -0.27$ e Å $^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å 2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.37952 (12)	0.28756 (9)	0.22024 (9)	0.0174 (2)
O2	0.32912 (12)	0.61791 (9)	0.36569 (9)	0.0186 (2)
O3	0.42615 (12)	0.02447 (10)	0.82668 (9)	0.0213 (2)
O4	0.80281 (12)	0.23741 (9)	0.86550 (9)	0.0191 (2)
N1	0.29107 (14)	0.41257 (11)	0.42005 (10)	0.0156 (2)
H1	0.295 (2)	0.3271 (10)	0.3921 (17)	0.028 (5)*
N2	0.47787 (14)	0.16551 (11)	0.70037 (10)	0.0148 (2)
H2	0.5491 (17)	0.2298 (13)	0.6951 (15)	0.016 (4)*

C1	0.40891 (17)	0.24580 (13)	0.10565 (13)	0.0190 (3)
H1a	0.4304	0.1570	0.0749	0.023*
C2	0.40293 (17)	0.34894 (14)	0.04255 (13)	0.0190 (3)
H2a	0.4180	0.3456	-0.0391	0.023*
C3	0.36958 (16)	0.46370 (13)	0.12235 (12)	0.0173 (3)
H3	0.3581	0.5517	0.1044	0.021*
C4	0.35755 (16)	0.42216 (12)	0.22885 (12)	0.0155 (3)
C5	0.32496 (16)	0.49452 (12)	0.34438 (12)	0.0145 (2)
C6	0.24862 (16)	0.44495 (13)	0.53444 (12)	0.0144 (2)
C7	0.23946 (17)	0.57726 (13)	0.59222 (12)	0.0169 (3)
H7	0.2603	0.6501	0.5537	0.020*
C8	0.19978 (17)	0.60133 (13)	0.70591 (12)	0.0182 (3)
H8	0.1950	0.6913	0.7455	0.022*
C9	0.16695 (17)	0.49624 (14)	0.76291 (12)	0.0192 (3)
H9	0.1386	0.5138	0.8404	0.023*
C10	0.17609 (17)	0.36502 (13)	0.70536 (12)	0.0174 (3)
H10	0.1539	0.2928	0.7443	0.021*
C11	0.21726 (16)	0.33732 (12)	0.59167 (12)	0.0142 (2)
C12	0.21741 (16)	0.19410 (12)	0.53056 (12)	0.0142 (3)
C13	0.08880 (17)	0.13778 (13)	0.41514 (12)	0.0164 (3)
H13	0.0094	0.1937	0.3731	0.020*
C14	0.07515 (17)	0.00244 (13)	0.36117 (12)	0.0182 (3)
H14	-0.0121	-0.0337	0.2825	0.022*
C15	0.18976 (17)	-0.08003 (13)	0.42285 (12)	0.0178 (3)
H15	0.1790	-0.1734	0.3877	0.021*
C16	0.31983 (16)	-0.02613 (13)	0.53571 (12)	0.0161 (3)
H16	0.3991	-0.0826	0.5770	0.019*
C17	0.33501 (16)	0.11065 (12)	0.58905 (12)	0.0141 (2)
C18	0.52125 (16)	0.10827 (12)	0.80562 (12)	0.0150 (3)
C19	0.69553 (16)	0.15138 (12)	0.89706 (12)	0.0155 (3)
C20	0.77797 (18)	0.11564 (13)	1.01001 (13)	0.0195 (3)
H20	0.7319	0.0577	1.0529	0.023*
C21	0.94791 (18)	0.18268 (15)	1.05134 (13)	0.0226 (3)
H21	1.0380	0.1779	1.1272	0.027*
C22	0.95558 (18)	0.25413 (15)	0.96175 (13)	0.0226 (3)
H22	1.0546	0.3090	0.9652	0.027*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0214 (5)	0.0138 (4)	0.0181 (4)	0.0017 (4)	0.0076 (4)	0.0032 (3)
O2	0.0226 (5)	0.0136 (4)	0.0217 (5)	0.0018 (4)	0.0085 (4)	0.0058 (4)
O3	0.0197 (5)	0.0241 (5)	0.0206 (5)	-0.0031 (4)	0.0051 (4)	0.0095 (4)
O4	0.0162 (5)	0.0212 (5)	0.0196 (5)	-0.0015 (4)	0.0041 (4)	0.0065 (4)
N1	0.0211 (6)	0.0113 (5)	0.0162 (5)	0.0023 (4)	0.0082 (4)	0.0031 (4)
N2	0.0136 (5)	0.0140 (5)	0.0169 (5)	-0.0005 (4)	0.0037 (4)	0.0057 (4)
C1	0.0176 (6)	0.0193 (6)	0.0187 (6)	-0.0010 (5)	0.0066 (5)	-0.0011 (5)
C2	0.0166 (6)	0.0231 (6)	0.0170 (6)	-0.0003 (5)	0.0058 (5)	0.0029 (5)

supplementary materials

C3	0.0160 (6)	0.0178 (6)	0.0185 (6)	0.0005 (5)	0.0056 (5)	0.0050 (5)
C4	0.0143 (6)	0.0132 (6)	0.0185 (6)	0.0005 (5)	0.0036 (5)	0.0044 (5)
C5	0.0124 (5)	0.0151 (6)	0.0158 (6)	0.0013 (4)	0.0031 (5)	0.0050 (5)
C6	0.0132 (6)	0.0158 (6)	0.0140 (6)	0.0022 (4)	0.0031 (5)	0.0039 (5)
C7	0.0176 (6)	0.0147 (6)	0.0177 (6)	0.0030 (5)	0.0038 (5)	0.0044 (5)
C8	0.0182 (6)	0.0163 (6)	0.0176 (6)	0.0056 (5)	0.0026 (5)	0.0012 (5)
C9	0.0198 (6)	0.0230 (7)	0.0143 (6)	0.0049 (5)	0.0048 (5)	0.0031 (5)
C10	0.0178 (6)	0.0192 (6)	0.0157 (6)	0.0015 (5)	0.0044 (5)	0.0060 (5)
C11	0.0120 (6)	0.0143 (6)	0.0148 (6)	0.0019 (4)	0.0016 (5)	0.0037 (4)
C12	0.0153 (6)	0.0140 (6)	0.0150 (6)	-0.0001 (5)	0.0063 (5)	0.0047 (5)
C13	0.0178 (6)	0.0162 (6)	0.0154 (6)	0.0026 (5)	0.0044 (5)	0.0053 (5)
C14	0.0194 (6)	0.0181 (6)	0.0153 (6)	-0.0015 (5)	0.0040 (5)	0.0018 (5)
C15	0.0211 (6)	0.0139 (6)	0.0193 (6)	-0.0002 (5)	0.0089 (5)	0.0018 (5)
C16	0.0157 (6)	0.0146 (6)	0.0202 (6)	0.0034 (5)	0.0078 (5)	0.0053 (5)
C17	0.0141 (6)	0.0156 (6)	0.0137 (6)	0.0000 (5)	0.0057 (5)	0.0040 (4)
C18	0.0154 (6)	0.0147 (6)	0.0156 (6)	0.0032 (5)	0.0055 (5)	0.0032 (5)
C19	0.0163 (6)	0.0147 (6)	0.0166 (6)	0.0018 (5)	0.0064 (5)	0.0036 (5)
C20	0.0205 (7)	0.0206 (6)	0.0171 (6)	0.0020 (5)	0.0049 (5)	0.0049 (5)
C21	0.0183 (7)	0.0283 (7)	0.0178 (6)	0.0018 (5)	0.0017 (5)	0.0023 (5)
C22	0.0152 (6)	0.0277 (7)	0.0219 (7)	-0.0026 (5)	0.0032 (5)	0.0024 (5)

Geometric parameters (Å, °)

O1—C1	1.3695 (16)	C8—H8	0.9500
O1—C4	1.3749 (15)	C9—C10	1.3893 (18)
O2—C5	1.2295 (15)	C9—H9	0.9500
O3—C18	1.2267 (16)	C10—C11	1.3931 (18)
O4—C22	1.3613 (16)	C10—H10	0.9500
O4—C19	1.3716 (15)	C11—C12	1.4941 (17)
N1—C5	1.3572 (16)	C12—C17	1.3957 (18)
N1—C6	1.4093 (16)	C12—C13	1.4030 (17)
N1—H1	0.875 (9)	C13—C14	1.3853 (17)
N2—C18	1.3569 (16)	C13—H13	0.9500
N2—C17	1.4311 (16)	C14—C15	1.3884 (19)
N2—H2	0.877 (9)	C14—H14	0.9500
C1—C2	1.3521 (19)	C15—C16	1.3858 (18)
C1—H1a	0.9500	C15—H15	0.9500
C2—C3	1.4267 (19)	C16—C17	1.3961 (17)
C2—H2a	0.9500	C16—H16	0.9500
C3—C4	1.3550 (18)	C18—C19	1.4765 (18)
C3—H3	0.9500	C19—C20	1.3559 (18)
C4—C5	1.4715 (18)	C20—C21	1.4255 (19)
C6—C7	1.3999 (17)	C20—H20	0.9500
C6—C11	1.4067 (17)	C21—C22	1.345 (2)
C7—C8	1.3851 (18)	C21—H21	0.9500
C7—H7	0.9500	C22—H22	0.9500
C8—C9	1.3869 (19)		
C1—O1—C4	106.32 (10)	C11—C10—H10	119.4
C22—O4—C19	105.90 (10)	C10—C11—C6	118.87 (12)

C5—N1—C6	129.64 (11)	C10—C11—C12	119.07 (11)
C5—N1—H1	114.1 (12)	C6—C11—C12	121.97 (11)
C6—N1—H1	116.3 (12)	C17—C12—C13	118.24 (11)
C18—N2—C17	122.19 (10)	C17—C12—C11	122.04 (11)
C18—N2—H2	118.8 (10)	C13—C12—C11	119.58 (11)
C17—N2—H2	118.2 (10)	C14—C13—C12	121.42 (12)
C2—C1—O1	110.34 (11)	C14—C13—H13	119.3
C2—C1—H1a	124.8	C12—C13—H13	119.3
O1—C1—H1a	124.8	C13—C14—C15	119.56 (12)
C1—C2—C3	106.73 (12)	C13—C14—H14	120.2
C1—C2—H2a	126.6	C15—C14—H14	120.2
C3—C2—H2a	126.6	C16—C15—C14	120.02 (12)
C4—C3—C2	106.31 (12)	C16—C15—H15	120.0
C4—C3—H3	126.8	C14—C15—H15	120.0
C2—C3—H3	126.8	C15—C16—C17	120.36 (12)
C3—C4—O1	110.30 (11)	C15—C16—H16	119.8
C3—C4—C5	131.50 (12)	C17—C16—H16	119.8
O1—C4—C5	118.19 (11)	C12—C17—C16	120.35 (11)
O2—C5—N1	125.53 (12)	C12—C17—N2	120.43 (11)
O2—C5—C4	120.77 (11)	C16—C17—N2	119.14 (11)
N1—C5—C4	113.70 (11)	O3—C18—N2	124.03 (12)
C7—C6—C11	120.02 (11)	O3—C18—C19	119.98 (12)
C7—C6—N1	122.95 (11)	N2—C18—C19	115.98 (11)
C11—C6—N1	117.03 (11)	C20—C19—O4	110.52 (11)
C8—C7—C6	119.61 (12)	C20—C19—C18	131.14 (12)
C8—C7—H7	120.2	O4—C19—C18	118.26 (11)
C6—C7—H7	120.2	C19—C20—C21	106.02 (12)
C9—C8—C7	121.07 (12)	C19—C20—H20	127.0
C9—C8—H8	119.5	C21—C20—H20	127.0
C7—C8—H8	119.5	C22—C21—C20	106.47 (12)
C8—C9—C10	119.20 (12)	C22—C21—H21	126.8
C8—C9—H9	120.4	C20—C21—H21	126.8
C10—C9—H9	120.4	C21—C22—O4	111.09 (12)
C9—C10—C11	121.23 (12)	C21—C22—H22	124.5
C9—C10—H10	119.4	O4—C22—H22	124.5
C4—O1—C1—C2	-1.04 (14)	C10—C11—C12—C13	109.90 (14)
O1—C1—C2—C3	0.66 (15)	C6—C11—C12—C13	-66.72 (16)
C1—C2—C3—C4	0.00 (14)	C17—C12—C13—C14	1.43 (19)
C2—C3—C4—O1	-0.65 (14)	C11—C12—C13—C14	-174.25 (11)
C2—C3—C4—C5	179.98 (13)	C12—C13—C14—C15	0.66 (19)
C1—O1—C4—C3	1.04 (14)	C13—C14—C15—C16	-1.8 (2)
C1—O1—C4—C5	-179.49 (11)	C14—C15—C16—C17	0.93 (19)
C6—N1—C5—O2	2.2 (2)	C13—C12—C17—C16	-2.34 (18)
C6—N1—C5—C4	-177.63 (12)	C11—C12—C17—C16	173.22 (11)
C3—C4—C5—O2	-11.2 (2)	C13—C12—C17—N2	174.26 (11)
O1—C4—C5—O2	169.42 (11)	C11—C12—C17—N2	-10.18 (18)
C3—C4—C5—N1	168.56 (13)	C15—C16—C17—C12	1.20 (18)
O1—C4—C5—N1	-10.77 (16)	C15—C16—C17—N2	-175.44 (11)
C5—N1—C6—C7	-1.9 (2)	C18—N2—C17—C12	133.43 (13)

supplementary materials

C5—N1—C6—C11	178.99 (12)	C18—N2—C17—C16	-49.93 (17)
C11—C6—C7—C8	0.14 (19)	C17—N2—C18—O3	-15.7 (2)
N1—C6—C7—C8	-178.89 (12)	C17—N2—C18—C19	163.63 (11)
C6—C7—C8—C9	-0.7 (2)	C22—O4—C19—C20	0.32 (15)
C7—C8—C9—C10	0.7 (2)	C22—O4—C19—C18	-176.70 (11)
C8—C9—C10—C11	-0.1 (2)	O3—C18—C19—C20	-1.2 (2)
C9—C10—C11—C6	-0.48 (19)	N2—C18—C19—C20	179.49 (13)
C9—C10—C11—C12	-177.20 (12)	O3—C18—C19—O4	175.11 (11)
C7—C6—C11—C10	0.46 (18)	N2—C18—C19—O4	-4.22 (17)
N1—C6—C11—C10	179.55 (11)	O4—C19—C20—C21	-0.45 (15)
C7—C6—C11—C12	177.08 (11)	C18—C19—C20—C21	176.06 (13)
N1—C6—C11—C12	-3.83 (18)	C19—C20—C21—C22	0.40 (16)
C10—C11—C12—C17	-65.60 (16)	C20—C21—C22—O4	-0.22 (16)
C6—C11—C12—C17	117.78 (14)	C19—O4—C22—C21	-0.05 (15)

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$N2-H2\cdots O2^i$	0.88 (1)	2.12 (1)	2.970 (1)	165 (1)

Symmetry codes: (i) $-x+1, -y+1, -z+1$.

Fig. 1




Fig. 2

