

3,3,12,12-Tetramethyl-1,5,10,14-tetra-oxadispiro[5.2.5.2]hexadecane

Yong-Jun He, Min-Hao Xie,* Pei Zou, Ya-Ling Liu and Jun Wu

Jiangsu Institute of Nuclear Medicine, Wuxi 214063, People's Republic of China
Correspondence e-mail: yongjunhe001@hotmail.com


Received 18 May 2009; accepted 4 June 2009

Key indicators: single-crystal X-ray study; $T = 93\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.049; wR factor = 0.095; data-to-parameter ratio = 18.8.

The molecule of the title compound, $\text{C}_{16}\text{H}_{28}\text{O}_4$, is centrosymmetric. The cyclohexane ring and both six-membered dioxane rings adopt chair conformations. In the crystal, the molecules lie in layers in the (100) planes and the shortest intermolecular contacts are $\text{H}\cdots\text{H}$ (2.30 \AA).

Related literature

The title compound is an intermediate in the synthesis of Frovatriptan, a 5-HT₁-like agonist, see: Borrett *et al.* (1999). For details of the synthesis, see: Babler & Spina (1984); Borrett *et al.* (1999). For a related structure, see: Luger *et al.* (1972).



Experimental

Crystal data

$\text{C}_{16}\text{H}_{28}\text{O}_4$	$V = 772.1 (9)\text{ \AA}^3$
$M_r = 284.38$	$Z = 2$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
$a = 12.639 (8)\text{ \AA}$	$\mu = 0.09\text{ mm}^{-1}$
$b = 5.838 (4)\text{ \AA}$	$T = 93\text{ K}$
$c = 11.179 (7)\text{ \AA}$	$0.43 \times 0.37 \times 0.14\text{ mm}$
$\beta = 110.611 (8)^\circ$	

Data collection

Rigaku SPIDER diffractometer	1752 independent reflections
Absorption correction: none	1243 reflections with $I > 2\sigma(I)$
5721 measured reflections	$R_{\text{int}} = 0.048$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.049$	93 parameters
$wR(F^2) = 0.095$	H-atom parameters constrained
$S = 1.00$	$\Delta\rho_{\text{max}} = 0.27\text{ e \AA}^{-3}$
1752 reflections	$\Delta\rho_{\text{min}} = -0.19\text{ e \AA}^{-3}$

Data collection: *RAPID-AUTO* (Rigaku 2004); cell refinement: *RAPID-AUTO*; data reduction: *RAPID-AUTO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The authors acknowledge financial support from Jiangsu Institute of Nuclear Medicine.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BI2376).

References

- Babler, J. H. & Spina, K. P. (1984). *Synth. Commun.* **14**, 39–44.
- Borrett, G. T., Kitteringham, J., Porter, R. A., Shipton, M. R., Vimal, M. & Young, R. C. (1999). US Patent No. 5 962 501.
- Luger, P., Plieth, K. & Ruban, G. (1972). *Acta Cryst. B* **28**, 706–710.
- Rigaku (2004). *RAPID-AUTO*. Rigaku Corporation, Tokyo, Japan.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.

supplementary materials

Acta Cryst. (2009). E65, o1559 [doi:10.1107/S1600536809021291]

3,3,12,12-Tetramethyl-1,5,10,14-tetraoxadispiro[5.2.5.2]hexadecane

Y.-J. He, M.-H. Xie, P. Zou, Y.-L. Liu and J. Wu

Comment

The title compound is useful as an intermediate in the synthesis of Frovatriptan, a 5-HT₁-like agonist (Babler & Spina, 1984; Borrett *et al.*, 1999). The molecular structure is shown in Fig. 1. Three six-membered rings exhibit chair conformations, with C—C bond lengths in the range 1.516 (2)–1.527 (2) Å and C—C—C angles in the range 106.59 (13)–112.55 (13)°; these agree well with the values in other cyclohexane derivatives described in the literature (Luger *et al.*, 1972).

Experimental

The title compound was obtained by reaction of 1,4-cyclohexanedione (20 mmol), 2,2-dimethyl-1,3-propanediol (40 mmol) and sulfuric acid (0.1 mmol) in hexane (20 ml). The mixture was stirred for 6 h at 333 K. Colourless block-shaped crystals suitable for X-ray diffraction analysis were grown at the bottom of the vessel after 7 days slow evaporation of the solution at room temperature.

Refinement

H atoms were placed in calculated positions with C—H = 0.99 and 0.98 Å for methylene and methyl H atoms, respectively, and refined as riding with $U_{\text{iso}}(\text{H}) = 1.2$ or $1.5U_{\text{eq}}(\text{C})$.

Figures



Fig. 1. The molecular structure with displacement ellipsoids drawn at the 30% probability level for non-H atoms. Non-labelled atoms are related to labelled atoms by the symmetry operator: $1 - x, 1 - y, 1 - z$.

3,3,12,12-Tetramethyl-1,5,10,14-tetraoxadispiro[5.2.5.2]hexadecane

Crystal data

C ₁₆ H ₂₈ O ₄	$F_{000} = 312$
$M_r = 284.38$	$D_x = 1.223 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Melting point: 430 K
Hall symbol: -P 2ybc	Mo $K\alpha$ radiation
$a = 12.639 (8) \text{ \AA}$	$\lambda = 0.71073 \text{ \AA}$
$b = 5.838 (4) \text{ \AA}$	Cell parameters from 2071 reflections
$c = 11.179 (7) \text{ \AA}$	$\theta = 3.3\text{--}27.5^\circ$
$\beta = 110.611 (8)^\circ$	$\mu = 0.09 \text{ mm}^{-1}$
	$T = 93 \text{ K}$

supplementary materials

$V = 772.1(9) \text{ \AA}^3$ Block, colourless
 $Z = 2$ $0.43 \times 0.37 \times 0.14 \text{ mm}$

Data collection

Rigaku SPIDER diffractometer
Radiation source: Rotating Anode
Monochromator: graphite
 $T = 93 \text{ K}$
 ω scans
Absorption correction: none
5721 measured reflections
1752 independent reflections

1243 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.048$
 $\theta_{\text{max}} = 27.5^\circ$
 $\theta_{\text{min}} = 3.4^\circ$
 $h = -16 \rightarrow 16$
 $k = -7 \rightarrow 7$
 $l = -13 \rightarrow 14$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.049$
 $wR(F^2) = 0.095$
 $S = 1.00$
1752 reflections
93 parameters
Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map
Hydrogen site location: inferred from neighbouring sites
H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0116P)^2 + 0.36P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} < 0.001$
 $\Delta\rho_{\text{max}} = 0.27 \text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.19 \text{ e \AA}^{-3}$
Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.68024 (9)	0.62454 (18)	0.57453 (10)	0.0219 (3)
O2	0.66546 (9)	0.37232 (18)	0.73088 (10)	0.0221 (3)
C1	0.43232 (13)	0.7060 (3)	0.48564 (15)	0.0224 (4)
H1A	0.3649	0.7851	0.4912	0.027*

H1B	0.4768	0.8194	0.4574	0.027*
C2	0.50381 (13)	0.6171 (3)	0.61780 (15)	0.0220 (4)
H2A	0.5299	0.7478	0.6774	0.026*
H2B	0.4570	0.5166	0.6505	0.026*
C3	0.60572 (13)	0.4836 (3)	0.61294 (15)	0.0204 (4)
C4	0.74097 (14)	0.7887 (3)	0.66956 (15)	0.0232 (4)
H4A	0.6868	0.8977	0.6843	0.028*
H4B	0.7928	0.8768	0.6380	0.028*
C5	0.80864 (13)	0.6730 (3)	0.79466 (15)	0.0217 (4)
C6	0.72591 (14)	0.5256 (3)	0.83191 (15)	0.0222 (4)
H6A	0.7677	0.4360	0.9093	0.027*
H6B	0.6715	0.6257	0.8526	0.027*
C7	0.85930 (15)	0.8559 (3)	0.89653 (16)	0.0284 (4)
H7A	0.9083	0.9571	0.8691	0.034*
H7B	0.9038	0.7820	0.9773	0.034*
H7C	0.7983	0.9459	0.9085	0.034*
C8	0.90174 (14)	0.5242 (3)	0.77798 (16)	0.0288 (4)
H8A	0.8680	0.4097	0.7113	0.035*
H8B	0.9430	0.4466	0.8587	0.035*
H8C	0.9541	0.6206	0.7532	0.035*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0275 (6)	0.0188 (6)	0.0206 (6)	-0.0030 (5)	0.0099 (5)	-0.0001 (5)
O2	0.0292 (6)	0.0154 (6)	0.0197 (6)	-0.0016 (5)	0.0060 (5)	0.0013 (5)
C1	0.0282 (9)	0.0151 (8)	0.0240 (9)	0.0018 (6)	0.0093 (7)	-0.0016 (7)
C2	0.0262 (9)	0.0184 (8)	0.0217 (9)	0.0008 (6)	0.0087 (7)	-0.0019 (7)
C3	0.0254 (9)	0.0167 (8)	0.0188 (8)	-0.0007 (6)	0.0074 (7)	0.0024 (7)
C4	0.0289 (9)	0.0160 (8)	0.0240 (9)	-0.0042 (7)	0.0083 (7)	0.0000 (7)
C5	0.0258 (9)	0.0182 (8)	0.0209 (9)	-0.0010 (7)	0.0083 (7)	0.0002 (7)
C6	0.0277 (9)	0.0207 (8)	0.0175 (9)	0.0002 (7)	0.0073 (7)	0.0003 (7)
C7	0.0331 (10)	0.0254 (9)	0.0253 (10)	-0.0041 (8)	0.0084 (8)	-0.0010 (8)
C8	0.0281 (9)	0.0291 (10)	0.0293 (10)	0.0016 (7)	0.0105 (8)	0.0008 (8)

Geometric parameters (\AA , $^\circ$)

O1—C3	1.4258 (19)	C4—H4A	0.990
O1—C4	1.4362 (19)	C4—H4B	0.990
O2—C3	1.4248 (19)	C5—C6	1.521 (2)
O2—C6	1.4328 (19)	C5—C8	1.526 (2)
C1—C3 ⁱ	1.516 (2)	C5—C7	1.527 (2)
C1—C2	1.526 (2)	C6—H6A	0.990
C1—H1A	0.990	C6—H6B	0.990
C1—H1B	0.990	C7—H7A	0.980
C2—C3	1.523 (2)	C7—H7B	0.980
C2—H2A	0.990	C7—H7C	0.980
C2—H2B	0.990	C8—H8A	0.980

supplementary materials

C3—C1 ⁱ	1.516 (2)	C8—H8B	0.980
C4—C5	1.517 (2)	C8—H8C	0.980
C3—O1—C4	113.58 (12)	H4A—C4—H4B	108.0
C3—O2—C6	113.98 (12)	C4—C5—C6	106.59 (13)
C3 ⁱ —C1—C2	112.55 (13)	C4—C5—C8	110.46 (14)
C3 ⁱ —C1—H1A	109.1	C6—C5—C8	110.18 (14)
C2—C1—H1A	109.1	C4—C5—C7	109.17 (14)
C3 ⁱ —C1—H1B	109.1	C6—C5—C7	109.84 (14)
C2—C1—H1B	109.1	C8—C5—C7	110.50 (14)
H1A—C1—H1B	107.8	O2—C6—C5	111.39 (13)
C3—C2—C1	111.07 (13)	O2—C6—H6A	109.3
C3—C2—H2A	109.4	C5—C6—H6A	109.3
C1—C2—H2A	109.4	O2—C6—H6B	109.3
C3—C2—H2B	109.4	C5—C6—H6B	109.3
C1—C2—H2B	109.4	H6A—C6—H6B	108.0
H2A—C2—H2B	108.0	C5—C7—H7A	109.5
O2—C3—O1	110.54 (12)	C5—C7—H7B	109.5
O2—C3—C1 ⁱ	105.59 (13)	H7A—C7—H7B	109.5
O1—C3—C1 ⁱ	106.05 (13)	C5—C7—H7C	109.5
O2—C3—C2	112.32 (13)	H7A—C7—H7C	109.5
O1—C3—C2	111.86 (13)	H7B—C7—H7C	109.5
C1 ⁱ —C3—C2	110.12 (13)	C5—C8—H8A	109.5
O1—C4—C5	111.45 (13)	C5—C8—H8B	109.5
O1—C4—H4A	109.3	H8A—C8—H8B	109.5
C5—C4—H4A	109.3	C5—C8—H8C	109.5
O1—C4—H4B	109.3	H8A—C8—H8C	109.5
C5—C4—H4B	109.3	H8B—C8—H8C	109.5
C3 ⁱ —C1—C2—C3	56.01 (19)	C1—C2—C3—C1 ⁱ	-54.63 (18)
C6—O2—C3—O1	54.99 (16)	C3—O1—C4—C5	57.12 (17)
C6—O2—C3—C1 ⁱ	169.25 (12)	O1—C4—C5—C6	-54.38 (17)
C6—O2—C3—C2	-70.72 (16)	O1—C4—C5—C8	65.32 (17)
C4—O1—C3—O2	-55.06 (16)	O1—C4—C5—C7	-172.97 (13)
C4—O1—C3—C1 ⁱ	-169.04 (12)	C3—O2—C6—C5	-56.61 (17)
C4—O1—C3—C2	70.90 (16)	C4—C5—C6—O2	54.05 (17)
C1—C2—C3—O2	-172.00 (12)	C8—C5—C6—O2	-65.83 (17)
C1—C2—C3—O1	63.02 (16)	C7—C5—C6—O2	172.21 (13)

Symmetry codes: (i) $-x+1, -y+1, -z+1$.

Fig. 1

