

4-Bromo-3-{N-[2-(3,4-dimethoxyphenyl)-ethyl]-N-methylsulfamoyl}-5-methylbenzoic acid monohydrate

C. Suneel Manohar Babu,^a Helen P. Kavitha,^b Jasmine P. Vennila,^c G. Chakkavarthi^d and V. Manivannan^{e*}

^aNicholas Piramal Research Centre, Nicholas Piramal India Limited, Mumbai 400 063, India, ^bDepartment of Chemistry, SRM University, Ramapuram, Chennai 600 089, India, ^cDepartment of Physics, Panimalar Institute of Technology, Chennai 600 095, India, ^dDepartment of Physics, CPCL Polytechnic College, Chennai 600 068, India, and ^eDepartment of Research and Development, PRIST University, Vallam, Thanjavur 613 403, Tamil Nadu, India

Correspondence e-mail: manivan_1999@yahoo.com


Received 30 May 2009; accepted 10 June 2009

Key indicators: single-crystal X-ray study; $T = 295\text{ K}$; mean $\sigma(\text{C-C}) = 0.005\text{ \AA}$; R factor = 0.040; wR factor = 0.107; data-to-parameter ratio = 16.8.

In the title compound, $\text{C}_{19}\text{H}_{22}\text{BrNO}_6\text{S}\cdot\text{H}_2\text{O}$, the dihedral angle between the planes of the two benzene rings is $3.1(1)^\circ$. These rings are stacked over one another with their centroids separated by $3.769(2)\text{ \AA}$, indicating weak $\pi-\pi$ interactions. In the crystal structure, molecules are linked by $\text{O}-\text{H}\cdots\text{O}$ and $\text{O}-\text{H}\cdots(\text{O},\text{O})$ hydrogen bonds involving the water molecule, forming a two-dimensional network parallel to (001).

Related literature

For the biological activity of sulfonamides, see: Cates (1986); Steele & Beran (1984); Benedetti (1987); Mengelers *et al.* (1997). For related structures, see: Babu *et al.* (2009*a,b*); Shad *et al.* (2009); For graph-set notation, see: Bernstein *et al.* (1995)



Experimental

Crystal data

$\text{C}_{19}\text{H}_{22}\text{BrNO}_6\text{S}\cdot\text{H}_2\text{O}$

$M_r = 490.36$

Orthorhombic, $P2_12_12_1$

$a = 7.7938(2)\text{ \AA}$

$b = 7.8280(2)\text{ \AA}$

$c = 34.6549(8)\text{ \AA}$

$V = 2114.29(9)\text{ \AA}^3$

$Z = 4$

Mo $K\alpha$ radiation

$\mu = 2.08\text{ mm}^{-1}$

$T = 295\text{ K}$

$0.15 \times 0.12 \times 0.10\text{ mm}$

Data collection

Bruker Kappa APEXII area-detector diffractometer

Absorption correction: multi-scan (*SADABS*; Sheldrick, 1996)

$T_{\min} = 0.745$, $T_{\max} = 0.819$

12583 measured reflections

4619 independent reflections

3427 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.028$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.040$

$wR(F^2) = 0.107$

$S = 1.05$

4619 reflections

275 parameters

3 restraints

H atoms treated by a mixture of independent and constrained refinement

$\Delta\rho_{\max} = 0.36\text{ e \AA}^{-3}$

$\Delta\rho_{\min} = -0.30\text{ e \AA}^{-3}$

Absolute structure: Flack (1983), 1915 Friedel pairs

Flack parameter: -0.004 (10)

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O2—H2A \cdots O1W	0.82	1.73	2.535 (4)	169
O1W—H1W \cdots O1 ⁱ	0.79 (3)	1.97 (4)	2.732 (5)	162 (6)
O1W—H2W \cdots O5 ⁱⁱ	0.75 (3)	2.46 (4)	3.066 (5)	139 (5)
O1W—H2W \cdots O6 ⁱⁱ	0.75 (3)	2.14 (4)	2.828 (5)	154 (5)

Symmetry codes: (i) $-x + 1, y - \frac{1}{2}, -z + \frac{1}{2}$; (ii) $x + 1, y - 1, z$.

Data collection: *APEX2* (Bruker, 2004); cell refinement: *SAINT* (Bruker, 2004); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2009); software used to prepare material for publication: *SHELXL97*.

The authors acknowledge the Sophisticated Analytical Instrument Facility, Indian Institute of Technology-Madras, Chennai, for the data collection.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CI2818).

References

- Babu, C. S. M., Kavitha, H. P., Arulmozhi, R., Vennila, J. P. & Manivannan, V. (2009*a*). *Acta Cryst. E65*, o1098.
Babu, C. S. M., Kavitha, H. P., Kavipriya, R., Vennila, J. P. & Manivannan, V. (2009*b*). *Acta Cryst. E65*, o921.
Benedetti, P. G. D. (1987). *Advances in Drug Research*, Vol. 16, edited by B. Testa, pp. 227–279. London, New York: Academic Press.
Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
Bruker (2004). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
Cates, L. A. (1986). *Sulfa Drugs. Handbook of Chemotherapeutic agents*, Vol. I, edited by M. Verderame, pp. 1–29. Boca Raton, Florida: CRC Press.
Flack, H. D. (1983). *Acta Cryst. A39*, 876–881.
Mengelers, M. J., Hougee, P. E., Jansson, L. H. & Van Miert, A. S. (1997). *J. Vet. Pharmacol. Ther.* **20**, 276–283.
Shad, H. A., Tahir, M. N. & Chohan, Z. H. (2009). *Acta Cryst. E65*, o98–o99.
Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
Spek, A. L. (2009). *Acta Cryst. D65*, 148–155.
Steele, J. H. & Beran, G. W. (1984). *Perspectives in the Uses of Antibiotics and Sulfonamides*, in *Handbook Series in Zoonoses, Section D: Antibiotics, Sulfonamides and Public Health*, Vol. 1, edited by J. H. Steele & G. W. Beran, p. 9. Boca Raton, Florida: CRC Press.

supplementary materials

Acta Cryst. (2009). E65, o1568 [doi:10.1107/S1600536809022041]

4-Bromo-3-{N-[2-(3,4-dimethoxyphenyl)ethyl]-N-methylsulfamoyl}-5-methylbenzoic acid monohydrate

C. S. M. Babu, H. P. Kavitha, J. P. Vennila, G. Chakkavarthi and V. Manivannan

Comment

Sulfonamides are a class of anti-microbial agents that have seen extensive use in medicine. They are the first agents to be used for the treatment of bacterial infection (Cates, 1986). Sulfonamides are used to treat atrophic rhinitis in swine and heart water in cattle and many other diseases in a variety of animals (Steele & Beran, 1984). In addition sulfonamides have a variety of biological activities such as antibacterial, antimalarial and antileprotic agents (Benedetti, 1987; Mengelers *et al.*, 1997).

The geometric parameters of the title molecule agree well with those reported for similar structures (Babu *et al.*, 2009*a,b*; Shad *et al.*, 2009). In the molecular structure, the two benzene rings are stacked over one another with their centroids separated by 3.769 (2) Å, indicating a weak π - π interaction. The dihedral angle between the two benzene rings is 3.1 (1) $^{\circ}$. A distorted tetrahedral geometry [O3—S1—O4 = 118.7 (2) $^{\circ}$ and O3—S1—N1 = 108.8 (2) $^{\circ}$] is observed around S1 atom.

In the crystal structure, O—H \cdots O hydrogen bonds (Table 1) link the molecules into a two-dimensional network parallel to the (0 0 1). The O1W—H2W \cdots O5 and O1W—H2W \cdots O6 bifurcated donor bonds generate an $R_1^{2}(5)$ ring motif (Bernstein *et al.*, 1995).

Experimental

A solution of 2-(3,4-dimethoxyphenyl)-*N*-methyl ethanamine (1 g, 0.0051 mol) in ethyl acetate (20 ml) was charged into a round bottom flask equipped with thermometer pocket, condenser and guard tube. Then, pyridine (0.81 g, 0.0102 mol) was added at 25–30°C. After 5 min stirring, 2-bromo-5-(chlorosulfonyl)-3-methylbenzoic acid (0.96 g, 0.00307 mol) was charged into the reaction mass and heated to 45–50°C and maintained for 5–6 h. The completion of the reaction was checked by thin layer chromatography (1:1 hexane-ethyl acetate) and the reaction mass was cooled to 25–30 °C and quenched with 20 ml of water. Then, the aqueous layer was separated and the ethyl acetate layer was washed twice with 10% sodium chloride solution and dried over 2 g of anhydrous sodium sulfate. The solvent was distilled under vacuum at 35–40°C and the crude compound isolated. The crude compound was purified through column chromatography using hexane and ethyl acetate as eluents.

Refinement

Atoms H1W and H2W were located in a difference difference Fourier map and refined freely. Other H atoms were positioned geometrically and refined using a riding model with C-H = 0.93 Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ for aromatic, C-H = 0.97 Å and $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$ for methylene, C-H = 0.96 Å and $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$ for methyl and O-H = 0.82 Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{O})$ for water H atoms. The O1W-H1W and O1W-H2W distances were restrained to 0.82 (4) Å and atoms O2 and C7 were subjected to a rigid bond restraint.

supplementary materials

Figures




Fig. 1. The molecular structure of the title compound, with atom labels and 30% probability displacement ellipsoids for non-H atoms. The dashed line denotes a hydrogen bond.

4-Bromo-3-{N-[2-(3,4-dimethoxyphenyl)ethyl]-N-methylsulfamoyl}-5-methylbenzoic acid monohydrate

Crystal data

C ₁₉ H ₂₂ BrNO ₆ S·H ₂ O	$F_{000} = 1008$
$M_r = 490.36$	$D_x = 1.540 \text{ Mg m}^{-3}$
Orthorhombic, P2 ₁ 2 ₁ 2 ₁	Mo K α radiation
Hall symbol: P 2ac 2ab	$\lambda = 0.71073 \text{ \AA}$
$a = 7.7938 (2) \text{ \AA}$	Cell parameters from 12583 reflections
$b = 7.8280 (2) \text{ \AA}$	$\theta = 1.2\text{--}27.2^\circ$
$c = 34.6549 (8) \text{ \AA}$	$\mu = 2.08 \text{ mm}^{-1}$
$V = 2114.29 (9) \text{ \AA}^3$	$T = 295 \text{ K}$
$Z = 4$	Block, colourless
	$0.15 \times 0.12 \times 0.10 \text{ mm}$

Data collection

Bruker Kappa APEXII area-detector diffractometer	4619 independent reflections
Radiation source: fine-focus sealed tube	3427 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.028$
$T = 295 \text{ K}$	$\theta_{\text{max}} = 27.2^\circ$
ω and φ scans	$\theta_{\text{min}} = 1.2^\circ$
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	$h = -10 \rightarrow 6$
$T_{\text{min}} = 0.745$, $T_{\text{max}} = 0.819$	$k = -9 \rightarrow 10$
12583 measured reflections	$l = -44 \rightarrow 44$

Refinement

Refinement on F^2	Hydrogen site location: inferred from neighbouring sites
Least-squares matrix: full	H atoms treated by a mixture of independent and constrained refinement
$R[F^2 > 2\sigma(F^2)] = 0.040$	$w = 1/[\sigma^2(F_o^2) + (0.0512P)^2]$
$wR(F^2) = 0.107$	where $P = (F_o^2 + 2F_c^2)/3$
	$(\Delta/\sigma)_{\text{max}} = 0.001$

$S = 1.05$
 4619 reflections
 275 parameters
 3 restraints
 Primary atom site location: structure-invariant direct methods
 Secondary atom site location: difference Fourier map

$\Delta\rho_{\max} = 0.36 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.29 \text{ e \AA}^{-3}$
 Extinction correction: none
 Absolute structure: Flack (1983), 1915 Friedel pairs
 Flack parameter: -0.004 (10)

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.4622 (4)	-0.0786 (5)	0.33480 (10)	0.0437 (9)
C2	0.4708 (4)	-0.1063 (5)	0.37399 (10)	0.0409 (8)
H2	0.5171	-0.2075	0.3835	0.049*
C3	0.4107 (4)	0.0161 (4)	0.39913 (9)	0.0379 (8)
C4	0.3406 (4)	0.1659 (5)	0.38498 (10)	0.0428 (8)
C5	0.3306 (5)	0.1973 (5)	0.34527 (11)	0.0490 (9)
C6	0.3940 (5)	0.0735 (5)	0.32070 (11)	0.0495 (9)
H6	0.3910	0.0921	0.2942	0.059*
C7	0.5272 (5)	-0.2101 (5)	0.30749 (10)	0.0500 (9)
C8	0.2547 (7)	0.3601 (5)	0.32931 (12)	0.0714 (12)
H8A	0.1351	0.3657	0.3358	0.107*
H8B	0.2674	0.3615	0.3018	0.107*
H8C	0.3133	0.4566	0.3402	0.107*
C9	0.1792 (6)	0.0519 (6)	0.49724 (11)	0.0665 (12)
H9A	0.0653	0.0908	0.4915	0.100*
H9B	0.2535	0.1485	0.5007	0.100*
H9C	0.1774	-0.0148	0.5205	0.100*
C10	0.1537 (5)	-0.2158 (5)	0.45884 (11)	0.0529 (10)
H10A	0.1564	-0.2827	0.4824	0.063*
H10B	0.2153	-0.2791	0.4392	0.063*
C11	-0.0297 (5)	-0.1940 (6)	0.44630 (12)	0.0606 (11)
H11A	-0.0842	-0.3054	0.4459	0.073*
H11B	-0.0889	-0.1251	0.4654	0.073*
C12	-0.0524 (5)	-0.1125 (5)	0.40751 (11)	0.0511 (10)
C13	-0.0043 (5)	-0.1982 (6)	0.37419 (13)	0.0625 (11)
H13	0.0431	-0.3069	0.3762	0.075*
C14	-0.0251 (5)	-0.1263 (6)	0.33830 (12)	0.0579 (11)
H14	0.0064	-0.1866	0.3163	0.069*
C15	-0.0922 (5)	0.0336 (6)	0.33519 (10)	0.0527 (10)
C16	-0.1403 (4)	0.1245 (5)	0.36809 (11)	0.0484 (9)
C17	-0.1194 (4)	0.0509 (5)	0.40371 (10)	0.0481 (9)
H17	-0.1507	0.1116	0.4257	0.058*
C18	-0.0805 (8)	0.0331 (7)	0.26626 (12)	0.0896 (16)
H18A	0.0405	0.0099	0.2655	0.134*
H18B	-0.1111	0.1043	0.2448	0.134*
H18C	-0.1428	-0.0725	0.2648	0.134*
C19	-0.2491 (6)	0.3849 (5)	0.39391 (12)	0.0633 (10)

supplementary materials

H19A	-0.3342	0.3264	0.4089	0.095*
H19B	-0.2942	0.4926	0.3854	0.095*
H19C	-0.1489	0.4042	0.4094	0.095*
N1	0.2420 (5)	-0.0523 (4)	0.46561 (7)	0.0491 (7)
O1	0.5144 (5)	-0.1988 (5)	0.27278 (7)	0.0826 (10)
O2	0.5954 (4)	-0.3405 (4)	0.32450 (7)	0.0634 (7)
H2A	0.6330	-0.4068	0.3082	0.095*
O3	0.5148 (4)	0.1079 (4)	0.46804 (8)	0.0690 (8)
O4	0.5175 (4)	-0.1965 (4)	0.45044 (7)	0.0609 (8)
O5	-0.2047 (4)	0.2842 (4)	0.36146 (7)	0.0624 (8)
O6	-0.1214 (4)	0.1172 (4)	0.30090 (7)	0.0683 (8)
S1	0.43421 (12)	-0.03348 (13)	0.44937 (2)	0.0467 (2)
Br1	0.25758 (6)	0.33631 (5)	0.418916 (12)	0.06382 (16)
O1W	0.7092 (5)	-0.5755 (5)	0.28115 (11)	0.0720 (10)
H2W	0.749 (6)	-0.644 (5)	0.2931 (13)	0.074 (18)*
H1W	0.641 (6)	-0.627 (7)	0.2690 (14)	0.09 (2)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0415 (18)	0.050 (2)	0.0399 (18)	-0.0069 (17)	-0.0028 (15)	-0.0013 (17)
C2	0.0412 (17)	0.040 (2)	0.0419 (18)	-0.0027 (15)	-0.0011 (15)	0.0009 (17)
C3	0.0372 (16)	0.042 (2)	0.0350 (16)	-0.0057 (15)	-0.0030 (14)	0.0002 (16)
C4	0.0365 (16)	0.0367 (19)	0.055 (2)	-0.0073 (16)	0.0003 (15)	-0.0056 (18)
C5	0.0495 (19)	0.042 (2)	0.056 (2)	-0.0081 (17)	-0.0055 (17)	0.0097 (19)
C6	0.057 (2)	0.054 (2)	0.0380 (18)	-0.0078 (19)	-0.0009 (16)	0.0117 (19)
C7	0.055 (2)	0.057 (2)	0.038 (2)	-0.0041 (16)	0.0011 (16)	-0.0068 (16)
C8	0.084 (3)	0.059 (3)	0.072 (3)	0.001 (3)	-0.001 (3)	0.021 (2)
C9	0.089 (3)	0.063 (3)	0.048 (2)	0.005 (2)	0.016 (2)	-0.013 (2)
C10	0.073 (3)	0.043 (2)	0.043 (2)	-0.0030 (19)	0.0064 (18)	0.0065 (17)
C11	0.061 (2)	0.064 (3)	0.057 (2)	-0.008 (2)	0.008 (2)	0.011 (2)
C12	0.0433 (18)	0.058 (3)	0.052 (2)	-0.0118 (18)	0.0007 (18)	-0.0030 (19)
C13	0.063 (2)	0.056 (3)	0.068 (3)	-0.003 (2)	0.003 (2)	-0.011 (2)
C14	0.063 (2)	0.060 (3)	0.050 (2)	0.000 (2)	0.0018 (19)	-0.015 (2)
C15	0.051 (2)	0.066 (3)	0.0408 (19)	0.000 (2)	-0.0015 (16)	-0.010 (2)
C16	0.0401 (18)	0.057 (3)	0.048 (2)	0.0007 (17)	-0.0030 (16)	-0.012 (2)
C17	0.0434 (19)	0.063 (3)	0.0382 (18)	-0.0067 (18)	0.0028 (15)	-0.0125 (19)
C18	0.127 (4)	0.097 (4)	0.045 (2)	0.009 (4)	0.014 (3)	-0.014 (3)
C19	0.066 (2)	0.063 (2)	0.062 (2)	0.007 (3)	0.001 (2)	-0.014 (2)
N1	0.0617 (17)	0.0490 (16)	0.0367 (14)	-0.0005 (19)	0.0051 (16)	-0.0077 (13)
O1	0.117 (3)	0.096 (3)	0.0341 (14)	0.021 (2)	-0.0022 (16)	-0.0001 (15)
O2	0.085 (2)	0.0635 (18)	0.0413 (13)	0.0139 (16)	-0.0039 (13)	-0.0106 (13)
O3	0.0662 (17)	0.086 (2)	0.0551 (15)	-0.0162 (16)	-0.0150 (13)	-0.0185 (16)
O4	0.0693 (17)	0.073 (2)	0.0410 (13)	0.0241 (15)	-0.0093 (13)	0.0071 (15)
O5	0.077 (2)	0.0691 (18)	0.0412 (14)	0.0187 (15)	0.0002 (13)	-0.0074 (13)
O6	0.087 (2)	0.078 (2)	0.0400 (14)	0.0153 (17)	-0.0033 (14)	-0.0105 (15)
S1	0.0499 (5)	0.0569 (6)	0.0334 (4)	0.0032 (5)	-0.0070 (4)	-0.0053 (4)
Br1	0.0684 (3)	0.0479 (2)	0.0752 (3)	0.0062 (2)	0.0019 (3)	-0.01215 (19)

O1W	0.089 (3)	0.072 (2)	0.0551 (19)	0.017 (2)	-0.0025 (18)	-0.0076 (19)
-----	-----------	-----------	-------------	-----------	--------------	--------------

Geometric parameters (\AA , $^\circ$)

C1—C2	1.377 (5)	C11—H11B	0.97
C1—C6	1.392 (6)	C12—C13	1.387 (6)
C1—C7	1.487 (5)	C12—C17	1.388 (6)
C2—C3	1.377 (5)	C13—C14	1.375 (6)
C2—H2	0.93	C13—H13	0.93
C3—C4	1.383 (5)	C14—C15	1.361 (6)
C3—S1	1.793 (3)	C14—H14	0.93
C4—C5	1.400 (5)	C15—O6	1.375 (5)
C4—Br1	1.893 (4)	C15—C16	1.395 (5)
C5—C6	1.381 (6)	C16—O5	1.366 (5)
C5—C8	1.510 (5)	C16—C17	1.372 (5)
C6—H6	0.93	C17—H17	0.93
C7—O1	1.210 (4)	C18—O6	1.406 (5)
C7—O2	1.293 (5)	C18—H18A	0.96
C8—H8A	0.96	C18—H18B	0.96
C8—H8B	0.96	C18—H18C	0.96
C8—H8C	0.96	C19—O5	1.417 (5)
C9—N1	1.451 (5)	C19—H19A	0.96
C9—H9A	0.96	C19—H19B	0.96
C9—H9B	0.96	C19—H19C	0.96
C9—H9C	0.96	N1—S1	1.607 (4)
C10—N1	1.472 (5)	O2—H2A	0.82
C10—C11	1.504 (6)	O3—S1	1.428 (3)
C10—H10A	0.97	O4—S1	1.432 (3)
C10—H10B	0.97	O1W—H2W	0.75 (3)
C11—C12	1.498 (6)	O1W—H1W	0.79 (3)
C11—H11A	0.97		
C2—C1—C6	120.0 (3)	H11A—C11—H11B	107.5
C2—C1—C7	120.1 (3)	C13—C12—C17	117.9 (4)
C6—C1—C7	119.9 (3)	C13—C12—C11	120.6 (4)
C3—C2—C1	119.9 (3)	C17—C12—C11	121.5 (4)
C3—C2—H2	120.1	C14—C13—C12	121.5 (4)
C1—C2—H2	120.1	C14—C13—H13	119.2
C2—C3—C4	120.0 (3)	C12—C13—H13	119.2
C2—C3—S1	115.4 (3)	C15—C14—C13	119.6 (4)
C4—C3—S1	124.6 (3)	C15—C14—H14	120.2
C3—C4—C5	121.3 (3)	C13—C14—H14	120.2
C3—C4—Br1	120.8 (3)	C14—C15—O6	124.7 (4)
C5—C4—Br1	117.9 (3)	C14—C15—C16	120.5 (4)
C6—C5—C4	117.6 (3)	O6—C15—C16	114.8 (4)
C6—C5—C8	120.4 (4)	O5—C16—C17	125.4 (3)
C4—C5—C8	122.0 (4)	O5—C16—C15	115.3 (3)
C5—C6—C1	121.3 (3)	C17—C16—C15	119.3 (4)
C5—C6—H6	119.3	C16—C17—C12	121.1 (4)
C1—C6—H6	119.3	C16—C17—H17	119.4

supplementary materials

O1—C7—O2	123.0 (4)	C12—C17—H17	119.4
O1—C7—C1	123.6 (4)	O6—C18—H18A	109.5
O2—C7—C1	113.4 (3)	O6—C18—H18B	109.5
C5—C8—H8A	109.5	H18A—C18—H18B	109.5
C5—C8—H8B	109.5	O6—C18—H18C	109.5
H8A—C8—H8B	109.5	H18A—C18—H18C	109.5
C5—C8—H8C	109.5	H18B—C18—H18C	109.5
H8A—C8—H8C	109.5	O5—C19—H19A	109.5
H8B—C8—H8C	109.5	O5—C19—H19B	109.5
N1—C9—H9A	109.5	H19A—C19—H19B	109.5
N1—C9—H9B	109.5	O5—C19—H19C	109.5
H9A—C9—H9B	109.5	H19A—C19—H19C	109.5
N1—C9—H9C	109.5	H19B—C19—H19C	109.5
H9A—C9—H9C	109.5	C9—N1—C10	116.8 (3)
H9B—C9—H9C	109.5	C9—N1—S1	121.8 (3)
N1—C10—C11	113.0 (3)	C10—N1—S1	117.3 (2)
N1—C10—H10A	109.0	C7—O2—H2A	109.5
C11—C10—H10A	109.0	C16—O5—C19	117.7 (3)
N1—C10—H10B	109.0	C15—O6—C18	118.5 (4)
C11—C10—H10B	109.0	O3—S1—O4	118.65 (17)
H10A—C10—H10B	107.8	O3—S1—N1	108.82 (16)
C12—C11—C10	114.8 (3)	O4—S1—N1	109.40 (17)
C12—C11—H11A	108.6	O3—S1—C3	108.50 (17)
C10—C11—H11A	108.6	O4—S1—C3	105.31 (16)
C12—C11—H11B	108.6	N1—S1—C3	105.34 (15)
C10—C11—H11B	108.6	H2W—O1W—H1W	102 (6)
C6—C1—C2—C3	0.5 (5)	C13—C14—C15—C16	-0.3 (6)
C7—C1—C2—C3	179.9 (3)	C14—C15—C16—O5	179.5 (4)
C1—C2—C3—C4	0.5 (5)	O6—C15—C16—O5	-1.7 (5)
C1—C2—C3—S1	-178.5 (3)	C14—C15—C16—C17	0.0 (6)
C2—C3—C4—C5	-0.6 (5)	O6—C15—C16—C17	178.8 (3)
S1—C3—C4—C5	178.3 (3)	O5—C16—C17—C12	-179.9 (3)
C2—C3—C4—Br1	-179.8 (2)	C15—C16—C17—C12	-0.5 (5)
S1—C3—C4—Br1	-0.9 (4)	C13—C12—C17—C16	1.1 (5)
C3—C4—C5—C6	-0.2 (5)	C11—C12—C17—C16	-179.8 (3)
Br1—C4—C5—C6	179.0 (3)	C11—C10—N1—C9	65.3 (4)
C3—C4—C5—C8	-179.9 (4)	C11—C10—N1—S1	-136.7 (3)
Br1—C4—C5—C8	-0.7 (5)	C17—C16—O5—C19	2.0 (5)
C4—C5—C6—C1	1.2 (5)	C15—C16—O5—C19	-177.4 (3)
C8—C5—C6—C1	-179.1 (4)	C14—C15—O6—C18	0.3 (6)
C2—C1—C6—C5	-1.4 (5)	C16—C15—O6—C18	-178.5 (4)
C7—C1—C6—C5	179.2 (3)	C9—N1—S1—O3	-6.1 (4)
C2—C1—C7—O1	175.4 (4)	C10—N1—S1—O3	-163.0 (3)
C6—C1—C7—O1	-5.1 (6)	C9—N1—S1—O4	124.9 (3)
C2—C1—C7—O2	-3.0 (5)	C10—N1—S1—O4	-31.9 (3)
C6—C1—C7—O2	176.4 (3)	C9—N1—S1—C3	-122.3 (3)
N1—C10—C11—C12	66.2 (5)	C10—N1—S1—C3	80.9 (3)
C10—C11—C12—C13	68.6 (5)	C2—C3—S1—O3	129.1 (3)
C10—C11—C12—C17	-110.5 (4)	C4—C3—S1—O3	-49.9 (3)

C17—C12—C13—C14	−1.4 (6)	C2—C3—S1—O4	1.1 (3)
C11—C12—C13—C14	179.5 (4)	C4—C3—S1—O4	−177.9 (3)
C12—C13—C14—C15	1.0 (6)	C2—C3—S1—N1	−114.5 (3)
C13—C14—C15—O6	−178.9 (4)	C4—C3—S1—N1	66.5 (3)

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
O2—H2A···O1W	0.82	1.73	2.535 (4)	169
O1W—H1W···O1 ⁱ	0.79 (3)	1.97 (4)	2.732 (5)	162 (6)
O1W—H2W···O5 ⁱⁱ	0.75 (3)	2.46 (4)	3.066 (5)	139 (5)
O1W—H2W···O6 ⁱⁱ	0.75 (3)	2.14 (4)	2.828 (5)	154 (5)

Symmetry codes: (i) $-x+1, y-1/2, -z+1/2$; (ii) $x+1, y-1, z$.

supplementary materials

Fig. 1

