

rac-7-Oxabicyclo[2.2.1]heptane-2,3-dicarboxylic acid–2-amino-1,3,4-thiadiazole–water (1/1/1)

Na Wang, Qiu-Yue Lin* and Yan-Jun Wang

Zhejiang Key Laboratory for Reactive Chemistry on Solid Surfaces, Institute of Physical Chemistry, Zhejiang Normal University, Jinhua, Zhejiang 321004, People's Republic of China, and, College of Chemistry and Life Science, Zhejiang Normal University, Jinhua 321004, Zhejiang, People's Republic of China
Correspondence e-mail: sky51@zjnu.cn


Received 27 May 2009; accepted 9 June 2009

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.046; wR factor = 0.133; data-to-parameter ratio = 16.0.

The title compound, $\text{C}_8\text{H}_{10}\text{O}_5 \cdot \text{C}_2\text{H}_3\text{N}_3\text{S} \cdot \text{H}_2\text{O}$, was synthesized by the reaction of 2-amino-1,3,4-thiadiazole with norcantharidin. The crystal structure is stabilized by $\text{N}-\text{H} \cdots \text{O}$, $\text{N}-\text{H} \cdots \text{N}$, $\text{O}-\text{H} \cdots \text{O}$ and $\text{O}-\text{H} \cdots \text{N}$ hydrogen bonds. In addition, weak $\pi-\pi$ interactions are observed between symmetry-related thiadiazole ring systems [centroid–centroid distance = $3.9110(3)$ Å, interplanar spacing = 3.4845 Å].

Related literature

7-Oxabicyclo[2.2.1]heptane-2,3-dicarboxylic anhydride (norcantharidin) is a lower toxicity anticancer drug, see: Shimi & Zaki (1982).



Experimental

Crystal data

$\text{C}_8\text{H}_{10}\text{O}_5 \cdot \text{C}_2\text{H}_3\text{N}_3\text{S} \cdot \text{H}_2\text{O}$
 $M_r = 305.31$

Monoclinic, $P2_1/n$
 $a = 5.7678(5)$ Å

$b = 18.4267(15)$ Å
 $c = 12.7546(11)$ Å
 $\beta = 101.336(6)^\circ$
 $V = 1329.1(2)$ Å³
 $Z = 4$

Mo $K\alpha$ radiation
 $\mu = 0.27$ mm⁻¹
 $T = 296$ K
 $0.30 \times 0.16 \times 0.09$ mm

Data collection

Bruker APEXII area-detector diffractometer
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)
 $T_{\min} = 0.949$, $T_{\max} = 0.977$

10820 measured reflections
2995 independent reflections
2026 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.040$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.046$
 $wR(F^2) = 0.133$
 $S = 1.05$
2995 reflections
187 parameters
3 restraints

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.30$ e Å⁻³
 $\Delta\rho_{\min} = -0.29$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
$\text{N1}-\text{H1A} \cdots \text{O4}^{\text{ii}}$	0.86	2.11	2.930 (3)	160
$\text{N1}-\text{H1C} \cdots \text{N3}^{\text{iii}}$	0.86	2.15	2.994 (3)	166
$\text{N1}-\text{H1C} \cdots \text{N2}^{\text{ii}}$	0.86	2.69	3.519 (3)	161
$\text{O2}-\text{H2A} \cdots \text{O1W}^{\text{iii}}$	0.82	1.81	2.626 (2)	176
$\text{O5}-\text{H5B} \cdots \text{N2}^{\text{iv}}$	0.82	1.85	2.664 (2)	172
$\text{O1W}-\text{H1WA} \cdots \text{O3}^{\text{v}}$	0.859 (17)	1.910 (17)	2.766 (2)	175 (3)
$\text{O1W}-\text{H1WB} \cdots \text{O4}^{\text{vi}}$	0.819 (17)	2.51 (3)	3.151 (3)	137 (3)
$\text{O1W}-\text{H1WB} \cdots \text{O1}^{\text{vi}}$	0.819 (17)	2.55 (3)	3.061 (3)	122 (3)

Symmetry codes: (i) $x + \frac{1}{2}, -y - \frac{1}{2}, z + \frac{1}{2}$; (ii) $x + 1, y, z$; (iii) $-x + \frac{1}{2}, y - \frac{1}{2}, -z + \frac{1}{2}$; (iv) $x - \frac{1}{2}, -y - \frac{1}{2}, z - \frac{1}{2}$; (v) $-x - \frac{1}{2}, y + \frac{1}{2}, -z + \frac{1}{2}$; (vi) $-x, -y, -z$.

Data collection: APEX2 (Bruker, 2006); cell refinement: SAINT (Bruker, 2006); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXL97.

The authors thank the Natural Science Foundation of Zhejiang Province, China (grant No. Y407301) for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: AT2800).

References

- Bruker (2006). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.
Sheldrick, G. M. (2008). Acta Cryst. A64, 112–122.
Shimi, I. R. & Zaki, Z. (1982). Eur. J. Cancer Clin. Oncol. 18, 785–793.

supplementary materials

Acta Cryst. (2009). E65, o1590 [doi:10.1107/S1600536809021825]

***rac*-7-Oxabicyclo[2.2.1]heptane-2,3-dicarboxylic acid-2-amino-1,3,4-thiadiazole-water (1/1/1)**

N. Wang, Q.-Y. Lin and Y.-J. Wang

Comment

7-Oxabicyclo[2.2.1]heptane-2,3-dicarboxylic anhydride (norcantharidin) derived from cantharidin is a lower toxicity anti-cancer drug (Shimi & Zaki, 1982). The title compound was synthesized by the reaction of 2-amino-1,3,4-thiadiazole with 7-oxabicyclo[2.2.1]heptane-2,3-dicarboxylic anhydride (norcantharidin). In this paper, we reports its structure.

X-ray crystallography measurement confirmed the molecular structure and the atom connectivity for the title compound (Fig. 1). The crystal structure is stabilized by N—H···O, N—H···N, O—H···O and O—H···N hydrogen bonds (Table 1). Further, weak π - π interactions are observed between symmetry related thiadiazole ring systems [centroid-centroid distance of 3.9110 (3) Å and interplanar spacing of 3.4845 Å].

Experimental

7-Oxabicyclo[2.2.1]heptane-2,3-dicarboxylic anhydride and 2-amino-1,3,4-thiadiazole were dissolved in tetrahydrofuran and the mixture was stirred for 6 h at room temperature. The clear solution was left undisturbed for days to give colourless crystals of the compound.

Refinement

The H atoms bonded to C and N atoms were positioned geometrically and refined using ariding model [C—H = 0.93- 0.98 Å, N—H = 0.86 Å and O—H = 0.82 Å and $U_{\text{iso}}(\text{H}) = 1.2$ or $1.5U_{\text{eq}}(\text{C,N,O})$]. The H atoms of the water molecule were located in a difference Fourier maps and refined with O—H distance restraints of 0.85 (2) and $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{O})$.

Figures




Fig. 1. A view of the molecule of (I) showing the atom-labelling scheme with displacement ellipsoids drawn at the 30% probability.

***rac*-7-Oxabicyclo[2.2.1]heptane-2,3-dicarboxylic acid-2-amino-1,3,4-thiadiazole-water (1/1/1)**

Crystal data

$\text{C}_8\text{H}_{10}\text{O}_5 \cdot \text{C}_2\text{H}_3\text{N}_3\text{S} \cdot \text{H}_2\text{O}$

$M_r = 305.31$

Monoclinic, $P2_1/n$

$F_{000} = 640$

$D_x = 1.526 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation

supplementary materials

Hall symbol: -P 2yn	$\lambda = 0.71073 \text{ \AA}$
$a = 5.7678 (5) \text{ \AA}$	Cell parameters from 1905 reflections
$b = 18.4267 (15) \text{ \AA}$	$\theta = 2.0\text{--}27.6^\circ$
$c = 12.7546 (11) \text{ \AA}$	$\mu = 0.27 \text{ mm}^{-1}$
$\beta = 101.336 (6)^\circ$	$T = 296 \text{ K}$
$V = 1329.1 (2) \text{ \AA}^3$	Block, colourless
$Z = 4$	$0.30 \times 0.16 \times 0.09 \text{ mm}$

Data collection

Bruker APEXII area-detector diffractometer	2995 independent reflections
Radiation source: fine-focus sealed tube	2026 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.040$
$T = 296 \text{ K}$	$\theta_{\text{max}} = 27.6^\circ$
ω scans	$\theta_{\text{min}} = 2.0^\circ$
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	$h = -6 \rightarrow 7$
$T_{\text{min}} = 0.949$, $T_{\text{max}} = 0.977$	$k = -24 \rightarrow 23$
10820 measured reflections	$l = -16 \rightarrow 15$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.046$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.133$	$w = 1/[\sigma^2(F_o^2) + (0.062P)^2 + 0.3174P]$
$S = 1.05$	where $P = (F_o^2 + 2F_c^2)/3$
2995 reflections	$(\Delta/\sigma)_{\text{max}} = 0.001$
187 parameters	$\Delta\rho_{\text{max}} = 0.30 \text{ e \AA}^{-3}$
3 restraints	$\Delta\rho_{\text{min}} = -0.29 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -

factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	-0.2976 (5)	0.05357 (14)	0.3265 (2)	0.0537 (7)
H1B	-0.3880	0.0947	0.3053	0.064*
C2	-0.0094 (4)	-0.03835 (13)	0.37115 (19)	0.0401 (6)
C3	-0.7049 (4)	-0.14263 (12)	-0.02775 (19)	0.0371 (5)
H3A	-0.7478	-0.1570	-0.1031	0.044*
C4	-0.4729 (4)	-0.10527 (12)	0.11367 (19)	0.0378 (5)
H4A	-0.3228	-0.0881	0.1563	0.045*
C5	-0.5597 (4)	-0.17663 (11)	0.15371 (18)	0.0308 (5)
H5A	-0.6555	-0.1649	0.2069	0.037*
C6	-0.7289 (4)	-0.20482 (11)	0.05131 (18)	0.0320 (5)
H6A	-0.8916	-0.2080	0.0630	0.038*
C7	-0.8388 (4)	-0.07593 (13)	-0.0002 (2)	0.0436 (6)
H7A	-0.9932	-0.0888	0.0135	0.052*
H7B	-0.8573	-0.0399	-0.0566	0.052*
C8	-0.6732 (4)	-0.04919 (12)	0.1016 (2)	0.0451 (6)
H8A	-0.7506	-0.0499	0.1626	0.054*
H8B	-0.6158	-0.0006	0.0925	0.054*
C9	-0.3734 (4)	-0.23059 (12)	0.20304 (18)	0.0339 (5)
C10	-0.6556 (4)	-0.27519 (12)	0.00717 (19)	0.0357 (5)
S1	-0.00230 (12)	0.04890 (3)	0.32282 (6)	0.0514 (2)
N1	0.1766 (3)	-0.08184 (12)	0.39063 (18)	0.0534 (6)
H1A	0.1631	-0.1250	0.4145	0.064*
H1C	0.3110	-0.0670	0.3794	0.064*
N2	-0.2197 (3)	-0.05840 (10)	0.38689 (17)	0.0431 (5)
N3	-0.3854 (4)	-0.00394 (11)	0.36078 (19)	0.0511 (6)
O1	-0.4621 (3)	-0.12182 (8)	0.00484 (13)	0.0399 (4)
O1W	0.3491 (3)	0.19399 (12)	0.19534 (16)	0.0573 (5)
O2	-0.1540 (3)	-0.20904 (9)	0.21051 (15)	0.0470 (5)
H2A	-0.0641	-0.2405	0.2405	0.071*
O3	-0.4251 (3)	-0.28854 (9)	0.23715 (14)	0.0453 (4)
O4	-0.4568 (3)	-0.28772 (9)	-0.00388 (16)	0.0527 (5)
O5	-0.8343 (3)	-0.31955 (9)	-0.02355 (17)	0.0575 (5)
H5B	-0.7870	-0.3568	-0.0474	0.086*
H1WA	0.213 (4)	0.1986 (18)	0.212 (2)	0.086*
H1WB	0.317 (5)	0.2024 (19)	0.1311 (15)	0.086*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0577 (16)	0.0398 (14)	0.064 (2)	0.0017 (11)	0.0119 (14)	0.0069 (13)
C2	0.0431 (13)	0.0412 (13)	0.0358 (14)	-0.0098 (10)	0.0075 (10)	-0.0022 (10)
C3	0.0419 (13)	0.0369 (12)	0.0311 (13)	0.0077 (9)	0.0038 (10)	-0.0009 (10)

supplementary materials

C4	0.0383 (12)	0.0342 (12)	0.0394 (14)	-0.0035 (9)	0.0041 (10)	0.0000 (10)
C5	0.0310 (11)	0.0334 (11)	0.0289 (12)	0.0019 (8)	0.0081 (9)	0.0004 (9)
C6	0.0280 (11)	0.0316 (11)	0.0371 (14)	0.0013 (8)	0.0081 (9)	-0.0036 (9)
C7	0.0443 (13)	0.0368 (12)	0.0486 (16)	0.0118 (10)	0.0062 (11)	0.0015 (11)
C8	0.0606 (16)	0.0306 (12)	0.0448 (16)	0.0032 (10)	0.0119 (12)	-0.0033 (11)
C9	0.0339 (12)	0.0390 (12)	0.0300 (13)	0.0015 (9)	0.0094 (9)	-0.0002 (10)
C10	0.0366 (13)	0.0334 (12)	0.0365 (14)	0.0016 (9)	0.0058 (10)	-0.0027 (9)
S1	0.0575 (4)	0.0399 (4)	0.0583 (5)	-0.0095 (3)	0.0150 (3)	0.0073 (3)
N1	0.0475 (12)	0.0419 (12)	0.0697 (17)	-0.0015 (9)	0.0089 (11)	0.0107 (11)
N2	0.0455 (11)	0.0354 (10)	0.0480 (13)	-0.0044 (8)	0.0085 (9)	0.0048 (9)
N3	0.0462 (12)	0.0434 (12)	0.0625 (15)	0.0023 (9)	0.0083 (11)	0.0072 (11)
O1	0.0417 (9)	0.0391 (9)	0.0425 (10)	0.0032 (7)	0.0170 (7)	0.0082 (7)
O1W	0.0381 (10)	0.0837 (14)	0.0524 (12)	-0.0079 (9)	0.0140 (9)	-0.0121 (11)
O2	0.0315 (9)	0.0499 (10)	0.0589 (12)	0.0004 (7)	0.0068 (8)	0.0103 (8)
O3	0.0374 (9)	0.0435 (9)	0.0558 (12)	0.0049 (7)	0.0108 (8)	0.0171 (8)
O4	0.0406 (10)	0.0447 (10)	0.0747 (14)	0.0032 (7)	0.0163 (9)	-0.0214 (9)
O5	0.0411 (10)	0.0362 (9)	0.0954 (16)	-0.0057 (7)	0.0137 (9)	-0.0213 (10)

Geometric parameters (Å, °)

C1—N3	1.287 (3)	C6—H6A	0.9800
C1—S1	1.715 (3)	C7—C8	1.534 (3)
C1—H1B	0.9300	C7—H7A	0.9700
C2—N2	1.320 (3)	C7—H7B	0.9700
C2—N1	1.323 (3)	C8—H8A	0.9700
C2—S1	1.725 (2)	C8—H8B	0.9700
C3—O1	1.433 (3)	C9—O3	1.212 (3)
C3—C7	1.529 (3)	C9—O2	1.311 (3)
C3—C6	1.550 (3)	C10—O4	1.205 (3)
C3—H3A	0.9800	C10—O5	1.313 (3)
C4—O1	1.434 (3)	N1—H1A	0.8600
C4—C5	1.530 (3)	N1—H1C	0.8600
C4—C8	1.535 (3)	N2—N3	1.381 (3)
C4—H4A	0.9800	O1W—H1WA	0.859 (17)
C5—C9	1.508 (3)	O1W—H1WB	0.819 (17)
C5—C6	1.558 (3)	O2—H2A	0.8200
C5—H5A	0.9800	O5—H5B	0.8200
C6—C10	1.507 (3)		
N3—C1—S1	115.3 (2)	C5—C6—H6A	110.7
N3—C1—H1B	122.4	C3—C7—C8	101.18 (17)
S1—C1—H1B	122.4	C3—C7—H7A	111.5
N2—C2—N1	122.5 (2)	C8—C7—H7A	111.5
N2—C2—S1	113.71 (18)	C3—C7—H7B	111.5
N1—C2—S1	123.80 (18)	C8—C7—H7B	111.5
O1—C3—C7	103.14 (18)	H7A—C7—H7B	109.4
O1—C3—C6	102.49 (17)	C7—C8—C4	101.53 (18)
C7—C3—C6	109.34 (19)	C7—C8—H8A	111.5
O1—C3—H3A	113.6	C4—C8—H8A	111.5
C7—C3—H3A	113.6	C7—C8—H8B	111.5

C6—C3—H3A	113.6	C4—C8—H8B	111.5
O1—C4—C5	102.67 (17)	H8A—C8—H8B	109.3
O1—C4—C8	102.77 (18)	O3—C9—O2	122.9 (2)
C5—C4—C8	108.84 (18)	O3—C9—C5	121.68 (19)
O1—C4—H4A	113.8	O2—C9—C5	115.39 (19)
C5—C4—H4A	113.8	O4—C10—O5	123.7 (2)
C8—C4—H4A	113.8	O4—C10—C6	123.5 (2)
C9—C5—C4	116.94 (17)	O5—C10—C6	112.67 (18)
C9—C5—C6	114.05 (17)	C1—S1—C2	86.76 (12)
C4—C5—C6	101.48 (17)	C2—N1—H1A	120.0
C9—C5—H5A	108.0	C2—N1—H1C	120.0
C4—C5—H5A	108.0	H1A—N1—H1C	120.0
C6—C5—H5A	108.0	C2—N2—N3	111.89 (19)
C10—C6—C3	108.99 (18)	C1—N3—N2	112.4 (2)
C10—C6—C5	115.12 (17)	C3—O1—C4	96.42 (15)
C3—C6—C5	100.22 (17)	H1WA—O1W—H1WB	101 (2)
C10—C6—H6A	110.7	C9—O2—H2A	109.5
C3—C6—H6A	110.7	C10—O5—H5B	109.5
O1—C4—C5—C9	90.1 (2)	C6—C5—C9—O3	-62.0 (3)
C8—C4—C5—C9	-161.49 (19)	C4—C5—C9—O2	2.8 (3)
O1—C4—C5—C6	-34.65 (19)	C6—C5—C9—O2	120.9 (2)
C8—C4—C5—C6	73.8 (2)	C3—C6—C10—O4	65.8 (3)
O1—C3—C6—C10	-86.2 (2)	C5—C6—C10—O4	-45.8 (3)
C7—C3—C6—C10	164.91 (18)	C3—C6—C10—O5	-110.5 (2)
O1—C3—C6—C5	35.06 (19)	C5—C6—C10—O5	137.9 (2)
C7—C3—C6—C5	-73.9 (2)	N3—C1—S1—C2	0.4 (2)
C9—C5—C6—C10	-10.1 (3)	N2—C2—S1—C1	-0.1 (2)
C4—C5—C6—C10	116.51 (19)	N1—C2—S1—C1	179.7 (2)
C9—C5—C6—C3	-126.86 (18)	N1—C2—N2—N3	-179.9 (2)
C4—C5—C6—C3	-0.22 (19)	S1—C2—N2—N3	-0.1 (3)
O1—C3—C7—C8	-34.6 (2)	S1—C1—N3—N2	-0.5 (3)
C6—C3—C7—C8	73.9 (2)	C2—N2—N3—C1	0.4 (3)
C3—C7—C8—C4	0.3 (2)	C7—C3—O1—C4	56.17 (19)
O1—C4—C8—C7	33.9 (2)	C6—C3—O1—C4	-57.40 (18)
C5—C4—C8—C7	-74.5 (2)	C5—C4—O1—C3	57.35 (18)
C4—C5—C9—O3	179.9 (2)	C8—C4—O1—C3	-55.63 (18)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N1—H1A...O4 ⁱ	0.86	2.11	2.930 (3)	160
N1—H1C...N3 ⁱⁱ	0.86	2.15	2.994 (3)	166
N1—H1C...N2 ⁱⁱ	0.86	2.69	3.519 (3)	161
O2—H2A...O1W ⁱⁱⁱ	0.82	1.81	2.626 (2)	176
O5—H5B...N2 ^{iv}	0.82	1.85	2.664 (2)	172
O1W—H1WA...O3 ^v	0.859 (17)	1.910 (17)	2.766 (2)	175 (3)
O1W—H1WB...O4 ^{vi}	0.819 (17)	2.51 (3)	3.151 (3)	137 (3)

supplementary materials

O1W—H1WB...O1^{vi}

0.819 (17)

2.55 (3)

3.061 (3)

122 (3)

Symmetry codes: (i) $x+1/2, -y-1/2, z+1/2$; (ii) $x+1, y, z$; (iii) $-x+1/2, y-1/2, -z+1/2$; (iv) $x-1/2, -y-1/2, z-1/2$; (v) $-x-1/2, y+1/2, -z+1/2$; (vi) $-x, -y, -z$.

Fig. 1

