

## N-Phenylformamide

B. Thimme Gowda,<sup>a\*</sup> Sabine Foro<sup>b</sup> and Hartmut Fuess<sup>b</sup>

<sup>a</sup>Department of Chemistry, Mangalore University, Mangalagangothri 574 199, Mangalore, India, and <sup>b</sup>Institute of Materials Science, Darmstadt University of Technology, Petersenstrasse 23, D-64287 Darmstadt, Germany  
Correspondence e-mail: gowdabt@yahoo.com


Received 5 June 2009; accepted 13 June 2009

Key indicators: single-crystal X-ray study;  $T = 298$  K; mean  $\sigma(\text{C}-\text{C}) = 0.003$  Å;  $R$  factor = 0.037;  $wR$  factor = 0.115; data-to-parameter ratio = 14.0.

There are two independent molecules in the asymmetric unit of the title compound,  $\text{C}_7\text{H}_7\text{NO}$ . The conformation of the N—H bond in the structure is *syn* to the C=O bond in one of the molecules and *anti* in the other. In the crystal, molecules are packed into chains diagonally in the *ac* plane via N—H $\cdots$ O hydrogen bonds.

## Related literature

For related structures, see: Gowda *et al.* (2006); Brown (1966). For our study of the effect of ring and side chain substitutions on the crystal structures of aromatic amides, see: Gowda *et al.* (2000, 2007, 2009).



## Experimental

## Crystal data

$\text{C}_7\text{H}_7\text{NO}$   
 $M_r = 121.14$   
Monoclinic,  $C2/c$   
 $a = 30.923$  (3) Å

$b = 6.1737$  (6) Å  
 $c = 14.814$  (1) Å  
 $\beta = 113.14$  (1)°  
 $V = 2600.6$  (4) Å<sup>3</sup>

$Z = 16$   
Mo  $K\alpha$  radiation  
 $\mu = 0.08$  mm<sup>-1</sup>

$T = 298$  K  
 $0.48 \times 0.44 \times 0.40$  mm

## Data collection

Oxford Diffraction Xcalibur diffractometer with a Sapphire CCD detector  
Absorption correction: multi-scan (*CrysAlis RED*; Oxford)

Diffraction, 2009)  
 $T_{\min} = 0.966$ ,  $T_{\max} = 0.969$   
8394 measured reflections  
2383 independent reflections  
1679 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.016$

## Refinement

$R[F^2 > 2\sigma(F^2)] = 0.037$   
 $wR(F^2) = 0.115$   
 $S = 1.12$   
2383 reflections  
170 parameters

H atoms treated by a mixture of independent and constrained refinement  
 $\Delta\rho_{\max} = 0.11$  e Å<sup>-3</sup>  
 $\Delta\rho_{\min} = -0.10$  e Å<sup>-3</sup>

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N1}-\text{H1N}\cdots\text{O2}^{\text{i}}$	0.888 (16)	1.936 (16)	2.8239 (17)	178.1 (14)
$\text{N2}-\text{H2N}\cdots\text{O1}^{\text{ii}}$	0.857 (16)	2.007 (16)	2.8637 (17)	177.0 (14)

Symmetry codes: (i)  $-x, y, -z + \frac{1}{2}$ ; (ii)  $x, -y, z - \frac{1}{2}$ .

Data collection: *CrysAlis CCD* (Oxford Diffraction, 2009); cell refinement: *CrysAlis RED* (Oxford Diffraction, 2009); data reduction: *CrysAlis RED*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2009); software used to prepare material for publication: *SHELXL97*.

BTG thanks the Alexander von Humboldt Foundation, Bonn, Germany for the resumption of his research fellowship.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: RK2151).

## References

- Brown, C. J. (1966). *Acta Cryst.* **21**, 442–445.  
Gowda, B. T., Foro, S., Terao, H. & Fuess, H. (2009). *Acta Cryst.* **E65**, o1039.  
Gowda, B. T., Paulus, H. & Fuess, H. (2000). *Z. Naturforsch. Teil A*, **55**, 791–800.  
Gowda, B. T., Paulus, H., Svoboda, I. & Fuess, H. (2007). *Z. Naturforsch. Teil A*, **62**, 331–337.  
Gowda, B. T., Shilpa & Lakshmi, J. K. (2006). *Z. Naturforsch. Teil A*, **61**, 595–599.  
Oxford Diffraction (2009). *CrysAlis CCD* and *CrysAlis RED*. Oxford Diffraction Ltd, Yarnton, England.  
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

**supplementary materials**

*Acta Cryst.* (2009). E65, o1633 [ doi:10.1107/S1600536809022776 ]

## *N*-Phenylformamide

**B. T. Gowda, S. Foro and H. Fues**

### Comment

As part of a study of the effect of ring and side chain substitutions on the crystal structures of aromatic amides (Gowda *et al.*, 2000; 2007; 2009), the structure of *N*-(phenyl)–formamide (**I**) has been determined. The asymmetric unit contains two independent molecules (Fig. 1). The conformation of the N–H bond is *syn* to the C=O bond in the side chain, in one of the molecules and is *anti* in the other, in contrast to the *anti* conformation observed in *N*-(2,6-dichlorophenyl)–formamide (Gowda *et al.*, 2000 and *N*-(phenyl)–acetamide (Brown *et al.*, 1966). The molecules in (**I**) are linked through intermolecular N–H...O hydrogen bonding (Tab. 1) and the chains formed diagonally as viewed in the *ac* plane (Fig. 2).

### Experimental

The purity of the commercial sample (Aldrich Chemicals) was checked by determining its melting point and characterized by recording its infrared and NMR spectra (Gowda *et al.*, 2006). The single crystals used in *X*-ray diffraction studies were grown in ethanol solution by slow evaporation at room temperature.

### Refinement

The H atoms were located in difference map and their positional parameters were refined with C–H = 0.9300 Å with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ . The N-bonded H atoms refined freely.

### Figures




Fig. 1. Molecular structure of (**I**), showing the atom labelling scheme. Displacement ellipsoids are drawn at the 50% probability level. H atoms are represented as a small spheres of arbitrary radii.




Fig. 2. Molecular packing of (I) with hydrogen bonding shown as dashed lines.

### N-Phenylformamide

#### Crystal data

$C_7H_7NO$

$M_r = 121.14$

Monoclinic,  $C2/c$

Hall symbol:  $-C 2yc$

$a = 30.923 (3) \text{ \AA}$

$b = 6.1737 (6) \text{ \AA}$

$c = 14.814 (1) \text{ \AA}$

$\beta = 113.14 (1)^\circ$

$V = 2600.6 (4) \text{ \AA}^3$

$Z = 16$

$F_{000} = 1024$

$D_x = 1.238 \text{ Mg m}^{-3}$

Mo  $K\alpha$  radiation,  $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 2475 reflections

$\theta = 2.7\text{--}28.2^\circ$

$\mu = 0.08 \text{ mm}^{-1}$

$T = 298 \text{ K}$

Needle, colourless

$0.48 \times 0.44 \times 0.40 \text{ mm}$

#### Data collection

Oxford Diffraction Xcalibur  
diffractometer with a Sapphire CCD detector

Radiation source: Fine-focus sealed tube

Monochromator: Graphite

$T = 298 \text{ K}$

$\omega$  and  $\varphi$  scans

Absorption correction: multi-scan  
(CrysAlis RED; Oxford Diffraction, 2009)

$T_{\min} = 0.966$ ,  $T_{\max} = 0.969$

8394 measured reflections

2383 independent reflections

1679 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.016$

$\theta_{\max} = 25.4^\circ$

$\theta_{\min} = 2.8^\circ$

$h = -37 \rightarrow 36$

$k = -7 \rightarrow 7$

$l = -17 \rightarrow 17$

#### Refinement

Refinement on  $F^2$

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.037$

Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent and constrained refinement

$w = 1/[\sigma^2(F_o^2) + (0.0651P)^2]$

$wR(F^2) = 0.115$	where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.12$	$(\Delta/\sigma)_{\max} < 0.001$
2383 reflections	$\Delta\rho_{\max} = 0.11 \text{ e } \text{\AA}^{-3}$
170 parameters	$\Delta\rho_{\min} = -0.10 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: SHELXL97 (Sheldrick, 2008), $F_c^* = kFc[1+0.001xFc^2\lambda^3/\sin(2\theta)]^{-1/4}$
Secondary atom site location: difference Fourier map	Extinction coefficient: 0.0028 (7)

*Special details*

**Experimental.** Absorption correction: CrysAlis RED (Oxford Diffraction, 2009); empirical absorption correction using spherical harmonics, implemented in SCALE3 ABSPACK scaling algorithm.

**Geometry.** All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > 2\sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.09999 (4)	0.21410 (18)	0.62537 (8)	0.0921 (4)
N1	0.04841 (4)	0.34831 (19)	0.47961 (9)	0.0694 (4)
H1N	0.0202 (6)	0.322 (2)	0.4332 (11)	0.083*
C1	0.07229 (4)	0.5289 (2)	0.46365 (9)	0.0561 (3)
C2	0.04734 (5)	0.6657 (3)	0.38753 (10)	0.0727 (4)
H2	0.0159	0.6367	0.3491	0.087*
C3	0.06857 (6)	0.8441 (3)	0.36825 (12)	0.0893 (5)
H3	0.0515	0.9351	0.3164	0.107*
C4	0.11462 (6)	0.8895 (3)	0.42461 (12)	0.0890 (5)
H4	0.1289	1.0119	0.4119	0.107*
C5	0.13948 (6)	0.7535 (3)	0.49972 (12)	0.0828 (5)
H5	0.1709	0.7834	0.5378	0.099*
C6	0.11890 (5)	0.5734 (2)	0.51989 (11)	0.0687 (4)
H6	0.1363	0.4818	0.5712	0.082*
C7	0.06296 (6)	0.2094 (2)	0.55435 (12)	0.0792 (5)
H7	0.0426	0.0966	0.5518	0.095*
O2	0.04140 (4)	0.2563 (2)	0.16582 (9)	0.0996 (4)
N2	0.11861 (4)	0.1904 (2)	0.22825 (8)	0.0628 (3)
H2N	0.1131 (5)	0.067 (3)	0.1994 (10)	0.075*
C8	0.16640 (5)	0.2389 (2)	0.28235 (9)	0.0565 (3)
C9	0.18179 (6)	0.4299 (3)	0.33258 (13)	0.0862 (5)

## supplementary materials

---

H9	0.1603	0.5369	0.3309	0.103*
C10	0.22921 (7)	0.4618 (3)	0.38543 (14)	0.1024 (6)
H10	0.2394	0.5901	0.4202	0.123*
C11	0.26131 (6)	0.3093 (3)	0.38760 (14)	0.0963 (6)
H11	0.2932	0.3324	0.4235	0.116*
C12	0.24605 (6)	0.1230 (3)	0.33668 (13)	0.0919 (5)
H12	0.2678	0.0184	0.3371	0.110*
C13	0.19895 (5)	0.0867 (2)	0.28451 (11)	0.0750 (4)
H13	0.1891	-0.0425	0.2503	0.090*
C14	0.08175 (6)	0.3109 (3)	0.21441 (12)	0.0771 (4)
H14	0.0867	0.4471	0.2436	0.093*

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
O1	0.0813 (8)	0.0791 (7)	0.0881 (8)	-0.0120 (6)	0.0032 (6)	0.0215 (5)
N1	0.0527 (7)	0.0658 (8)	0.0743 (8)	-0.0096 (6)	0.0082 (6)	0.0067 (6)
C1	0.0522 (8)	0.0564 (8)	0.0570 (7)	-0.0001 (6)	0.0186 (6)	-0.0032 (6)
C2	0.0644 (9)	0.0780 (10)	0.0644 (9)	-0.0023 (7)	0.0131 (7)	0.0049 (7)
C3	0.1006 (14)	0.0823 (11)	0.0750 (10)	-0.0058 (10)	0.0236 (9)	0.0207 (9)
C4	0.0993 (14)	0.0849 (12)	0.0826 (11)	-0.0263 (10)	0.0355 (10)	0.0063 (9)
C5	0.0682 (10)	0.0902 (12)	0.0847 (11)	-0.0210 (9)	0.0242 (8)	0.0026 (9)
C6	0.0555 (8)	0.0699 (9)	0.0742 (9)	-0.0033 (7)	0.0184 (7)	0.0063 (7)
C7	0.0714 (10)	0.0648 (9)	0.0897 (11)	-0.0116 (8)	0.0189 (9)	0.0098 (8)
O2	0.0556 (7)	0.1164 (10)	0.1092 (9)	0.0152 (6)	0.0132 (6)	0.0123 (7)
N2	0.0567 (7)	0.0615 (7)	0.0645 (7)	0.0048 (6)	0.0175 (6)	-0.0040 (5)
C8	0.0557 (8)	0.0602 (8)	0.0522 (7)	-0.0013 (6)	0.0197 (6)	0.0026 (6)
C9	0.0790 (12)	0.0746 (10)	0.1005 (12)	-0.0037 (8)	0.0303 (9)	-0.0202 (9)
C10	0.0939 (14)	0.0939 (13)	0.1047 (13)	-0.0311 (11)	0.0233 (11)	-0.0271 (10)
C11	0.0647 (11)	0.1055 (15)	0.0999 (13)	-0.0158 (11)	0.0122 (9)	0.0068 (11)
C12	0.0601 (10)	0.0924 (12)	0.1119 (13)	0.0070 (9)	0.0218 (9)	0.0067 (10)
C13	0.0618 (9)	0.0680 (9)	0.0873 (10)	0.0029 (7)	0.0209 (8)	-0.0064 (7)
C14	0.0667 (11)	0.0739 (10)	0.0865 (11)	0.0132 (8)	0.0256 (9)	0.0104 (8)

### Geometric parameters ( $\text{\AA}$ , $^\circ$ )

O1—C7	1.2132 (17)	O2—C14	1.2178 (18)
N1—C7	1.3314 (19)	N2—C14	1.3082 (19)
N1—C1	1.4076 (17)	N2—C8	1.4089 (17)
N1—H1N	0.888 (16)	N2—H2N	0.857 (16)
C1—C2	1.3771 (19)	C8—C13	1.3684 (19)
C1—C6	1.3797 (18)	C8—C9	1.375 (2)
C2—C3	1.369 (2)	C9—C10	1.378 (2)
C2—H2	0.9300	C9—H9	0.9300
C3—C4	1.367 (2)	C10—C11	1.359 (3)
C3—H3	0.9300	C10—H10	0.9300
C4—C5	1.365 (2)	C11—C12	1.354 (3)
C4—H4	0.9300	C11—H11	0.9300
C5—C6	1.371 (2)	C12—C13	1.373 (2)

C5—H5	0.9300	C12—H12	0.9300
C6—H6	0.9300	C13—H13	0.9300
C7—H7	0.9300	C14—H14	0.9300
C7—N1—C1	128.45 (13)	C14—N2—C8	128.51 (14)
C7—N1—H1N	115.8 (10)	C14—N2—H2N	115.9 (10)
C1—N1—H1N	115.7 (10)	C8—N2—H2N	115.6 (10)
C2—C1—C6	119.24 (13)	C13—C8—C9	118.76 (14)
C2—C1—N1	117.46 (12)	C13—C8—N2	117.69 (12)
C6—C1—N1	123.29 (12)	C9—C8—N2	123.55 (13)
C3—C2—C1	120.27 (14)	C8—C9—C10	119.64 (16)
C3—C2—H2	119.9	C8—C9—H9	120.2
C1—C2—H2	119.9	C10—C9—H9	120.2
C4—C3—C2	120.51 (15)	C11—C10—C9	121.22 (17)
C4—C3—H3	119.7	C11—C10—H10	119.4
C2—C3—H3	119.7	C9—C10—H10	119.4
C5—C4—C3	119.30 (15)	C12—C11—C10	118.93 (17)
C5—C4—H4	120.4	C12—C11—H11	120.5
C3—C4—H4	120.4	C10—C11—H11	120.5
C4—C5—C6	121.06 (15)	C11—C12—C13	120.85 (17)
C4—C5—H5	119.5	C11—C12—H12	119.6
C6—C5—H5	119.5	C13—C12—H12	119.6
C5—C6—C1	119.62 (14)	C8—C13—C12	120.59 (15)
C5—C6—H6	120.2	C8—C13—H13	119.7
C1—C6—H6	120.2	C12—C13—H13	119.7
O1—C7—N1	126.94 (14)	O2—C14—N2	124.22 (16)
O1—C7—H7	116.5	O2—C14—H14	117.9
N1—C7—H7	116.5	N2—C14—H14	117.9
C7—N1—C1—C2	-171.26 (15)	C14—N2—C8—C13	-178.73 (14)
C7—N1—C1—C6	9.0 (2)	C14—N2—C8—C9	1.3 (2)
C6—C1—C2—C3	-0.2 (2)	C13—C8—C9—C10	-1.4 (2)
N1—C1—C2—C3	180.00 (14)	N2—C8—C9—C10	178.58 (15)
C1—C2—C3—C4	-0.5 (3)	C8—C9—C10—C11	1.1 (3)
C2—C3—C4—C5	0.8 (3)	C9—C10—C11—C12	0.0 (3)
C3—C4—C5—C6	-0.5 (3)	C10—C11—C12—C13	-0.8 (3)
C4—C5—C6—C1	-0.2 (2)	C9—C8—C13—C12	0.7 (2)
C2—C1—C6—C5	0.5 (2)	N2—C8—C13—C12	-179.34 (13)
N1—C1—C6—C5	-179.70 (14)	C11—C12—C13—C8	0.4 (3)
C1—N1—C7—O1	1.5 (3)	C8—N2—C14—O2	179.62 (13)

Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ )

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N1—H1N $\cdots$ O2 <sup>i</sup>	0.888 (16)	1.936 (16)	2.8239 (17)	178.1 (14)
N2—H2N $\cdots$ O1 <sup>ii</sup>	0.857 (16)	2.007 (16)	2.8637 (17)	177.0 (14)

Symmetry codes: (i)  $-x, y, -z+1/2$ ; (ii)  $x, -y, z-1/2$ .

Fig. 1






Fig. 2

