

3,3'-Di-*tert*-butyl-5,5'-dimethoxy-biphenyl-2,2'-diol

Zhong-Xiang Du^{a*} and Ling-Zhi Wang^b

^aDepartment of Chemistry, Luoyang Normal University, Luoyang, Henan 471022, People's Republic of China, and ^bEquipment Department, Luoyang Normal University, Luoyang, Henan 471022, People's Republic of China
Correspondence e-mail: dzx6281@126.com


Received 13 June 2009; accepted 16 June 2009

Key indicators: single-crystal X-ray study; $T = 291\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$; R factor = 0.041; wR factor = 0.112; data-to-parameter ratio = 15.8.

The title compound, $\text{C}_{22}\text{H}_{30}\text{O}_4$, displays twofold rotational symmetry. The two benzene rings are almost perpendicular to each other, forming a dihedral angle of $89.8(6)^\circ$. In the crystal, molecules are linked into an extended one-dimensional chain structure *via* intermolecular $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds.

Related literature

For the various methods of preparing di-BHA [a dimer of 3-*tert*-butyl-4-hydroxyanisole], see: Hewgill & Hewitt (1967); Jarl *et al.* (2004); Masahiro *et al.* (2005); Seiichiro *et al.* (2004).



Experimental

Crystal data

$\text{C}_{22}\text{H}_{30}\text{O}_4$
 $M_r = 358.46$

Tetragonal, $I4_1/a$
 $a = 13.4289(8)\text{ \AA}$

$c = 23.127(3)\text{ \AA}$
 $V = 4170.5(6)\text{ \AA}^3$
 $Z = 8$
Mo $K\alpha$ radiation

$\mu = 0.08\text{ mm}^{-1}$
 $T = 291\text{ K}$
 $0.49 \times 0.49 \times 0.38\text{ mm}$

Data collection

Bruker APEXII CCD area-detector diffractometer
Absorption correction: multi-scan (*SADABS*; Sheldrick, 1996)
 $T_{\min} = 0.963$, $T_{\max} = 0.972$

13638 measured reflections
1958 independent reflections
1542 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.025$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.041$
 $wR(F^2) = 0.112$
 $S = 1.04$
1938 reflections

123 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.14\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.13\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O2—H2 \cdots O1 ⁱ	0.82	2.08	2.7592 (15)	140

Symmetry code: (i) $x, y - \frac{1}{2}, -z$.

Data collection: *APEX2* (Bruker, 2004); cell refinement: *SAINT* (Bruker, 2004); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

This work was supported financially by the National Natural Science Foundation of China (No. 20771054).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: AT2819).

References

- Bruker (2004). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
Hewgill, F. R. & Hewitt, D. G. (1967). *J. Chem. Soc. C*, pp. 726–730.
Jarl, I. V., Alison, C. H., Samuel, N., Rafael, S., Allison, M. M., Martin, L., Anthony, L. S., Christian, M. & Dieter, V. (2004). *Adv. Synth. Catal.* **346**, 993–1003.
Masahiro, O., Kanae, T. S., Takao, K., Kentaro, O., Masako, S., Shiro, U., Keiichi, H. & Toyoshige, E. (2005). *Biol. Pharm. Bull.* **28**, 1120–1122.
Seiichiro, F., Mariko, I. & Ichiro, Y. (2004). *Internet J. Mol. Des.* pp. 241–246.
Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.

supplementary materials

Acta Cryst. (2009). E65, o1664 [doi:10.1107/S1600536809023071]

3,3'-Di-*tert*-butyl-5,5'-dimethoxybiphenyl-2,2'-diol

Z.-X. Du and L.-Z. Wang

Comment

In the previous literatures, several methods for preparing di-BHA [a dimer of 3-*tert*-butyl-4-hydroxyanisole (BHA)] have been reported (Hewgill & Hewitt, 1967; Masahiro *et al.*, 2005; Jarl *et al.*, 2004; Seiichiro *et al.*, 2004), but its single-crystal and precise molecular structure has not been investigated so far. Here we describe the structure of the title compound, (I), (Fig. 1).

Di-BHA shows 2-fold rotational symmetry characters, where the 2-fold rotation axis is perpendicular to the C6—C6A bond. The oxygen atoms are almost coplanar with their own benzene ring—the largest deviation from the least-squares plane was found for O1 (or O1A), with an atom-plane distance of 0.017 Å. The two benzene rings have a dihedral angle of 89.8°, indicating that they are almost perpendicular to each other. The phenolic hydroxyl donor and methoxyl acceptor are involved in intermolecular hydrogen bonds and they extend di-BHA molecules into a one-dimensional chain structure along the *b* axis (Table 1, Fig. 2), thus stabilizing di-BHA in the solid state.

Experimental

An easy preparation method improved by Jarl *et al.* (2004) was adopted in our experiment. A solution of [K₃Fe(CN)₆] (0.1 mol, 3.29 g) and KOH (0.1 mol, 5.61 g) in water (100 ml) was prepared and was added dropwise to a solution of 3-*tert*-butyl-4-hydroxyanisole (0.1 mol, 1.80 g) in acetone (10 ml) over 3 h at room temperature. After vigorous agitation, yellow rice-shaped precipitate was obtained and filtered. Then the solid product was extracted with CH₂Cl₂ (3 × 50 ml), and the organic phase was dried over Na₂SO₄. After removal of CH₂Cl₂ under vacuum, a light brown solid was obtained. It turned into white crystal substance after washed with anhydrous ethanol (3 × 50 ml). Dissolve the white crystal substance in CH₂Cl₂ and filter the solution. About 4 days later, colourless block-shaped crystals suitable for X-ray diffraction analysis were appeared by slow evaporation in a yield of 63%. m. p. 510–511 K. Analysis, found: C 73.57, H 8.44%; C₂₂H₃₀O₄ requires: C 73.65, H 8.37%. IR (KBr, ν , cm⁻¹): 3412.6(v O—H), 1594.2, 1455.6(v (C₆H₆), skeleton), 1396.2, 1365.4(ν (CH₃)₃-C, skeleton), 1215.3, 1138.8(vC-O), 784.1(γ(C=C—H)).

Refinement

H atoms bonded to C were positioned geometrically with C—H distance of 0.93–0.96 Å, and treated as riding atoms, with $U_{\text{iso}}(\text{H})=1.2$ or $1.5U_{\text{eq}}(\text{C})$. The O—H hydrogen atom was located in a difference Fourier map and the applied restraint of the O—H distance was 0.820 Å, with $U_{\text{iso}}(\text{H})=1.5U_{\text{eq}}(\text{O})$.

supplementary materials

Figures




Fig. 1. Molecular structure of (I), with displacement ellipsoids drawn at the 25% probability level. Atoms with suffix A are at the symmetry position $(-x, -y + 3/2, z)$.




Fig. 2. The crystal packing of (I), showing hydrogen bonds as dashed lines along b axis. H atoms on C atoms have been omitted.

3,3'-Di-*tert*-butyl-5,5'-dimethoxybiphenyl-2,2'-diol

Crystal data

C ₂₂ H ₃₀ O ₄	Z = 8
$M_r = 358.46$	$F_{000} = 1552$
Tetragonal, $I4_1/a$	$D_x = 1.142 \text{ Mg m}^{-3}$
Hall symbol: -I 4ad	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
$a = 13.4289(8) \text{ \AA}$	Cell parameters from 3994 reflections
$b = 13.4289(8) \text{ \AA}$	$\theta = 3.0\text{--}25.5^\circ$
$c = 23.127(3) \text{ \AA}$	$\mu = 0.08 \text{ mm}^{-1}$
$\alpha = 90^\circ$	$T = 291 \text{ K}$
$\beta = 90^\circ$	Block, colourless
$\gamma = 90^\circ$	$0.49 \times 0.49 \times 0.38 \text{ mm}$
$V = 4170.5(6) \text{ \AA}^3$	

Data collection

Bruker APEXII CCD area-detector diffractometer	1938 independent reflections
Radiation source: fine-focus sealed tube	1542 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.025$
$T = 291 \text{ K}$	$\theta_{\max} = 25.5^\circ$
ϕ and ω scans	$\theta_{\min} = 2.8^\circ$
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	$h = -16 \rightarrow 15$
$T_{\min} = 0.963$, $T_{\max} = 0.972$	$k = -15 \rightarrow 16$
13638 measured reflections	$l = -28 \rightarrow 27$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites

$R[F^2 > 2\sigma(F^2)] = 0.041$	H-atom parameters constrained
$wR(F^2) = 0.112$	$w = 1/[\sigma^2(F_o^2) + (0.0476P)^2 + 2.3626P]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.04$	$(\Delta/\sigma)_{\max} < 0.001$
1938 reflections	$\Delta\rho_{\max} = 0.14 \text{ e } \text{\AA}^{-3}$
123 parameters	$\Delta\rho_{\min} = -0.13 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes)

are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.11953 (11)	0.79586 (10)	0.02439 (6)	0.0416 (4)
C2	0.18758 (11)	0.87570 (10)	0.02195 (6)	0.0416 (4)
C3	0.16557 (11)	0.95179 (11)	-0.01654 (6)	0.0431 (4)
H3	0.2085	1.0060	-0.0188	0.052*
C4	0.08231 (11)	0.95010 (10)	-0.05172 (6)	0.0408 (3)
C5	0.01714 (11)	0.87114 (10)	-0.04926 (6)	0.0417 (4)
H5	-0.0385	0.8696	-0.0732	0.050*
C6	0.03503 (10)	0.79350 (10)	-0.01059 (6)	0.0387 (3)
C7	0.28338 (12)	0.87786 (13)	0.05857 (7)	0.0546 (4)
C8	0.34307 (17)	0.97382 (18)	0.04844 (11)	0.0950 (8)
H8A	0.3018	1.0305	0.0566	0.142*
H8B	0.3647	0.9764	0.0089	0.142*
H8C	0.4001	0.9746	0.0735	0.142*
C9	0.34957 (16)	0.78944 (19)	0.04258 (11)	0.0878 (7)
H9A	0.4062	0.7876	0.0679	0.132*
H9B	0.3718	0.7964	0.0033	0.132*
H9C	0.3123	0.7288	0.0465	0.132*
C10	0.25901 (15)	0.87337 (17)	0.12326 (8)	0.0725 (6)

supplementary materials

H10A	0.2230	0.9322	0.1342	0.109*
H10B	0.3197	0.8695	0.1451	0.109*
H10C	0.2190	0.8157	0.1310	0.109*
C11	0.00097 (17)	1.02543 (14)	-0.13233 (9)	0.0733 (6)
H11A	0.0167	0.9698	-0.1567	0.110*
H11B	0.0028	1.0856	-0.1547	0.110*
H11C	-0.0644	1.0168	-0.1163	0.110*
O1	0.07070 (9)	1.03143 (8)	-0.08749 (5)	0.0586 (3)
O2	0.13978 (10)	0.72001 (8)	0.06230 (6)	0.0665 (4)
H2	0.0997	0.6747	0.0574	0.100*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0462 (8)	0.0351 (7)	0.0437 (8)	-0.0028 (6)	-0.0020 (6)	0.0010 (6)
C2	0.0421 (8)	0.0389 (8)	0.0438 (8)	-0.0058 (6)	-0.0027 (6)	-0.0030 (6)
C3	0.0453 (8)	0.0369 (8)	0.0470 (8)	-0.0119 (6)	0.0005 (7)	-0.0021 (6)
C4	0.0483 (8)	0.0313 (7)	0.0427 (8)	-0.0033 (6)	-0.0002 (6)	0.0021 (6)
C5	0.0409 (8)	0.0403 (8)	0.0440 (8)	-0.0042 (6)	-0.0047 (6)	-0.0015 (6)
C6	0.0393 (8)	0.0338 (7)	0.0428 (8)	-0.0048 (6)	0.0025 (6)	-0.0024 (6)
C7	0.0491 (9)	0.0582 (10)	0.0565 (10)	-0.0083 (8)	-0.0133 (8)	-0.0002 (8)
C8	0.0763 (14)	0.1044 (17)	0.1042 (18)	-0.0472 (13)	-0.0422 (13)	0.0238 (14)
C9	0.0574 (12)	0.1129 (18)	0.0930 (16)	0.0194 (12)	-0.0211 (11)	-0.0159 (14)
C10	0.0739 (13)	0.0843 (14)	0.0595 (11)	-0.0043 (10)	-0.0228 (10)	-0.0031 (10)
C11	0.0946 (15)	0.0568 (11)	0.0686 (12)	-0.0020 (10)	-0.0309 (11)	0.0131 (9)
O1	0.0791 (8)	0.0406 (6)	0.0562 (7)	-0.0134 (5)	-0.0178 (6)	0.0115 (5)
O2	0.0759 (9)	0.0465 (7)	0.0770 (9)	-0.0174 (6)	-0.0289 (7)	0.0210 (6)

Geometric parameters (\AA , $^\circ$)

C1—O2	1.3711 (18)	C8—H8A	0.9600
C1—C6	1.394 (2)	C8—H8B	0.9600
C1—C2	1.410 (2)	C8—H8C	0.9600
C2—C3	1.387 (2)	C9—H9A	0.9600
C2—C7	1.541 (2)	C9—H9B	0.9600
C3—C4	1.383 (2)	C9—H9C	0.9600
C3—H3	0.9300	C10—H10A	0.9600
C4—C5	1.3761 (19)	C10—H10B	0.9600
C4—O1	1.3788 (17)	C10—H10C	0.9600
C5—C6	1.395 (2)	C11—O1	1.400 (2)
C5—H5	0.9300	C11—H11A	0.9600
C6—C6 ⁱ	1.500 (3)	C11—H11B	0.9600
C7—C9	1.529 (3)	C11—H11C	0.9600
C7—C10	1.533 (3)	O2—H2	0.8200
C7—C8	1.535 (3)		
O2—C1—C6	121.03 (13)	C7—C8—H8B	109.5
O2—C1—C2	117.52 (13)	H8A—C8—H8B	109.5
C6—C1—C2	121.45 (13)	C7—C8—H8C	109.5

C3—C2—C1	116.61 (13)	H8A—C8—H8C	109.5
C3—C2—C7	121.12 (13)	H8B—C8—H8C	109.5
C1—C2—C7	122.25 (13)	C7—C9—H9A	109.5
C4—C3—C2	122.53 (13)	C7—C9—H9B	109.5
C4—C3—H3	118.7	H9A—C9—H9B	109.5
C2—C3—H3	118.7	C7—C9—H9C	109.5
C5—C4—O1	124.28 (13)	H9A—C9—H9C	109.5
C5—C4—C3	120.16 (13)	H9B—C9—H9C	109.5
O1—C4—C3	115.55 (12)	C7—C10—H10A	109.5
C4—C5—C6	119.54 (13)	C7—C10—H10B	109.5
C4—C5—H5	120.2	H10A—C10—H10B	109.5
C6—C5—H5	120.2	C7—C10—H10C	109.5
C1—C6—C5	119.70 (12)	H10A—C10—H10C	109.5
C1—C6—C6 ⁱ	121.90 (13)	H10B—C10—H10C	109.5
C5—C6—C6 ⁱ	118.31 (12)	O1—C11—H11A	109.5
C9—C7—C10	109.25 (17)	O1—C11—H11B	109.5
C9—C7—C8	108.15 (18)	H11A—C11—H11B	109.5
C10—C7—C8	107.07 (16)	O1—C11—H11C	109.5
C9—C7—C2	109.75 (14)	H11A—C11—H11C	109.5
C10—C7—C2	110.95 (14)	H11B—C11—H11C	109.5
C8—C7—C2	111.58 (14)	C4—O1—C11	118.33 (12)
C7—C8—H8A	109.5	C1—O2—H2	109.5
O2—C1—C2—C3	-179.81 (14)	O2—C1—C6—C6 ⁱ	-2.8 (2)
C6—C1—C2—C3	0.6 (2)	C2—C1—C6—C6 ⁱ	176.75 (13)
O2—C1—C2—C7	1.9 (2)	C4—C5—C6—C1	-1.1 (2)
C6—C1—C2—C7	-177.67 (14)	C4—C5—C6—C6 ⁱ	-177.55 (13)
C1—C2—C3—C4	-1.0 (2)	C3—C2—C7—C9	-117.04 (18)
C7—C2—C3—C4	177.31 (14)	C1—C2—C7—C9	61.1 (2)
C2—C3—C4—C5	0.3 (2)	C3—C2—C7—C10	122.12 (17)
C2—C3—C4—O1	179.81 (14)	C1—C2—C7—C10	-59.7 (2)
O1—C4—C5—C6	-178.71 (14)	C3—C2—C7—C8	2.8 (2)
C3—C4—C5—C6	0.7 (2)	C1—C2—C7—C8	-179.02 (17)
O2—C1—C6—C5	-179.17 (14)	C5—C4—O1—C11	-13.7 (2)
C2—C1—C6—C5	0.4 (2)	C3—C4—O1—C11	166.78 (16)

Symmetry codes: (i) $-x, -y+3/2, z$.

Hydrogen-bond geometry (\AA , $^\circ$)

$D—\text{H}\cdots A$	$D—\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D—\text{H}\cdots A$
O2—H2 ⁱⁱ —O1 ⁱⁱ	0.82	2.08	2.7592 (15)	140

Symmetry codes: (ii) $x, y-1/2, -z$.

supplementary materials

Fig. 1




Fig. 2

