

## 1-(2-Methylimidazo[1,2-a]pyridin-3-yl)-3,3-bis(methylsulfanyl)prop-2-enone monohydrate

Yvon Bibila Mayaya Bisseyou,<sup>a\*</sup> Drissa Sissouma,<sup>b</sup> Severin D. Goulian Bi,<sup>a</sup> Mahama Ouattara<sup>c</sup> and R. C. A. Yao-Kakou<sup>a</sup>

<sup>a</sup>Laboratoire de Cristallographie et Physique Moléculaire, UFR SSMT, Université de Cocody, 22 BP 582 Abidjan 22, Côte d'Ivoire, <sup>b</sup>Laboratoire de Chimie Organique Structurale, UFR SSMT, Université de Cocody, 22 BP 582 Abidjan 22, Côte d'Ivoire, and <sup>c</sup>Laboratoire de Chimie Thérapeutique et Synthèse de Médicaments, UFR Sciences Pharmaceutiques, Université de Cocody, 01 BP V 34 Abidjan 01, Côte d'Ivoire

Correspondence e-mail: bibilamayayabisseyou@yahoo.fr


Received 15 May 2009; accepted 18 June 2009

Key indicators: single-crystal X-ray study;  $T = 295\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$ ;  $R$  factor = 0.047;  $wR$  factor = 0.097; data-to-parameter ratio = 12.0.

The title compound,  $\text{C}_{13}\text{H}_{14}\text{N}_2\text{OS}_2\cdot\text{H}_2\text{O}$ , appears in the form of bimolecular aggregate in which molecular components are linked by  $\text{O}-\text{H}\cdots\text{N}$  hydrogen bonding. The nine-membered imidazo[1,2-a]pyridine system is almost planar, with a mean deviation of  $0.026(1)\text{ \AA}$ . An intramolecular  $\text{C}-\text{H}\cdots\text{O}$  hydrogen bond forms within the imidazo[1,2-a]pyridine system. The crystal packing is consolidated by  $\text{O}-\text{H}\cdots\text{O}$  and  $\text{C}-\text{H}\cdots\text{O}$  hydrogen bonds, forming a supramolecular structure consisting of perpendicular infinite molecular chains running along the  $a$  and  $c$  axes.

### Related literature

For related structures, see: Bibila Mayaya Bisseyou *et al.* (2007, 2009); Duan *et al.* (2006). For hydrogen-bond motifs, see: Bernstein *et al.* (1995).



### Experimental

#### Crystal data

$\text{C}_{13}\text{H}_{14}\text{N}_2\text{OS}_2\cdot\text{H}_2\text{O}$   
 $M_r = 296.41$   
Orthorhombic,  $Pbca$   
 $a = 5.1405(1)\text{ \AA}$   
 $b = 17.7653(3)\text{ \AA}$   
 $c = 31.3919(6)\text{ \AA}$

$V = 2866.79(9)\text{ \AA}^3$   
 $Z = 8$   
Mo  $K\alpha$  radiation  
 $\mu = 0.37\text{ mm}^{-1}$   
 $T = 295\text{ K}$   
 $0.25 \times 0.15 \times 0.15\text{ mm}$

#### Data collection

Nonius KappaCCD diffractometer  
Absorption correction: multi-scan  
*DENZO/SCALEPACK*  
(Otwinowski & Minor, 1997)  
 $T_{\min} = 0.90$ ,  $T_{\max} = 0.95$

11253 measured reflections  
4202 independent reflections  
2344 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.05$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.047$   
 $wR(F^2) = 0.097$   
 $S = 0.99$   
4202 reflections

172 parameters  
H-atom parameters constrained  
 $\Delta\rho_{\max} = 0.18\text{ e \AA}^{-3}$   
 $\Delta\rho_{\min} = -0.25\text{ e \AA}^{-3}$

**Table 1**  
Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C4—H4 $\cdots$ O1	0.95	2.23	2.840 (4)	121
C3—H3 $\cdots$ O1 <sup>i</sup>	0.93	2.45	3.242 (4)	143
O2w—H22w $\cdots$ O2w <sup>ii</sup>	0.82	2.05	2.862 (4)	168
O2w—H21w $\cdots$ N2	0.82	2.06	2.849 (4)	164

Symmetry codes: (i)  $x + \frac{1}{2}, -y + \frac{1}{2}, -z$ ; (ii)  $x + \frac{1}{2}, y, -z + \frac{1}{2}$ .

Data collection: *COLLECT* (Nonius, 1997); cell refinement: *DENZO/SCALEPACK* (Otwinowski & Minor, 1997); data reduction: *DENZO/SCALEPACK*; program(s) used to solve structure: *SIR92* (Altomare *et al.*, 1994); program(s) used to refine structure: *CRYSTALS* (Betteridge *et al.*, 2003); molecular graphics: *ORTEP-3* (Farrugia, 1997) and *PLATON* (Spek, 2009); software used to prepare material for publication: *CRYSTALS*.

The authors thank the Spectropôle Service of the Faculty of Sciences and Techniques of Saint Jérôme (France) for the use of their diffractometer.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BQ2145).

### References

- Altomare, A., Casciaro, G., Giacovazzo, C., Guagliardi, A., Burla, M. C., Polidori, G. & Camalli, M. (1994). *J. Appl. Cryst.* **27**, 435.  
Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.  
Betteridge, P. W., Carruthers, J. R., Cooper, R. I., Prout, K. & Watkin, D. J. (2003). *J. Appl. Cryst.* **36**, 1487.  
Bibila Mayaya Bisseyou, Y., Adohi-Krou, A., Ouattara, M. P., Sissouma, D., Zoro-Diamé, E. G. & Ebby, N. (2009). *Z. Kristallogr. New Cryst. Struct.* **224**, 81–82.  
Bibila Mayaya Bisseyou, Y., Soro, A. P., Sissouma, D., Giorgi, M. & Ebby, N. (2007). *Acta Cryst.* **E63**, o4758–o4759.  
Duan, G.-Y., Tu, C.-B., Sun, Y.-W., Zhang, D.-T. & Wang, J.-W. (2006). *Acta Cryst.* **E62**, o1141–o1142.

Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.  
Nonius (1997). COLLECT. Nonius BV, Delft, The Netherlands.

Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276,  
*Macromolecular Crystallography*, Part A, edited by C. W. Carter Jr & R. M.  
Sweet, pp. 307–326. New York: Academic Press.  
Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.

## **supplementary materials**

*Acta Cryst.* (2009). E65, o1698–o1699 [doi:10.1107/S1600536809023514]

## 1-(2-Methylimidazo[1,2-*a*]pyridin-3-yl)-3,3-bis(methylsulfanyl)prop-2-enone monohydrate

**Y. Bibila Mayaya Bisseyou, D. Sissouma, S. D. Goulizan Bi, M. Ouattara and R. C. A. Yao-Kakou**

### Comment

As part of continuing work on heterocyclic compounds biologically active, we have synthesized new derivative in order to explore new potential activities or to improve known properties of this compound class. Asymmetric unit of title compound which crystallize as the monohydrate is consist of two independent molecular components connected between them by an O—H···N hydrogen bond forming thus one bimolecular aggregate (Fig. 1). In the imidazo[1,2-*a*]pyridine component, the values of bond lengths and angles of nine-membered imidazopyridine ring are similar to those found in previous studies (Bibila Mayaya Bisseyou *et al.*, 2007, 2009; Duan *et al.*, 2006). Imidazopyridine ring, essentially planar with mean deviation of 0.026 (1) Å, makes dihedral angles of 6.11 (23)° and 9.79 (15)° with P1(C9/C10/S1/C11) and P2(C9/C10/S2/C12) mean planes, respectively. Besides, we also note within of the same component, the presence of C—H···O intra-molecular hydrogen bond (Table 1) generating an S(6) motif (Bernstein *et al.*, 1995). The three-dimensional crystal packing is consolidated by inter-bimolecular aggregate hydrogen bonds (Fig. 2). Indeed, each bimolecular unit is linked to three others *via* O—H···O and C—H···O hydrogen bonds (Table 1) forming thus a supra-molecular structure consisted of to two perpendicular infinite chain types: the first one is established by bimolecular aggregates along *c* axis; the second one is parallel to *a* axis and constituted only by water molecules in zigzag fashion.

### Experimental

1-(2-methylimidazo[1,2-*a*]pyridine-3-yl) acetyl (1 g, 6.2 mmol) was dissolved in distilled dimethyl sulfoxide (15 ml), and the carbon disulfur (0.41 ml, 6.8 mmol) was added. After cooling of the mixture at 0°C, sodium hydrure (0.36 g, 15.4 mmol) was added. After stirring for 30 minutes at 0°C, the mixture was stirred at ambient temperature during 4 h. Solution was then cooled at 0°C and methyl iodide (2.5 molar equivalents) was added dropwise. The resulting mixture was then left under stirring during 24 h then poured into 50 ml ice-cold water. The precipitate was filtered and recrystallized from mixture of water-dioxane (2:1) to obtain orange single crystals of title compound (1.31 g; yield, 80%; m.p.: 430 K)

### Refinement

The H atoms were all located in a difference of Fourier map. They were all initially refined with soft restraints on the bond lengths and angles to regularize their geometry (C—H in the range 0.95–0.97 Å, O—H=0.82 Å and  $U_{\text{iso}}(\text{H})$  in the range 1.2–1.5 times  $U_{\text{eq}}$  of the parent atom), after which their positions were refined with riding constraints.

# supplementary materials

---

## Figures




Fig. 1. The molecular structure of the title compound and the atomic numbering scheme. Displacement ellipsoids are drawn at the 50% probability level. H atoms are shown as spheres of arbitrary radius. Dashed lines indicate hydrogen bonds.



Fig. 2. Crystal packing of the title compound, viewed down the  $a$  axis, showing the both perpendicular infinite chain types. The first one is established by bimolecular aggregates along  $c$  axis; the second one is parallel to  $a$  axis and constituted only by water molecules in zigzag fashion. H atoms not involved in hydrogen bonds have been omitted for clarity. Dashed lines indicate hydrogen bonds.

## 1-(2-Methylimidazo[1,2-a]pyridin-3-yl)-3,3-bis(methylsulfanyl)prop- 2-enone monohydrate

### Crystal data

$C_{13}H_{14}N_2OS_2H_2O$	$D_x = 1.373 \text{ Mg m}^{-3}$
$M_r = 296.41$	Melting point: 430 K
Orthorhombic, $Pbca$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2ac 2ab	Cell parameters from 11253 reflections
$a = 5.14050 (10) \text{ \AA}$	$\theta = 4\text{--}30^\circ$
$b = 17.7653 (3) \text{ \AA}$	$\mu = 0.37 \text{ mm}^{-1}$
$c = 31.3919 (6) \text{ \AA}$	$T = 295 \text{ K}$
$V = 2866.79 (9) \text{ \AA}^3$	Prism, orange
$Z = 8$	$0.25 \times 0.15 \times 0.15 \text{ mm}$
$F_{000} = 1248$	

### Data collection

Nonius KappaCCD diffractometer	2344 reflections with $I > 3\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.05$
$T = 295 \text{ K}$	$\theta_{\max} = 30.5^\circ$
$\varphi$ scans	$\theta_{\min} = 1.3^\circ$
Absorption correction: multi-scan DENZO/SCALEPACK (Otwinowski & Minor, 1997)	$h = 0 \rightarrow 7$
$T_{\min} = 0.90, T_{\max} = 0.95$	$k = 0 \rightarrow 25$
11253 measured reflections	$l = 0 \rightarrow 43$
4202 independent reflections	

### Refinement

Refinement on $F^2$	Hydrogen site location: inferred from neighbouring sites
Least-squares matrix: full	H-atom parameters constrained
$R[F^2 > 2\sigma(F^2)] = 0.047$	$w = 1/[\sigma^2(F^2) + (0.04P)^2 + 3.2P]$ ,

where  $P = [\max(F_0^2, 0) + 2F_c^2]/3$

$wR(F^2) = 0.097$        $(\Delta/\sigma)_{\max} = 0.0002$

$S = 0.99$        $\Delta\rho_{\max} = 0.18 \text{ e \AA}^{-3}$

2066 reflections       $\Delta\rho_{\min} = -0.25 \text{ e \AA}^{-3}$

172 parameters      Extinction correction: None

Primary atom site location: structure-invariant direct methods

### *Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.01264 (14)	0.40723 (4)	0.05322 (3)	0.0552
S2	-0.11568 (16)	0.46016 (4)	0.14025 (3)	0.0696
O1	0.3999 (4)	0.30577 (10)	0.05889 (6)	0.0542
N1	0.7820 (4)	0.21236 (11)	0.09663 (7)	0.0394
O2w	0.9388 (4)	0.07731 (12)	0.22997 (7)	0.0739
N2	0.8459 (5)	0.18741 (13)	0.16574 (8)	0.0559
C4	0.8270 (5)	0.20481 (13)	0.05354 (9)	0.0464
C7	0.9265 (5)	0.17257 (14)	0.12603 (9)	0.0474
C9	0.2551 (5)	0.35366 (14)	0.12396 (9)	0.0472
C5	0.6013 (5)	0.25573 (13)	0.11962 (8)	0.0410
C3	1.0207 (6)	0.15826 (14)	0.04040 (10)	0.0525
C2	1.1724 (6)	0.11813 (15)	0.06988 (11)	0.0573
C1	1.1255 (6)	0.12480 (15)	0.11196 (11)	0.0561
C6	0.6478 (5)	0.23742 (16)	0.16209 (9)	0.0494
C10	0.0715 (5)	0.40096 (14)	0.10800 (9)	0.0473
C8	0.4160 (5)	0.30513 (13)	0.09821 (9)	0.0417
C11	-0.2533 (6)	0.47334 (16)	0.04946 (12)	0.0704
C13	0.5083 (7)	0.2619 (2)	0.20158 (10)	0.0723
C12	0.0018 (9)	0.4409 (2)	0.19285 (13)	0.0992
H2	1.3072	0.0877	0.0601	0.0687*
H4	0.7197	0.2326	0.0346	0.0557*
H9	0.2801	0.3530	0.1530	0.0562*
H1	1.2232	0.0990	0.1323	0.0665*
H113	-0.2078	0.5201	0.0623	0.1052*
H112	-0.2875	0.4811	0.0199	0.1058*
H111	-0.4050	0.4535	0.0630	0.1059*
H3	1.0530	0.1538	0.0113	0.0637*
H132	0.5316	0.3152	0.2061	0.1075*
H133	0.3256	0.2531	0.1989	0.1079*
H131	0.5739	0.2357	0.2257	0.1077*
H22w	1.0875	0.0824	0.2392	0.1108*
H123	-0.0206	0.3876	0.1991	0.1488*
H21w	0.9353	0.1136	0.2140	0.1109*
H122	0.1833	0.4544	0.1939	0.1486*
H121	-0.0960	0.4706	0.2124	0.1492*

## supplementary materials

---

### *Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
S1	0.0444 (4)	0.0482 (4)	0.0731 (5)	0.0067 (3)	-0.0097 (4)	0.0037 (4)
S2	0.0549 (5)	0.0552 (4)	0.0988 (7)	0.0090 (4)	0.0082 (4)	-0.0195 (4)
O1	0.0542 (11)	0.0608 (12)	0.0477 (13)	0.0151 (9)	-0.0031 (10)	0.0060 (9)
N1	0.0375 (11)	0.0360 (10)	0.0448 (12)	-0.0011 (9)	-0.0041 (10)	0.0044 (9)
O2w	0.0700 (15)	0.0794 (15)	0.0723 (16)	-0.0039 (12)	-0.0076 (12)	0.0241 (12)
N2	0.0551 (15)	0.0623 (15)	0.0501 (15)	0.0054 (12)	-0.0097 (12)	0.0093 (12)
C4	0.0497 (16)	0.0397 (13)	0.0496 (17)	0.0002 (11)	0.0008 (12)	0.0037 (12)
C7	0.0416 (15)	0.0434 (13)	0.0572 (18)	0.0000 (11)	-0.0097 (13)	0.0060 (13)
C9	0.0414 (14)	0.0488 (14)	0.0515 (16)	0.0037 (12)	0.0006 (13)	-0.0051 (12)
C5	0.0362 (13)	0.0405 (13)	0.0464 (16)	-0.0010 (11)	-0.0001 (12)	0.0028 (11)
C3	0.0550 (17)	0.0431 (14)	0.0594 (18)	0.0028 (13)	0.0065 (14)	-0.0017 (12)
C2	0.0493 (17)	0.0444 (15)	0.078 (2)	0.0084 (13)	0.0018 (15)	-0.0043 (15)
C1	0.0497 (17)	0.0448 (14)	0.074 (2)	0.0076 (13)	-0.0120 (15)	0.0049 (14)
C6	0.0454 (15)	0.0552 (16)	0.0476 (17)	0.0005 (13)	-0.0047 (13)	0.0023 (13)
C10	0.0356 (14)	0.0381 (13)	0.068 (2)	-0.0033 (11)	0.0012 (12)	-0.0048 (12)
C8	0.0356 (13)	0.0400 (12)	0.0495 (17)	-0.0016 (10)	-0.0039 (12)	0.0026 (12)
C11	0.0485 (17)	0.0517 (17)	0.111 (3)	0.0064 (14)	-0.0165 (19)	0.0092 (18)
C13	0.076 (2)	0.098 (2)	0.0430 (18)	0.015 (2)	-0.0014 (17)	0.0030 (17)
C12	0.112 (3)	0.105 (3)	0.081 (3)	0.021 (3)	0.016 (2)	-0.035 (2)

### *Geometric parameters ( $\text{\AA}$ , $^\circ$ )*

S1—C10	1.750 (3)	C5—C6	1.393 (4)
S1—C11	1.806 (3)	C5—C8	1.459 (3)
S2—C10	1.748 (3)	C3—C2	1.405 (4)
S2—C12	1.791 (4)	C3—H3	0.930
O1—C8	1.237 (3)	C2—C1	1.348 (4)
N1—C4	1.379 (3)	C2—H2	0.931
N1—C7	1.380 (3)	C1—H1	0.932
N1—C5	1.406 (3)	C6—C13	1.496 (4)
O2w—H22w	0.822	C11—H113	0.952
O2w—H21w	0.817	C11—H112	0.954
N2—C7	1.340 (4)	C11—H111	0.955
N2—C6	1.356 (3)	C13—H132	0.966
C4—C3	1.358 (4)	C13—H133	0.956
C4—H4	0.948	C13—H131	0.951
C7—C1	1.401 (4)	C12—H123	0.974
C9—C10	1.359 (3)	C12—H122	0.964
C9—C8	1.443 (4)	C12—H121	0.953
C9—H9	0.922		
C10—S1—C11	103.68 (15)	C5—C6—N2	111.3 (2)
C10—S2—C12	103.50 (16)	C5—C6—C13	130.0 (3)
C4—N1—C7	121.1 (2)	N2—C6—C13	118.7 (3)
C4—N1—C5	131.9 (2)	S1—C10—S2	115.83 (15)

C7—N1—C5	107.0 (2)	S1—C10—C9	121.4 (2)
H22w—O2w—H21w	98.6	S2—C10—C9	122.7 (2)
C7—N2—C6	106.4 (2)	C5—C8—C9	118.4 (2)
N1—C4—C3	118.7 (3)	C5—C8—O1	120.6 (2)
N1—C4—H4	117.7	C9—C8—O1	121.0 (2)
C3—C4—H4	123.6	S1—C11—H113	110.7
N1—C7—N2	110.8 (2)	S1—C11—H112	107.3
N1—C7—C1	119.5 (3)	H113—C11—H112	109.3
N2—C7—C1	129.7 (3)	S1—C11—H111	110.4
C10—C9—C8	124.1 (3)	H113—C11—H111	109.6
C10—C9—H9	118.0	H112—C11—H111	109.5
C8—C9—H9	117.8	C6—C13—H132	110.3
N1—C5—C6	104.5 (2)	C6—C13—H133	110.5
N1—C5—C8	121.6 (2)	H132—C13—H133	107.1
C6—C5—C8	133.9 (2)	C6—C13—H131	110.4
C4—C3—C2	121.1 (3)	H132—C13—H131	108.6
C4—C3—H3	118.7	H133—C13—H131	109.9
C2—C3—H3	120.3	S2—C12—H123	109.3
C3—C2—C1	120.1 (3)	S2—C12—H122	108.2
C3—C2—H2	119.3	H123—C12—H122	110.5
C1—C2—H2	120.5	S2—C12—H121	108.0
C7—C1—C2	119.5 (3)	H123—C12—H121	110.3
C7—C1—H1	118.4	H122—C12—H121	110.5
C2—C1—H1	122.0		

*Hydrogen-bond geometry (Å, °)*

D—H···A	D—H	H···A	D···A	D—H···A
C4—H4···O1	0.95	2.23	2.840 (4)	121
C3—H3···O1 <sup>i</sup>	0.93	2.45	3.242 (4)	143
O2w—H22w···O2w <sup>ii</sup>	0.82	2.05	2.862 (4)	168
O2w—H21w···N2	0.82	2.06	2.849 (4)	164

Symmetry codes: (i)  $x+1/2, -y+1/2, -z$ ; (ii)  $x+1/2, y, -z+1/2$ .

## supplementary materials

---

Fig. 1




Fig. 2

