

4-Methyl-3-[4-(3-pyridyl)pyrimidin-2-yl-oxy]aniline

Shi-Gui Tang,^a Cheng-Long Yang,^b Jun-Hua Chen,^c Sheng Bi^d and Cheng Guo^{c*}

^aCollege of Biology and Pharmacy Engineering, Nanjing University of Technology, Xinmofan Road No. 5 Nanjing, Nanjing 210009, People's Republic of China,
^bCollege of Kang ni, Nanjing College of Engineering, Hongjing Road No. 1 Jiangning District, Nanjing, Nanjing 211167, People's Republic of China, ^cCollege of Science, Nanjing University of Technology, Xinmofan Road No. 5 Nanjing, Nanjing 210009, People's Republic of China, and ^dGuangdong Petroleum College of Technology, Chisha Road No. 23 Nanhui District, Fushan, Fushan 528222, People's Republic of China

Correspondence e-mail: guocheng@njut.edu.cn


Received 3 June 2009; accepted 21 June 2009

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.005\text{ \AA}$; R factor = 0.061; wR factor = 0.170; data-to-parameter ratio = 13.3.

In the title compound, $\text{C}_{16}\text{H}_{14}\text{N}_4\text{O}$, there are intermolecular $\text{N}-\text{H}\cdots\text{N}$ hydrogen bonds which may be effective in stabilizing the crystal. The title compound is an important medicament and is used in the synthesis of antitumour drugs.

Related literature

For bond-length data, see: Allen *et al.* (1987)



Experimental

Crystal data

$\text{C}_{16}\text{H}_{14}\text{N}_4\text{O}$
 $M_r = 278.31$

Monoclinic, $P2_1/c$
 $a = 8.5800(17)\text{ \AA}$

$b = 20.360(4)\text{ \AA}$
 $c = 8.0780(16)\text{ \AA}$
 $\beta = 98.29(3)^\circ$
 $V = 1396.4(5)\text{ \AA}^3$
 $Z = 4$

Mo $K\alpha$ radiation
 $\mu = 0.09\text{ mm}^{-1}$
 $T = 293\text{ K}$
 $0.20 \times 0.20 \times 0.10\text{ mm}$

Data collection

Enraf–Nonius CAD-4 diffractometer
Absorption correction: ψ scan (North *et al.*, 1968)
 $T_{\min} = 0.983$, $T_{\max} = 0.991$
2698 measured reflections

2526 independent reflections
1587 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.018$
3 standard reflections
every 200 reflections
intensity decay: 1%

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.061$
 $wR(F^2) = 0.170$
 $S = 1.01$
2526 reflections

190 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.28\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.30\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N1—H1A \cdots N4 ⁱ	0.86	2.47	3.214 (4)	145
N1—H1B \cdots N2 ⁱⁱ	0.86	2.43	3.166 (4)	144

Symmetry codes: (i) $-x, y + \frac{1}{2}, -z + \frac{3}{2}$; (ii) $-x, -y + 1, -z + 2$.

Data collection: *CAD-4 EXPRESS* (Enraf–Nonius, 1989); cell refinement: *CAD-4 EXPRESS*; data reduction: *XCAD4* (Harms & Wocadlo, 1995); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *SHELXL97*.

The authors thank the Center of Testing and Analysis, Nanjing University, for support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: PV2165).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Enraf–Nonius (1989). *CAD-4 EXPRESS*. Enraf–Nonius, Delft, The Netherlands.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Harms, K. & Wocadlo, S. (1995). *XCAD4*. University of Marburg, Germany.
- North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst. A* **24**, 351–359.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.

supplementary materials

Acta Cryst. (2009). E65, o1707 [doi:10.1107/S1600536809023770]

4-Methyl-3-[4-(3-pyridyl)pyrimidin-2-yloxy]aniline

S.-G. Tang, C.-L. Yang, J.-H. Chen, S. Bi and C. Guo

Comment

Some derivatives of benzenamine are important medical materials. We report here the crystal structure of the title compound, (I), which was synthesized by the reaction of *tert*-butyl-4-methyl-3-(4-(3-pyridinyl)pyrimidin-2-yloxy)phenylcarbamate and dichloromethane with trifluoroacetic acid. The molecular structure of (I) is shown in Fig. 1. The bond lengths and angles in (I) are within normal ranges (Allen *et al.* 1987). The structure is stabilized by N—H···N type hydrogen bonds (Table 1 and Fig. 2).

Experimental

In a three neck bottom flask containing dichloromethane (65 ml) and trifluoroacetic acid (20 ml) was added *tert*-butyl-4-methyl-3-(4-(3-pyridinyl)pyrimidin-2-yloxy)phenylcarbamate (7.5 g) at 273 K. After the addition of all chemicals, the flask was taken off the ice-water bath and the reaction was allowed to take place for 6 h at room temperature. Neutralized with sodium bicarbonate and separated the dichloromethane and aqueous layers. On evaporation of dichloromethane a solid product was obtained. Crystals of (I) suitable for X-ray diffraction were obtained by slow evaporation of a cyclohexane solution.

Refinement

All H atoms bonded to the C atoms were placed geometrically at the distances of 0.93, 0.96 and 0.86 Å, for aryl, methyl and amino H-atoms, respectively, and were included in the refinement in riding motion approximation with $U_{\text{iso}}(\text{H}) = 1.2$ or $1.5U_{\text{eq}}$ of the carrier atom.

Figures




Fig. 1. The molecular structure of (I), showing the atom-numbering scheme and displacement ellipsoids at the 50% probability level.




Fig. 2. A packing diagram of (I). The intermolecular hydrogen bonds are shown as dashed lines.

supplementary materials

4-Methyl-3-[4-(3-pyridyl)pyrimidin-2-yloxy]aniline

Crystal data

C ₁₆ H ₁₄ N ₄ O	$F_{000} = 584$
$M_r = 278.31$	$D_x = 1.324 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2ybc	Cell parameters from 25 reflections
$a = 8.5800 (17) \text{ \AA}$	$\theta = 9-13^\circ$
$b = 20.360 (4) \text{ \AA}$	$\mu = 0.09 \text{ mm}^{-1}$
$c = 8.0780 (16) \text{ \AA}$	$T = 293 \text{ K}$
$\beta = 98.29 (3)^\circ$	Block, colorless
$V = 1396.4 (5) \text{ \AA}^3$	$0.20 \times 0.20 \times 0.10 \text{ mm}$
$Z = 4$	

Data collection

Enraf–Nonius CAD-4 diffractometer	$R_{\text{int}} = 0.018$
Radiation source: fine-focus sealed tube	$\theta_{\text{max}} = 25.3^\circ$
Monochromator: graphite	$\theta_{\text{min}} = 2.0^\circ$
$T = 293 \text{ K}$	$h = -10 \rightarrow 0$
$\omega/2\theta$ scans	$k = 0 \rightarrow 24$
Absorption correction: ψ scan (North <i>et al.</i> , 1968)	$l = -9 \rightarrow 9$
$T_{\text{min}} = 0.983$, $T_{\text{max}} = 0.991$	3 standard reflections
2698 measured reflections	every 200 reflections
2526 independent reflections	intensity decay: 1%
1587 reflections with $I > 2\sigma(I)$	

Refinement

Refinement on F^2	Hydrogen site location: inferred from neighbouring sites
Least-squares matrix: full	H-atom parameters constrained
$R[F^2 > 2\sigma(F^2)] = 0.061$	$w = 1/[\sigma^2(F_o^2) + (0.06P)^2 + 1.4P]$
$wR(F^2) = 0.170$	where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.01$	$(\Delta/\sigma)_{\text{max}} < 0.001$
2526 reflections	$\Delta\rho_{\text{max}} = 0.28 \text{ e \AA}^{-3}$
190 parameters	$\Delta\rho_{\text{min}} = -0.30 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: SHELXL97 (Sheldrick, 2008)
Secondary atom site location: difference Fourier map	Extinction coefficient: 0.017 (4)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O	0.1194 (2)	0.51325 (11)	0.7760 (3)	0.0508 (6)
N1	-0.2002 (3)	0.66654 (13)	1.0241 (4)	0.0543 (8)
H1A	-0.2537	0.7024	1.0186	0.065*
H1B	-0.1955	0.6419	1.1111	0.065*
C1	0.1120 (5)	0.5926 (2)	0.4732 (5)	0.0680 (11)
H1C	0.1658	0.5515	0.4963	0.102*
H1D	0.1862	0.6258	0.4525	0.102*
H1E	0.0335	0.5879	0.3765	0.102*
N2	0.3246 (3)	0.44734 (12)	0.7623 (3)	0.0416 (7)
C2	0.0341 (4)	0.61219 (17)	0.6213 (4)	0.0460 (8)
N3	0.3632 (3)	0.56324 (13)	0.7919 (4)	0.0497 (7)
C3	-0.0526 (4)	0.66983 (17)	0.6214 (4)	0.0498 (9)
H3B	-0.0593	0.6973	0.5285	0.060*
C4	-0.1290 (4)	0.68813 (16)	0.7523 (4)	0.0480 (9)
H4A	-0.1857	0.7272	0.7468	0.058*
N4	0.4792 (4)	0.26228 (15)	0.6373 (4)	0.0649 (9)
C5	-0.1216 (3)	0.64847 (14)	0.8928 (4)	0.0405 (8)
C6	-0.0318 (3)	0.59152 (14)	0.8984 (4)	0.0395 (8)
H6A	-0.0230	0.5645	0.9921	0.047*
C7	0.0441 (3)	0.57516 (15)	0.7653 (4)	0.0403 (8)
C8	0.2772 (4)	0.50942 (15)	0.7753 (4)	0.0412 (8)
C9	0.4799 (4)	0.43896 (15)	0.7646 (4)	0.0406 (8)
C10	0.5823 (4)	0.49200 (17)	0.7776 (5)	0.0522 (9)
H10A	0.6898	0.4863	0.7773	0.063*
C11	0.5179 (4)	0.55349 (17)	0.7909 (5)	0.0556 (10)
H11A	0.5845	0.5898	0.7995	0.067*
C12	0.5341 (4)	0.37050 (15)	0.7517 (4)	0.0411 (8)
C13	0.4408 (4)	0.32551 (17)	0.6553 (5)	0.0526 (9)
H13A	0.3447	0.3400	0.5990	0.063*
C14	0.6177 (5)	0.24293 (19)	0.7200 (5)	0.0630 (11)
H14A	0.6476	0.1993	0.7104	0.076*
C15	0.7182 (5)	0.28438 (19)	0.8184 (5)	0.0619 (10)
H15A	0.8136	0.2687	0.8737	0.074*

supplementary materials

C16	0.6774 (4)	0.34860 (18)	0.8347 (5)	0.0531 (9)
H16A	0.7445	0.3772	0.9004	0.064*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O	0.0398 (13)	0.0379 (12)	0.0783 (18)	-0.0001 (10)	0.0208 (11)	-0.0024 (11)
N1	0.066 (2)	0.0395 (16)	0.0609 (19)	0.0056 (14)	0.0227 (16)	-0.0018 (14)
C1	0.069 (3)	0.083 (3)	0.055 (2)	-0.006 (2)	0.019 (2)	0.000 (2)
N2	0.0410 (16)	0.0358 (14)	0.0497 (17)	-0.0021 (12)	0.0123 (12)	-0.0049 (12)
C2	0.0391 (18)	0.053 (2)	0.046 (2)	-0.0059 (16)	0.0087 (15)	0.0002 (16)
N3	0.0463 (17)	0.0412 (16)	0.0619 (19)	-0.0045 (13)	0.0084 (14)	-0.0053 (14)
C3	0.0455 (19)	0.052 (2)	0.051 (2)	-0.0074 (17)	0.0033 (16)	0.0154 (17)
C4	0.0435 (19)	0.0358 (18)	0.065 (2)	0.0023 (15)	0.0084 (17)	0.0086 (16)
N4	0.062 (2)	0.0462 (18)	0.090 (3)	-0.0014 (16)	0.0242 (18)	-0.0145 (17)
C5	0.0369 (17)	0.0353 (17)	0.050 (2)	-0.0061 (14)	0.0070 (14)	-0.0037 (15)
C6	0.0381 (17)	0.0328 (17)	0.048 (2)	-0.0047 (14)	0.0077 (15)	0.0052 (14)
C7	0.0340 (17)	0.0313 (16)	0.057 (2)	-0.0029 (13)	0.0106 (15)	-0.0013 (15)
C8	0.0406 (18)	0.0393 (18)	0.0453 (19)	-0.0014 (15)	0.0116 (14)	-0.0034 (15)
C9	0.0393 (18)	0.0424 (18)	0.0414 (18)	-0.0007 (14)	0.0101 (14)	-0.0043 (14)
C10	0.0387 (18)	0.048 (2)	0.071 (3)	-0.0050 (16)	0.0125 (17)	-0.0057 (18)
C11	0.045 (2)	0.045 (2)	0.076 (3)	-0.0100 (16)	0.0074 (18)	-0.0075 (18)
C12	0.0387 (18)	0.0407 (18)	0.0462 (19)	-0.0019 (14)	0.0143 (15)	-0.0032 (15)
C13	0.0426 (19)	0.047 (2)	0.070 (2)	-0.0003 (16)	0.0138 (17)	-0.0118 (18)
C14	0.070 (3)	0.044 (2)	0.081 (3)	0.010 (2)	0.033 (2)	0.000 (2)
C15	0.058 (2)	0.057 (2)	0.072 (3)	0.018 (2)	0.012 (2)	0.006 (2)
C16	0.050 (2)	0.054 (2)	0.056 (2)	0.0034 (17)	0.0087 (17)	-0.0060 (17)

Geometric parameters (\AA , $^\circ$)

O—C8	1.357 (4)	N4—C14	1.336 (5)
O—C7	1.413 (4)	N4—C13	1.342 (4)
N1—C5	1.387 (4)	C5—C6	1.389 (4)
N1—H1A	0.8600	C6—C7	1.376 (4)
N1—H1B	0.8600	C6—H6A	0.9300
C1—C2	1.506 (5)	C9—C10	1.387 (4)
C1—H1C	0.9600	C9—C12	1.478 (4)
C1—H1D	0.9600	C10—C11	1.379 (5)
C1—H1E	0.9600	C10—H10A	0.9300
N2—C8	1.337 (4)	C11—H11A	0.9300
N2—C9	1.340 (4)	C12—C13	1.381 (5)
C2—C7	1.379 (5)	C12—C16	1.386 (5)
C2—C3	1.389 (5)	C13—H13A	0.9300
N3—C8	1.317 (4)	C14—C15	1.375 (5)
N3—C11	1.344 (4)	C14—H14A	0.9300
C3—C4	1.374 (5)	C15—C16	1.365 (5)
C3—H3B	0.9300	C15—H15A	0.9300
C4—C5	1.387 (4)	C16—H16A	0.9300
C4—H4A	0.9300		

C8—O—C7	119.9 (2)	C6—C7—O	115.6 (3)
C5—N1—H1A	120.0	C2—C7—O	120.8 (3)
C5—N1—H1B	120.0	N3—C8—N2	128.5 (3)
H1A—N1—H1B	120.0	N3—C8—O	119.8 (3)
C2—C1—H1C	109.5	N2—C8—O	111.7 (3)
C2—C1—H1D	109.5	N2—C9—C10	121.3 (3)
H1C—C1—H1D	109.5	N2—C9—C12	116.2 (3)
C2—C1—H1E	109.5	C10—C9—C12	122.5 (3)
H1C—C1—H1E	109.5	C11—C10—C9	117.1 (3)
H1D—C1—H1E	109.5	C11—C10—H10A	121.4
C8—N2—C9	115.6 (3)	C9—C10—H10A	121.4
C7—C2—C3	115.5 (3)	N3—C11—C10	122.8 (3)
C7—C2—C1	123.0 (3)	N3—C11—H11A	118.6
C3—C2—C1	121.5 (3)	C10—C11—H11A	118.6
C8—N3—C11	114.6 (3)	C13—C12—C16	117.5 (3)
C4—C3—C2	123.0 (3)	C13—C12—C9	120.1 (3)
C4—C3—H3B	118.5	C16—C12—C9	122.4 (3)
C2—C3—H3B	118.5	N4—C13—C12	124.5 (3)
C3—C4—C5	120.1 (3)	N4—C13—H13A	117.8
C3—C4—H4A	120.0	C12—C13—H13A	117.8
C5—C4—H4A	120.0	N4—C14—C15	123.1 (3)
C14—N4—C13	116.3 (3)	N4—C14—H14A	118.4
C4—C5—N1	120.1 (3)	C15—C14—H14A	118.4
C4—C5—C6	118.2 (3)	C16—C15—C14	119.7 (4)
N1—C5—C6	121.7 (3)	C16—C15—H15A	120.2
C7—C6—C5	119.9 (3)	C14—C15—H15A	120.2
C7—C6—H6A	120.0	C15—C16—C12	118.9 (4)
C5—C6—H6A	120.0	C15—C16—H16A	120.5
C6—C7—C2	123.2 (3)	C12—C16—H16A	120.5
C7—C2—C3—C4	−2.5 (5)	C7—O—C8—N2	−171.2 (3)
C1—C2—C3—C4	178.0 (3)	C8—N2—C9—C10	−0.8 (5)
C2—C3—C4—C5	0.0 (5)	C8—N2—C9—C12	179.5 (3)
C3—C4—C5—N1	−179.2 (3)	N2—C9—C10—C11	1.1 (5)
C3—C4—C5—C6	1.9 (5)	C12—C9—C10—C11	−179.2 (3)
C4—C5—C6—C7	−1.3 (4)	C8—N3—C11—C10	−1.5 (5)
N1—C5—C6—C7	179.8 (3)	C9—C10—C11—N3	0.1 (6)
C5—C6—C7—C2	−1.2 (5)	N2—C9—C12—C13	34.5 (4)
C5—C6—C7—O	−174.0 (3)	C10—C9—C12—C13	−145.2 (3)
C3—C2—C7—C6	3.1 (5)	N2—C9—C12—C16	−145.1 (3)
C1—C2—C7—C6	−177.4 (3)	C10—C9—C12—C16	35.2 (5)
C3—C2—C7—O	175.4 (3)	C14—N4—C13—C12	0.1 (5)
C1—C2—C7—O	−5.0 (5)	C16—C12—C13—N4	0.2 (5)
C8—O—C7—C6	−119.7 (3)	C9—C12—C13—N4	−179.3 (3)
C8—O—C7—C2	67.4 (4)	C13—N4—C14—C15	−0.3 (6)
C11—N3—C8—N2	1.9 (5)	N4—C14—C15—C16	0.0 (6)
C11—N3—C8—O	179.5 (3)	C14—C15—C16—C12	0.4 (6)
C9—N2—C8—N3	−0.8 (5)	C13—C12—C16—C15	−0.5 (5)
C9—N2—C8—O	−178.5 (3)	C9—C12—C16—C15	179.1 (3)

supplementary materials

C7—O—C8—N3


10.9 (5)

Hydrogen-bond geometry (\AA , $^\circ$)

$D\text{—H}\cdots A$	$D\text{—H}$	$\text{H}\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
N1—H1A…N4 ⁱ	0.86	2.47	3.214 (4)	145
N1—H1B…N2 ⁱⁱ	0.86	2.43	3.166 (4)	144

Symmetry codes: (i) $-x, y+1/2, -z+3/2$; (ii) $-x, -y+1, -z+2$.

Fig. 1



supplementary materials

Fig. 2

