

Chlorido(pentane-2,4-dionato- κ^2O,O')-(1,10-phenanthroline- κ^2N,N')copper(II)

 Yew Sent Wong,^a Chew Hee Ng^a and Seik Weng Ng^{b*}

^aFaculty of Engineering and Science, Universiti Tunku Abdul Rahman, 53300 Kuala Lumpur, Malaysia, and ^bDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia

Correspondence e-mail: seikweng@um.edu.my


Received 9 July 2009; accepted 10 July 2009

Key indicators: single-crystal X-ray study; $T = 163$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.028; wR factor = 0.073; data-to-parameter ratio = 16.2.

The Cu^{II} atom in the title compound, $[\text{Cu}(\text{C}_5\text{H}_7\text{O}_2)\text{Cl}(\text{C}_{12}\text{H}_8\text{N}_2)]$, shows a distorted square-planar coordination; the chelating N and O atoms occupy the basal sites and the Cl atom the apical site. The square-pyramidal character along the Berry $D_{3h}-C_{4v}$ pseudorotation pathway is 92%.

Related literature

For the synthesis and electronic spectrum, see: Kwik & Ang (1978). For isostructural $\text{CuBr}(\text{C}_{12}\text{H}_8\text{N}_2)(\text{C}_5\text{H}_7\text{O}_2)$, see: Onawumi *et al.* (2008).



Experimental

Crystal data

$[\text{Cu}(\text{C}_5\text{H}_7\text{O}_2)\text{Cl}(\text{C}_{12}\text{H}_8\text{N}_2)]$
 $M_r = 378.30$

Triclinic, $P\bar{1}$
 $a = 7.5436$ (1) Å

$b = 9.0347$ (2) Å
 $c = 11.9399$ (2) Å
 $\alpha = 85.638$ (1)°
 $\beta = 72.329$ (1)°
 $\gamma = 85.716$ (1)°
 $V = 771.97$ (2) Å³

$Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 1.60$ mm⁻¹
 $T = 163$ K
 $0.35 \times 0.35 \times 0.15$ mm

Data collection

Bruker SMART APEX
 diffractometer
 Absorption correction: multi-scan
 (SADABS; Sheldrick, 1996)
 $T_{\text{min}} = 0.605$, $T_{\text{max}} = 0.796$

5384 measured reflections
 3406 independent reflections
 3151 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.015$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.028$
 $wR(F^2) = 0.073$
 $S = 1.03$
 3406 reflections

210 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.34$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.43$ e Å⁻³

Table 1

Selected geometric parameters (Å, °).

Cu1—Cl1	2.4717 (6)	Cu1—N1	2.038 (2)
Cu1—O1	1.927 (2)	Cu1—N2	2.025 (2)
Cu1—O2	1.936 (2)		
O1—Cu1—O2	93.82 (7)	N1—Cu1—N2	81.04 (7)

Data collection: *APEX2* (Bruker, 2008); cell refinement: *APEX2*; data reduction: *SAINT* (Bruker, 2008); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *pubCIF* (Westrip, 2009).

The authors thank MOSTI (grant No. 02-02-11-SF0033) and the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: XU2552).

References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
 Bruker (2008). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Kwik, W. L. & Ang, H. G. (1978). *Aust. J. Chem.* **31**, 459–463.
 Onawumi, O. O. E., Faboya, O. O. P., Odunola, O. A., Prasad, T. K. & Rajasekharan, M. V. (2008). *Polyhedron*, **27**, 113–117.
 Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Westrip, S. P. (2009). *pubCIF*. In preparation.

supplementary materials

Acta Cryst. (2009). E65, m934 [doi:10.1107/S1600536809027111]

Chlorido(pentane-2,4-dionato- κ^2O,O')(1,10-phenanthroline- κ^2N,N')copper(II)

Y. S. Wong, C. H. Ng and S. W. Ng

Experimental

Copper chloride dihydrate (0.17 g, 1 mmol) and 1,10-phenanthroline monohydrate (0.20 g, 1 mmol) were reacted in a 1:1 v/v methanol-water mixture (5 ml) to give a light green precipitate. The copper dichloride phenanthroline adduct (0.047 g, 0.15 mmol) was dissolved in a methanol-water mixture (5 ml) and this was treated with excess acetylacetone (5 ml) and 0.1 M sodium hydroxide (5 ml). The dark green solution was heated at 343 K for 30 min. Bluish-green crystals separated from the cool solution.

Refinement

Carbon-bound H-atoms were placed in calculated positions (C—H 0.95–0.98 Å) and were included in the refinement in the riding model approximation, with $U_{iso}(H)$ set to 1.2–1.5 $U_{eq}(C)$.

Figures




Fig. 1. Thermal ellipsoid (Barbour, 2001) plot of $CuCl(C_{12}H_8N_2)(C_5H_7O_2)$ at the 70% probability level. Hydrogen atoms are drawn as spheres of arbitrary radii.

Chlorido(pentane-2,4-dionato- κ^2O,O')(1,10- phenanthroline- κ^2N,N')copper(II)

Crystal data

$[Cu(C_5H_7O_2)Cl(C_{12}H_8N_2)]$

$M_r = 378.30$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$a = 7.5436$ (1) Å

$b = 9.0347$ (2) Å

$c = 11.9399$ (2) Å

$\alpha = 85.638$ (1)°

$\beta = 72.329$ (1)°

$\gamma = 85.716$ (1)°

$V = 771.97$ (2) Å³

$Z = 2$

$F_{000} = 386$

$D_x = 1.627$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 4297 reflections

$\theta = 2.8$ – 28.3 °

$\mu = 1.60$ mm⁻¹

$T = 163$ K

Block, blue

$0.35 \times 0.35 \times 0.15$ mm

supplementary materials

Data collection

Bruker SMART APEX diffractometer	3406 independent reflections
Radiation source: fine-focus sealed tube	3151 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.015$
$T = 163$ K	$\theta_{\text{max}} = 27.5^\circ$
ω scans	$\theta_{\text{min}} = 1.8^\circ$
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	$h = -9 \rightarrow 9$
$T_{\text{min}} = 0.605$, $T_{\text{max}} = 0.796$	$k = -11 \rightarrow 11$
5384 measured reflections	$l = -15 \rightarrow 15$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.028$	H-atom parameters constrained
$wR(F^2) = 0.073$	$w = 1/[\sigma^2(F_o^2) + (0.0226P)^2 + 1.0753P]$
$S = 1.03$	where $P = (F_o^2 + 2F_c^2)/3$
3406 reflections	$(\Delta/\sigma)_{\text{max}} = 0.001$
210 parameters	$\Delta\rho_{\text{max}} = 0.34 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	$\Delta\rho_{\text{min}} = -0.43 \text{ e } \text{\AA}^{-3}$
	Extinction correction: none

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Cu1	0.74448 (4)	0.62243 (3)	0.25532 (2)	0.01986 (8)
Cl1	0.42757 (8)	0.74327 (6)	0.30310 (4)	0.02521 (12)
O1	0.7987 (2)	0.61537 (17)	0.40339 (13)	0.0275 (3)
O2	0.8805 (2)	0.79922 (17)	0.19426 (14)	0.0272 (3)
N1	0.7637 (2)	0.56785 (19)	0.08940 (15)	0.0197 (3)
N2	0.6667 (2)	0.41013 (19)	0.29186 (15)	0.0201 (3)
C1	0.8971 (4)	0.6884 (3)	0.5587 (2)	0.0391 (6)
H1A	0.7958	0.6294	0.6087	0.059*
H1B	0.8954	0.7830	0.5942	0.059*
H1C	1.0168	0.6335	0.5514	0.059*
C2	0.8716 (3)	0.7181 (3)	0.4385 (2)	0.0261 (5)
C3	0.9315 (3)	0.8491 (3)	0.3737 (2)	0.0313 (5)
H3	0.9691	0.9233	0.4126	0.038*
C4	0.9407 (3)	0.8800 (2)	0.2562 (2)	0.0256 (5)
C5	1.0312 (4)	1.0189 (3)	0.1944 (2)	0.0370 (6)
H5A	1.0362	1.0225	0.1113	0.056*
H5B	1.1580	1.0185	0.2005	0.056*

H5C	0.9584	1.1063	0.2314	0.056*
C6	0.8170 (3)	0.6495 (2)	-0.01113 (19)	0.0243 (4)
H6	0.8574	0.7464	-0.0101	0.029*
C7	0.8159 (3)	0.5982 (3)	-0.11824 (19)	0.0270 (5)
H7	0.8561	0.6597	-0.1884	0.032*
C8	0.7568 (3)	0.4593 (3)	-0.12245 (18)	0.0247 (4)
H8	0.7551	0.4239	-0.1950	0.030*
C9	0.6982 (3)	0.3695 (2)	-0.01631 (18)	0.0213 (4)
C10	0.6357 (3)	0.2224 (2)	-0.00981 (19)	0.0254 (4)
H10	0.6279	0.1814	-0.0790	0.030*
C11	0.5872 (3)	0.1408 (2)	0.0943 (2)	0.0263 (5)
H11	0.5476	0.0430	0.0966	0.032*
C12	0.5948 (3)	0.1995 (2)	0.20076 (18)	0.0216 (4)
C13	0.5500 (3)	0.1196 (2)	0.3117 (2)	0.0266 (5)
H13	0.5100	0.0210	0.3198	0.032*
C14	0.5653 (3)	0.1866 (2)	0.40724 (19)	0.0267 (5)
H14	0.5371	0.1341	0.4820	0.032*
C15	0.6226 (3)	0.3326 (2)	0.39506 (18)	0.0230 (4)
H15	0.6302	0.3778	0.4626	0.028*
C16	0.6529 (3)	0.3440 (2)	0.19648 (17)	0.0183 (4)
C17	0.7057 (3)	0.4300 (2)	0.08651 (17)	0.0183 (4)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cu1	0.02849 (15)	0.01578 (13)	0.01782 (13)	-0.00745 (10)	-0.00941 (10)	-0.00008 (9)
Cl1	0.0299 (3)	0.0241 (3)	0.0245 (2)	-0.0016 (2)	-0.0116 (2)	-0.00471 (19)
O1	0.0404 (9)	0.0230 (8)	0.0251 (8)	-0.0066 (7)	-0.0179 (7)	-0.0008 (6)
O2	0.0357 (9)	0.0206 (8)	0.0261 (8)	-0.0129 (7)	-0.0080 (7)	-0.0005 (6)
N1	0.0240 (9)	0.0180 (8)	0.0181 (8)	-0.0041 (7)	-0.0076 (7)	0.0001 (6)
N2	0.0254 (9)	0.0168 (8)	0.0189 (8)	-0.0031 (7)	-0.0075 (7)	-0.0007 (6)
C1	0.0551 (17)	0.0374 (14)	0.0368 (13)	0.0050 (12)	-0.0312 (13)	-0.0112 (11)
C2	0.0282 (11)	0.0258 (11)	0.0291 (11)	0.0042 (9)	-0.0149 (9)	-0.0102 (9)
C3	0.0356 (13)	0.0265 (12)	0.0385 (13)	-0.0064 (10)	-0.0179 (11)	-0.0102 (10)
C4	0.0211 (10)	0.0200 (10)	0.0355 (12)	-0.0043 (8)	-0.0060 (9)	-0.0071 (9)
C5	0.0362 (13)	0.0262 (12)	0.0447 (14)	-0.0134 (10)	-0.0020 (11)	-0.0083 (10)
C6	0.0290 (11)	0.0211 (10)	0.0238 (10)	-0.0082 (8)	-0.0092 (9)	0.0032 (8)
C7	0.0327 (12)	0.0289 (12)	0.0200 (10)	-0.0064 (9)	-0.0093 (9)	0.0051 (8)
C8	0.0282 (11)	0.0293 (12)	0.0183 (9)	-0.0032 (9)	-0.0088 (8)	-0.0022 (8)
C9	0.0209 (10)	0.0231 (10)	0.0210 (10)	-0.0020 (8)	-0.0073 (8)	-0.0018 (8)
C10	0.0308 (11)	0.0237 (11)	0.0246 (10)	-0.0040 (9)	-0.0105 (9)	-0.0073 (8)
C11	0.0335 (12)	0.0188 (10)	0.0291 (11)	-0.0075 (9)	-0.0110 (9)	-0.0046 (8)
C12	0.0237 (10)	0.0187 (10)	0.0221 (10)	-0.0032 (8)	-0.0059 (8)	-0.0016 (8)
C13	0.0327 (12)	0.0179 (10)	0.0280 (11)	-0.0058 (9)	-0.0071 (9)	0.0017 (8)
C14	0.0350 (12)	0.0213 (11)	0.0230 (10)	-0.0057 (9)	-0.0078 (9)	0.0049 (8)
C15	0.0289 (11)	0.0218 (10)	0.0185 (9)	-0.0042 (8)	-0.0070 (8)	0.0004 (8)
C16	0.0190 (9)	0.0163 (9)	0.0196 (9)	-0.0013 (7)	-0.0058 (7)	-0.0011 (7)
C17	0.0184 (9)	0.0177 (9)	0.0195 (9)	-0.0022 (7)	-0.0065 (7)	-0.0010 (7)

supplementary materials

Geometric parameters (Å, °)

Cu1—C11	2.4717 (6)	C5—H5C	0.9800
Cu1—O1	1.927 (2)	C6—C7	1.396 (3)
Cu1—O2	1.936 (2)	C6—H6	0.9500
Cu1—N1	2.038 (2)	C7—C8	1.372 (3)
Cu1—N2	2.025 (2)	C7—H7	0.9500
O1—C2	1.271 (3)	C8—C9	1.420 (3)
O2—C4	1.273 (3)	C8—H8	0.9500
N1—C6	1.328 (3)	C9—C17	1.399 (3)
N1—C17	1.357 (3)	C9—C10	1.433 (3)
N2—C15	1.332 (3)	C10—C11	1.359 (3)
N2—C16	1.360 (3)	C10—H10	0.9500
C1—C2	1.507 (3)	C11—C12	1.431 (3)
C1—H1A	0.9800	C11—H11	0.9500
C1—H1B	0.9800	C12—C16	1.401 (3)
C1—H1C	0.9800	C12—C13	1.418 (3)
C2—C3	1.390 (3)	C13—C14	1.370 (3)
C3—C4	1.391 (3)	C13—H13	0.9500
C3—H3	0.9500	C14—C15	1.402 (3)
C4—C5	1.506 (3)	C14—H14	0.9500
C5—H5A	0.9800	C15—H15	0.9500
C5—H5B	0.9800	C16—C17	1.435 (3)
O1—Cu1—O2	93.82 (7)	H5B—C5—H5C	109.5
O1—Cu1—N2	88.87 (7)	N1—C6—C7	122.4 (2)
O2—Cu1—N2	164.40 (7)	N1—C6—H6	118.8
O1—Cu1—N1	158.42 (7)	C7—C6—H6	118.8
O2—Cu1—N1	91.13 (7)	C8—C7—C6	120.18 (19)
N1—Cu1—N2	81.04 (7)	C8—C7—H7	119.9
O1—Cu1—C11	102.88 (5)	C6—C7—H7	119.9
O2—Cu1—C11	97.34 (5)	C7—C8—C9	118.7 (2)
N2—Cu1—C11	97.03 (5)	C7—C8—H8	120.6
N1—Cu1—C11	97.28 (5)	C9—C8—H8	120.6
C2—O1—Cu1	124.95 (14)	C17—C9—C10	119.20 (19)
C4—O2—Cu1	124.08 (15)	C17—C9—C8	116.96 (19)
C6—N1—C17	118.08 (18)	C10—C9—C8	123.8 (2)
C6—N1—Cu1	129.04 (15)	C11—C10—C9	120.7 (2)
C17—N1—Cu1	112.82 (13)	C11—C10—H10	119.7
C15—N2—C16	117.97 (18)	C9—C10—H10	119.7
C15—N2—Cu1	128.74 (15)	C10—C11—C12	121.3 (2)
C16—N2—Cu1	113.24 (13)	C10—C11—H11	119.4
C2—C1—H1A	109.5	C12—C11—H11	119.4
C2—C1—H1B	109.5	C16—C12—C13	116.90 (19)
H1A—C1—H1B	109.5	C16—C12—C11	118.84 (19)
C2—C1—H1C	109.5	C13—C12—C11	124.2 (2)
H1A—C1—H1C	109.5	C14—C13—C12	119.0 (2)
H1B—C1—H1C	109.5	C14—C13—H13	120.5
O1—C2—C3	125.0 (2)	C12—C13—H13	120.5

O1—C2—C1	115.3 (2)	C13—C14—C15	120.21 (19)
C3—C2—C1	119.7 (2)	C13—C14—H14	119.9
C2—C3—C4	125.1 (2)	C15—C14—H14	119.9
C2—C3—H3	117.4	N2—C15—C14	122.2 (2)
C4—C3—H3	117.4	N2—C15—H15	118.9
O2—C4—C3	125.5 (2)	C14—C15—H15	118.9
O2—C4—C5	116.0 (2)	N2—C16—C12	123.74 (18)
C3—C4—C5	118.4 (2)	N2—C16—C17	116.22 (18)
C4—C5—H5A	109.5	C12—C16—C17	120.03 (19)
C4—C5—H5B	109.5	N1—C17—C9	123.64 (18)
H5A—C5—H5B	109.5	N1—C17—C16	116.42 (18)
C4—C5—H5C	109.5	C9—C17—C16	119.93 (18)
H5A—C5—H5C	109.5		
O2—Cu1—O1—C2	10.33 (19)	C6—C7—C8—C9	-0.3 (3)
N2—Cu1—O1—C2	174.95 (19)	C7—C8—C9—C17	-0.2 (3)
N1—Cu1—O1—C2	113.2 (2)	C7—C8—C9—C10	-179.3 (2)
Cl1—Cu1—O1—C2	-88.10 (18)	C17—C9—C10—C11	-1.2 (3)
O1—Cu1—O2—C4	-11.32 (19)	C8—C9—C10—C11	177.9 (2)
N2—Cu1—O2—C4	-110.9 (3)	C9—C10—C11—C12	0.8 (4)
N1—Cu1—O2—C4	-170.31 (18)	C10—C11—C12—C16	0.2 (3)
Cl1—Cu1—O2—C4	92.21 (18)	C10—C11—C12—C13	-178.6 (2)
O1—Cu1—N1—C6	-115.3 (2)	C16—C12—C13—C14	0.1 (3)
O2—Cu1—N1—C6	-12.0 (2)	C11—C12—C13—C14	178.9 (2)
N2—Cu1—N1—C6	-178.4 (2)	C12—C13—C14—C15	0.7 (3)
Cl1—Cu1—N1—C6	85.57 (19)	C16—N2—C15—C14	0.7 (3)
O1—Cu1—N1—C17	67.4 (2)	Cu1—N2—C15—C14	177.97 (16)
O2—Cu1—N1—C17	170.77 (15)	C13—C14—C15—N2	-1.1 (4)
N2—Cu1—N1—C17	4.33 (14)	C15—N2—C16—C12	0.1 (3)
Cl1—Cu1—N1—C17	-91.68 (14)	Cu1—N2—C16—C12	-177.56 (16)
O1—Cu1—N2—C15	17.41 (19)	C15—N2—C16—C17	-178.63 (19)
O2—Cu1—N2—C15	117.6 (3)	Cu1—N2—C16—C17	3.7 (2)
N1—Cu1—N2—C15	178.3 (2)	C13—C12—C16—N2	-0.5 (3)
Cl1—Cu1—N2—C15	-85.43 (19)	C11—C12—C16—N2	-179.4 (2)
O1—Cu1—N2—C16	-165.19 (15)	C13—C12—C16—C17	178.19 (19)
O2—Cu1—N2—C16	-65.0 (3)	C11—C12—C16—C17	-0.7 (3)
N1—Cu1—N2—C16	-4.33 (14)	C6—N1—C17—C9	-0.3 (3)
Cl1—Cu1—N2—C16	91.97 (14)	Cu1—N1—C17—C9	177.29 (16)
Cu1—O1—C2—C3	-3.0 (3)	C6—N1—C17—C16	178.75 (19)
Cu1—O1—C2—C1	178.82 (16)	Cu1—N1—C17—C16	-3.7 (2)
O1—C2—C3—C4	-7.9 (4)	C10—C9—C17—N1	179.7 (2)
C1—C2—C3—C4	170.2 (2)	C8—C9—C17—N1	0.5 (3)
Cu1—O2—C4—C3	5.2 (3)	C10—C9—C17—C16	0.7 (3)
Cu1—O2—C4—C5	-175.75 (16)	C8—C9—C17—C16	-178.49 (19)
C2—C3—C4—O2	6.6 (4)	N2—C16—C17—N1	0.0 (3)
C2—C3—C4—C5	-172.4 (2)	C12—C16—C17—N1	-178.79 (19)
C17—N1—C6—C7	-0.3 (3)	N2—C16—C17—C9	179.10 (19)
Cu1—N1—C6—C7	-177.38 (17)	C12—C16—C17—C9	0.3 (3)
N1—C6—C7—C8	0.6 (4)		

Fig. 1

