

Hemiaquabis(2-fluorobenzoato- κ^2O,O')-bis(1,10-phenanthroline- κ^2N,N')lead(II) dihydrate

Su-Fang Ye and Bi-Song Zhang*

College of Materials Science and Chemical Engineering, Jinhua College of Profession and Technology, Jinhua, Zhejiang 321017, People's Republic of China

Correspondence e-mail: zbs_jy@163.com


Received 14 June 2009; accepted 13 July 2009

Key indicators: single-crystal X-ray study; $T = 290$ K; mean $\sigma(C-C) = 0.008$ Å; disorder in main residue; R factor = 0.032; wR factor = 0.078; data-to-parameter ratio = 22.0.

In the title compound, $[Pb(C_7H_4FO_2)_2(C_{12}H_8N_2)_2(H_2O)_{0.5}] \cdot 2H_2O$, the Pb^{II} atom is coordinated by four N atoms from two bidentate chelating 1,10-phenanthroline (phen) ligands, four O atoms from two 2-fluorobenzoate ligands and a half-occupied water molecule in an irregular coordination geometry. One carboxylate O atom and two F atoms are each disordered over two sites with occupancy factors of 0.558 (6) and 0.442 (6). The two crystallographically independent phen ligands are co-planar [dihedral angle $0.0(2)^\circ$]. Centroid-centroid distances of 3.659 (7) and 3.687 (7) Å indicate π - π stacking interactions between neighboring phen ligands. In the crystal, $O-H \cdots O$, $C-H \cdots F$ and $C-H \cdots O$ hydrogen bonds link the complex molecules and uncoordinated water molecules into a supramolecular network.

Related literature

For other complexes with a 2(or 4)-fluorobenzoate ligand, see: Zhang *et al.* (2005). For related structures, see: Zhang (2004, 2005, 2006*a,b,c*).



Experimental

Crystal data

$[Pb(C_7H_4FO_2)_2(C_{12}H_8N_2)_2 \cdot (H_2O)_{0.5}] \cdot 2H_2O$
 $M_r = 890.84$
 Triclinic, $P\bar{1}$
 $a = 9.833(2)$ Å
 $b = 11.568(2)$ Å
 $c = 15.766(3)$ Å
 $\alpha = 81.11(3)^\circ$

$\beta = 77.23(3)^\circ$
 $\gamma = 86.20(3)^\circ$
 $V = 1727.0(6)$ Å³
 $Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 4.95$ mm⁻¹
 $T = 290$ K
 $0.34 \times 0.19 \times 0.16$ mm

Data collection

Bruker SMART APEX CCD diffractometer
 Absorption correction: multi-scan (SADABS; Sheldrick, 1996)
 $T_{min} = 0.330$, $T_{max} = 0.448$

28426 measured reflections
 10232 independent reflections
 7916 reflections with $I > 2\sigma(I)$
 $R_{int} = 0.028$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.032$
 $wR(F^2) = 0.078$
 $S = 1.02$
 10232 reflections
 466 parameters

3 restraints
 H-atom parameters constrained
 $\Delta\rho_{max} = 1.33$ e Å⁻³
 $\Delta\rho_{min} = -0.77$ e Å⁻³

Table 1

Selected bond lengths (Å).

Pb1—O1	2.957 (8)	Pb1—N1	2.796 (3)
Pb1—O1'	2.866 (9)	Pb1—N2	2.656 (3)
Pb1—O2	2.631 (3)	Pb1—N3	2.768 (3)
Pb1—O3	2.575 (3)	Pb1—N4	2.906 (3)
Pb1—O4	2.570 (3)	Pb1—O7W	2.965 (7)

Table 2

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
O5W—H5WA \cdots O2	0.85	1.98	2.796 (4)	162
O5W—H5WB \cdots O6W	0.85	1.97	2.757 (5)	153
O6W—H6WA \cdots O5W ⁱ	0.85	1.98	2.809 (6)	163
O6W—H6WB \cdots O4	0.85	1.97	2.818 (3)	175
O6W—H6WA \cdots O5W ⁱ	0.85	1.98	2.809 (6)	163
O7W—H7WA \cdots O1'	0.85	1.98	2.496 (5)	118
O7W—H7WB \cdots O1' ⁱⁱⁱ	0.85	1.99	2.565 (2)	124
C8—H8 \cdots F1 ⁱⁱⁱ	0.93	2.54	3.310 (7)	141
C30—H30 \cdots F1 ⁱⁱⁱ	0.93	2.50	3.032 (12)	115
C29—H29 \cdots O3 ^{iv}	0.93	2.46	3.311 (5)	153

Symmetry codes: (i) $-x+2, -y+2, -z$; (ii) $-x+1, -y+2, -z+1$; (iii) $x+1, y, z$; (iv) $-x+1, -y+1, -z+1$.

Data collection: SMART (Bruker, 2007); cell refinement: SAINT (Bruker, 2007); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

The authors gratefully acknowledge financial support from the Jinhua Municipal Science and Technology (grant No. 2003-01-179).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HY2206).

References

Bruker (2007). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.

- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Zhang, B.-S. (2004). *Z. Kristallogr. New Cryst. Struct.* **219**, 483–484.
Zhang, B.-S. (2005). *Z. Kristallogr. New Cryst. Struct.* **220**, 73–74.
Zhang, B.-S. (2006a). *Acta Cryst.* **E62**, m2645–m2647.
Zhang, B.-S. (2006b). *Z. Kristallogr. New Cryst. Struct.* **221**, 191–194.
Zhang, B. S. (2006c). *Z. Kristallogr. New Cryst. Struct.* **221**, 355–356.
Zhang, B.-S., Zeng, X.-R., Yu, Y.-Y., Fang, X.-N. & Huang, C.-F. (2005). *Z. Kristallogr. New Cryst. Struct.* **220**, 75–76.

supplementary materials

Acta Cryst. (2009). E65, m936-m937 [doi:10.1107/S1600536809027524]

Hemiaquabis(2-fluorobenzoato- κ^2O,O')bis(1,10-phenanthroline- κ^2N,N')lead(II) dihydrate

S.-F. Ye and B.-S. Zhang

Comment

We have prepared the title complex by the hydrothermal reaction of freshly prepared $PbCO_3$ with 1,10-phenanthroline (phen) and 2-fluorobenzoic acid in CH_3OH/H_2O , and report here its crystal structure (Fig. 1). The title compound has a structure similar to those of complexes with halobenzoate ligands, $X-C_6H_4COO^-$, where X is F, Cl, Br and I (Zhang, 2004, 2005, 2006a,b,c; Zhang *et al.*, 2005). The asymmetric unit of the title compound consists of a $[Pb(C_7H_4FO_2)_2(C_{12}H_8N_2)_2(H_2O)_{0.5}]$ complex molecule and two uncoordinated water molecules. The Pb^{II} atom is coordinated by four N atoms from two bidentate chelating phen ligands, four O atoms from two 2-fluorobenzoate ligands and a half-occupied water molecule in an irregular coordination geometry, with $Pb-N$ bond lengths in the range of 2.656 (3) to 2.906 (3) Å and $Pb-O$ bond lengths in the range of 2.570 (3) to 2.965 (7) Å (Table 1). The centroid-centroid distances of 3.659 (7) and 3.687 (7) Å indicate $\pi-\pi$ stacking interactions between the neighboring phen ligands (Fig. 2). $O-H\cdots O$, $C-H\cdots F$ and $C-H\cdots O$ hydrogen bonds are also present (Table 2 and Fig. 3). A combination of the $\pi-\pi$ stacking interactions and hydrogen bonds leads to a supramolecular network.

Experimental

$Pb(CH_3COO)_2 \cdot 3H_2O$ (0.17 g, 0.45 mmol) was dissolved in appropriate amount of water, and then 1M Na_2CO_3 solution was added. $PbCO_3$ was obtained by filtration, which was then washed with distilled water for 5 times. The freshly prepared $PbCO_3$, phen (0.05 g, 0.25 mmol), 2-fluorobenzoic acid (0.04 g, 0.29 mmol), CH_3OH/H_2O (v/v = 1:2, 15 ml) were mixed and stirred for 1.5 h. Subsequently, the resulting cream suspension was heated in a 23 ml Teflon-lined stainless steel autoclave at 423 K for one week. After the autoclave was cooled to room temperature, the solid was filtered off. The resulting filtrate was allowed to stand at room temperature, and evaporation for 3 weeks afforded colorless transparent pillar-like single crystals. Analysis calculated for $C_{38}H_{29}F_2N_4O_{6.5}Pb$: C 51.19, H 3.26, N 6.28%; found: C 51.06, H 3.06, N 6.13%.

Refinement

The disordered O and F atoms on the ligands were refined isotropically. H atoms on C atoms were positioned geometrically and refined as riding atoms, with $C-H = 0.93$ Å and with $U_{iso}(H) = 1.2U_{eq}(C)$. H atoms of water molecules were located in a difference Fourier map and refined with restraints of $O-H = 0.85$ (1) Å and $U_{iso}(H) = 1.5U_{eq}(O)$. The largest peak in the final difference Fourier map is 0.24 Å from atom H35 and the deepest hole is 0.52 Å from atom F2'.

Figures




Fig. 1. Molecular structure of the title compound. Displacement ellipsoids are drawn at the 50% probability level. H atoms have been omitted for clarity. Dashed line shows the bond between Pb1 and half-occupied O7W.




Fig. 2. The π - π stacking interactions (dashed double arrows), with the centroid-centroid distances of 3.659 (7) and 3.687 (7) Å.




Fig. 3. The hydrogen bonds (thin dashed lines) in the title compound.

Hemiaquabis(2-fluorobenzoato- κ^2O,O')bis(1,10-phenanthroline- κ^2N,N')lead(II) dihydrate

Crystal data

[Pb(C₇H₄FO₂)₂(C₁₂H₈N₂)₂(H₂O)_{0.5}]₂·2H₂O

M_r = 890.84

Triclinic, $P\bar{1}$

Hall symbol: -P 1

a = 9.833 (2) Å

b = 11.568 (2) Å

c = 15.766 (3) Å

α = 81.11 (3)°

β = 77.23 (3)°

γ = 86.20 (3)°

V = 1727.0 (6) Å³

Z = 2

F_{000} = 874

D_x = 1.713 Mg m⁻³

Mo $K\alpha$ radiation, λ = 0.71073 Å

Cell parameters from 8657 reflections

θ = 2.7–30.5°

μ = 4.95 mm⁻¹

T = 290 K

Pillar-like, colorless

0.34 × 0.19 × 0.16 mm

Data collection

Bruker SMART APEX CCD
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

T = 290 K

ϕ and ω scans

10232 independent reflections

7916 reflections with $I > 2\sigma(I)$

R_{int} = 0.028

θ_{max} = 30.5°

θ_{min} = 2.7°

Absorption correction: multi-scan
(SADABS; Sheldrick, 1996) $h = -14 \rightarrow 13$
 $T_{\min} = 0.330$, $T_{\max} = 0.448$ $k = -16 \rightarrow 16$
 28426 measured reflections $l = -22 \rightarrow 22$

Refinement

Refinement on F^2 Secondary atom site location: difference Fourier map
 Least-squares matrix: full Hydrogen site location: inferred from neighbouring sites
 $R[F^2 > 2\sigma(F^2)] = 0.032$ H-atom parameters constrained
 $wR(F^2) = 0.078$ $w = 1/[\sigma^2(F_o^2) + (0.0433P)^2]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $S = 1.02$ $(\Delta/\sigma)_{\max} = 0.002$
 10232 reflections $\Delta\rho_{\max} = 1.33 \text{ e } \text{\AA}^{-3}$
 466 parameters $\Delta\rho_{\min} = -0.77 \text{ e } \text{\AA}^{-3}$
 3 restraints Extinction correction: none
 Primary atom site location: structure-invariant direct methods

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Pb1	0.694779 (12)	0.832399 (10)	0.325534 (7)	0.04599 (5)	
N1	0.8374 (4)	0.7952 (3)	0.4629 (2)	0.0650 (8)	
N2	0.9505 (3)	0.9112 (3)	0.29959 (19)	0.0529 (7)	
N3	0.4998 (3)	0.7983 (3)	0.22999 (19)	0.0535 (7)	
N4	0.4509 (3)	0.6922 (3)	0.4003 (2)	0.0570 (7)	
O1	0.6395 (9)	1.0811 (7)	0.3491 (5)	0.075 (2)*	0.442 (6)
O1'	0.5776 (10)	1.0611 (8)	0.3514 (6)	0.112 (3)*	0.558 (6)
O2	0.6834 (3)	1.0286 (2)	0.21751 (18)	0.0670 (7)	
O3	0.7824 (3)	0.6229 (2)	0.30117 (19)	0.0718 (8)	
O4	0.8193 (3)	0.7548 (2)	0.18316 (16)	0.0564 (6)	
O5W	0.8186 (4)	1.0878 (3)	0.0418 (2)	0.0933 (10)	
H5WA	0.7855	1.0837	0.0967	0.140*	
H5WB	0.8144	1.0180	0.0321	0.140*	
O6W	0.8963 (4)	0.8634 (3)	0.0076 (2)	0.1016 (11)	
H6WA	0.9847	0.8631	-0.0092	0.152*	
H6WB	0.8676	0.8313	0.0603	0.152*	
O7W	0.5357 (7)	0.9161 (8)	0.4888 (6)	0.114 (3)	0.50
H7WA	0.5897	0.9728	0.4679	0.170*	0.50
H7WB	0.5360	0.8824	0.5406	0.170*	0.50
F1	0.4835 (5)	1.1309 (4)	0.1281 (3)	0.0642 (16)*	0.442 (6)
F1'	0.6161 (11)	1.3066 (9)	0.3310 (7)	0.174 (4)*	0.558 (6)
F2	0.9722 (9)	0.4352 (8)	0.2413 (6)	0.1118 (17)*	0.442 (6)
F2'	0.9188 (7)	0.4233 (6)	0.2639 (5)	0.1118 (17)*	0.558 (6)
C1	0.7849 (6)	0.7420 (4)	0.5430 (3)	0.0860 (14)	
H1	0.6973	0.7103	0.5529	0.103*	

supplementary materials

C2	0.8501 (7)	0.7302 (5)	0.6125 (3)	0.0926 (15)	
H2	0.8076	0.6915	0.6672	0.111*	
C3	0.9756 (7)	0.7753 (4)	0.6002 (3)	0.0866 (15)	
H3	1.0213	0.7676	0.6467	0.104*	
C4	1.0395 (5)	0.8345 (4)	0.5173 (3)	0.0661 (11)	
C5	1.1716 (5)	0.8861 (4)	0.5000 (4)	0.0812 (14)	
H5	1.2202	0.8820	0.5448	0.097*	
C6	1.2265 (4)	0.9402 (5)	0.4201 (4)	0.0812 (14)	
H6	1.3146	0.9710	0.4099	0.097*	
C7	1.1553 (4)	0.9528 (3)	0.3497 (3)	0.0618 (9)	
C8	1.2079 (4)	1.0116 (4)	0.2665 (4)	0.0762 (13)	
H8	1.2944	1.0457	0.2548	0.091*	
C9	1.1344 (5)	1.0203 (4)	0.2018 (3)	0.0723 (11)	
H9	1.1690	1.0605	0.1461	0.087*	
C10	1.0081 (4)	0.9681 (4)	0.2211 (3)	0.0649 (10)	
H10	0.9593	0.9727	0.1764	0.078*	
C11	1.0242 (3)	0.9021 (3)	0.3638 (2)	0.0516 (8)	
C12	0.9641 (4)	0.8418 (3)	0.4497 (2)	0.0554 (8)	
C13	0.6226 (4)	1.0950 (4)	0.2690 (2)	0.0671 (10)	
C14	0.5605 (4)	1.2106 (4)	0.2336 (3)	0.0667 (10)	
C15	0.5581 (7)	1.3120 (6)	0.2687 (4)	0.1057 (18)	
H15	0.5976	1.3075	0.3176	0.127*	0.442 (6)
C16	0.5055 (10)	1.4165 (7)	0.2405 (7)	0.162 (4)	
H16	0.5106	1.4823	0.2668	0.195*	
C17	0.4438 (12)	1.4207 (9)	0.1708 (8)	0.189 (6)	
H17	0.4027	1.4909	0.1500	0.227*	
C18	0.4406 (8)	1.3262 (7)	0.1311 (5)	0.134 (3)	
H18	0.3991	1.3324	0.0830	0.161*	
C19	0.4987 (5)	1.2199 (5)	0.1615 (3)	0.0873 (15)	
H19	0.4965	1.1549	0.1337	0.105*	0.558 (6)
C21	0.5238 (5)	0.8488 (4)	0.1454 (3)	0.0667 (10)	
H21	0.6039	0.8915	0.1229	0.080*	
C22	0.4337 (5)	0.8395 (4)	0.0906 (3)	0.0769 (12)	
H22	0.4534	0.8762	0.0327	0.092*	
C23	0.3175 (5)	0.7773 (4)	0.1216 (3)	0.0787 (13)	
H23	0.2570	0.7707	0.0850	0.094*	
C24	0.2883 (4)	0.7231 (4)	0.2079 (3)	0.0644 (10)	
C25	0.1654 (5)	0.6569 (5)	0.2460 (4)	0.0849 (14)	
H25	0.1015	0.6494	0.2119	0.102*	
C26	0.1415 (4)	0.6059 (5)	0.3298 (4)	0.0864 (14)	
H26	0.0612	0.5635	0.3526	0.104*	
C27	0.2356 (4)	0.6151 (3)	0.3845 (3)	0.0640 (10)	
C28	0.2154 (5)	0.5624 (4)	0.4730 (4)	0.0818 (14)	
H28	0.1364	0.5190	0.4978	0.098*	
C29	0.3075 (5)	0.5735 (4)	0.5217 (3)	0.0794 (13)	
H29	0.2946	0.5380	0.5799	0.095*	
C30	0.4237 (4)	0.6403 (4)	0.4825 (3)	0.0700 (11)	
H30	0.4869	0.6489	0.5170	0.084*	
C31	0.3572 (3)	0.6795 (3)	0.3515 (2)	0.0508 (7)	

C32	0.3842 (3)	0.7350 (3)	0.2611 (2)	0.0519 (8)
C33	0.8114 (3)	0.6510 (3)	0.2198 (2)	0.0519 (8)
C34	0.8350 (4)	0.5560 (3)	0.1620 (3)	0.0624 (9)
C35	0.7819 (6)	0.5800 (5)	0.0821 (3)	0.1034 (18)
H35	0.7385	0.6508	0.0647	0.124*
C36	0.8019 (8)	0.4873 (7)	0.0339 (5)	0.130 (2)
H36	0.7718	0.4983	-0.0188	0.156*
C37	0.8619 (9)	0.3820 (7)	0.0576 (6)	0.133 (3)
H37	0.8681	0.3236	0.0224	0.160*
C38	0.9122 (7)	0.3607 (5)	0.1308 (5)	0.114 (2)
H38	0.9570	0.2897	0.1463	0.137*
C39	0.8944 (5)	0.4497 (4)	0.1824 (3)	0.0824 (12)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Pb1	0.05017 (8)	0.04824 (8)	0.04002 (7)	-0.00731 (5)	-0.01187 (5)	-0.00214 (5)
N1	0.084 (2)	0.066 (2)	0.0500 (17)	-0.0199 (17)	-0.0252 (16)	0.0001 (14)
N2	0.0529 (15)	0.0569 (17)	0.0524 (16)	-0.0064 (13)	-0.0191 (13)	-0.0061 (13)
N3	0.0539 (16)	0.0579 (17)	0.0510 (16)	-0.0040 (13)	-0.0176 (13)	-0.0055 (13)
N4	0.0508 (16)	0.0616 (19)	0.0534 (17)	-0.0048 (13)	-0.0081 (13)	0.0045 (14)
O2	0.0816 (18)	0.0576 (16)	0.0581 (15)	0.0021 (13)	-0.0133 (13)	-0.0013 (12)
O3	0.095 (2)	0.0565 (16)	0.0581 (16)	-0.0063 (14)	-0.0101 (14)	0.0023 (12)
O4	0.0677 (15)	0.0500 (14)	0.0494 (13)	-0.0075 (11)	-0.0086 (11)	-0.0034 (11)
O5W	0.113 (2)	0.082 (2)	0.0669 (18)	0.0181 (18)	0.0005 (17)	0.0069 (15)
O6W	0.130 (3)	0.080 (2)	0.074 (2)	0.008 (2)	0.006 (2)	0.0087 (17)
O7W	0.086 (5)	0.142 (7)	0.112 (6)	-0.007 (5)	0.002 (4)	-0.048 (5)
C1	0.117 (4)	0.078 (3)	0.065 (3)	-0.028 (3)	-0.029 (3)	0.009 (2)
C2	0.144 (5)	0.083 (3)	0.055 (2)	-0.006 (3)	-0.041 (3)	0.004 (2)
C3	0.133 (4)	0.072 (3)	0.069 (3)	0.026 (3)	-0.055 (3)	-0.019 (2)
C4	0.088 (3)	0.053 (2)	0.071 (2)	0.020 (2)	-0.042 (2)	-0.0250 (19)
C5	0.077 (3)	0.090 (3)	0.101 (4)	0.027 (2)	-0.056 (3)	-0.047 (3)
C6	0.053 (2)	0.094 (3)	0.115 (4)	0.006 (2)	-0.039 (2)	-0.045 (3)
C7	0.0517 (19)	0.062 (2)	0.080 (3)	0.0078 (16)	-0.0221 (18)	-0.028 (2)
C8	0.048 (2)	0.078 (3)	0.103 (4)	-0.0112 (19)	-0.006 (2)	-0.025 (3)
C9	0.067 (2)	0.072 (3)	0.072 (3)	-0.011 (2)	-0.004 (2)	-0.004 (2)
C10	0.060 (2)	0.076 (3)	0.057 (2)	-0.0078 (19)	-0.0125 (17)	-0.0020 (19)
C11	0.0530 (18)	0.0459 (18)	0.061 (2)	0.0070 (14)	-0.0187 (16)	-0.0182 (15)
C12	0.068 (2)	0.0455 (18)	0.061 (2)	0.0095 (16)	-0.0291 (18)	-0.0156 (15)
C13	0.081 (3)	0.074 (3)	0.0417 (18)	-0.012 (2)	-0.0091 (18)	0.0050 (17)
C14	0.067 (2)	0.060 (2)	0.061 (2)	-0.0025 (18)	0.0112 (19)	-0.0094 (18)
C15	0.114 (4)	0.096 (4)	0.101 (4)	-0.006 (3)	0.006 (3)	-0.035 (3)
C16	0.155 (8)	0.069 (4)	0.225 (11)	-0.004 (5)	0.062 (7)	-0.050 (6)
C17	0.191 (10)	0.097 (6)	0.227 (13)	0.070 (7)	0.024 (9)	0.003 (7)
C18	0.153 (6)	0.114 (5)	0.122 (5)	0.057 (5)	-0.036 (5)	0.007 (4)
C19	0.087 (3)	0.077 (3)	0.084 (3)	0.013 (2)	-0.011 (3)	0.016 (3)
C21	0.077 (3)	0.075 (3)	0.050 (2)	-0.004 (2)	-0.0205 (19)	-0.0026 (18)
C22	0.103 (3)	0.082 (3)	0.051 (2)	0.011 (3)	-0.032 (2)	-0.011 (2)

supplementary materials

C23	0.089 (3)	0.079 (3)	0.085 (3)	0.012 (2)	-0.049 (3)	-0.027 (2)
C24	0.059 (2)	0.063 (2)	0.082 (3)	0.0078 (17)	-0.030 (2)	-0.028 (2)
C25	0.060 (2)	0.085 (3)	0.123 (4)	-0.008 (2)	-0.031 (3)	-0.035 (3)
C26	0.050 (2)	0.087 (3)	0.126 (4)	-0.015 (2)	-0.009 (3)	-0.034 (3)
C27	0.0493 (19)	0.050 (2)	0.089 (3)	-0.0029 (15)	-0.0022 (19)	-0.0144 (19)
C28	0.067 (3)	0.057 (2)	0.105 (4)	-0.012 (2)	0.015 (3)	-0.007 (2)
C29	0.076 (3)	0.068 (3)	0.076 (3)	-0.002 (2)	0.009 (2)	0.009 (2)
C30	0.067 (2)	0.072 (3)	0.062 (2)	-0.0066 (19)	-0.0073 (19)	0.010 (2)
C31	0.0435 (16)	0.0431 (17)	0.064 (2)	0.0003 (13)	-0.0067 (15)	-0.0110 (15)
C32	0.0489 (17)	0.0456 (18)	0.064 (2)	0.0034 (14)	-0.0162 (15)	-0.0137 (15)
C33	0.0482 (17)	0.053 (2)	0.054 (2)	-0.0059 (14)	-0.0079 (14)	-0.0057 (15)
C34	0.064 (2)	0.051 (2)	0.070 (2)	-0.0070 (17)	-0.0073 (19)	-0.0102 (18)
C35	0.129 (4)	0.124 (5)	0.070 (3)	-0.039 (3)	-0.012 (3)	-0.051 (3)
C36	0.157 (6)	0.130 (6)	0.129 (5)	0.008 (5)	-0.065 (5)	-0.053 (5)
C37	0.171 (7)	0.100 (5)	0.143 (7)	0.008 (5)	-0.040 (6)	-0.056 (5)
C38	0.133 (5)	0.060 (3)	0.146 (6)	0.016 (3)	-0.023 (5)	-0.025 (4)
C39	0.097 (3)	0.072 (3)	0.079 (3)	0.005 (2)	-0.023 (3)	-0.009 (2)

Geometric parameters (Å, °)

Pb1—O1	2.957 (8)	C8—H8	0.9300
Pb1—O1'	2.866 (9)	C9—C10	1.369 (6)
Pb1—O2	2.631 (3)	C9—H9	0.9300
Pb1—O3	2.575 (3)	C10—H10	0.9300
Pb1—O4	2.570 (3)	C11—C12	1.441 (5)
Pb1—N1	2.796 (3)	C13—C14	1.511 (6)
Pb1—N2	2.656 (3)	C14—C15	1.368 (7)
Pb1—N3	2.768 (3)	C14—C19	1.390 (7)
Pb1—N4	2.906 (3)	C15—C16	1.334 (10)
Pb1—O7W	2.965 (7)	C15—H15	0.9300
N1—C1	1.324 (5)	C16—C17	1.361 (14)
N1—C12	1.351 (5)	C16—H16	0.9300
N2—C10	1.332 (5)	C17—C18	1.346 (14)
N2—C11	1.357 (4)	C17—H17	0.9300
N3—C21	1.346 (5)	C18—C19	1.383 (7)
N3—C32	1.349 (4)	C18—H18	0.9300
N4—C30	1.319 (5)	C19—H19	0.9300
N4—C31	1.352 (5)	C21—C22	1.386 (6)
O1—C13	1.294 (9)	C21—H21	0.9300
O1'—C13	1.285 (9)	C22—C23	1.348 (7)
O2—C13	1.234 (5)	C22—H22	0.9300
O3—C33	1.247 (4)	C23—C24	1.383 (6)
O4—C33	1.249 (4)	C23—H23	0.9300
O5W—H5WA	0.85	C24—C32	1.419 (5)
O5W—H5WB	0.85	C24—C25	1.438 (6)
O6W—H6WA	0.85	C25—C26	1.336 (7)
O6W—H6WB	0.85	C25—H25	0.9300
O7W—H7WA	0.85	C26—C27	1.417 (7)
O7W—H7WB	0.85	C26—H26	0.9300

F1—C19	1.258 (7)	C27—C31	1.406 (5)
F1'—C15	1.232 (11)	C27—C28	1.410 (7)
F2—F2'	0.578 (12)	C28—C29	1.335 (7)
F2—C39	1.313 (10)	C28—H28	0.9300
F2'—C39	1.344 (8)	C29—C30	1.393 (6)
C1—C2	1.373 (6)	C29—H29	0.9300
C1—H1	0.9300	C30—H30	0.9300
C2—C3	1.334 (7)	C31—C32	1.445 (5)
C2—H2	0.9300	C33—C34	1.508 (5)
C3—C4	1.411 (7)	C34—C39	1.350 (6)
C3—H3	0.9300	C34—C35	1.450 (7)
C4—C12	1.416 (5)	C35—C36	1.387 (8)
C4—C5	1.419 (7)	C35—H35	0.9300
C5—C6	1.331 (7)	C36—C37	1.354 (10)
C5—H5	0.9300	C36—H36	0.9300
C6—C7	1.422 (6)	C37—C38	1.335 (10)
C6—H6	0.9300	C37—H37	0.9300
C7—C8	1.385 (6)	C38—C39	1.388 (8)
C7—C11	1.410 (5)	C38—H38	0.9300
C8—C9	1.364 (7)		
O4—Pb1—O3	50.53 (9)	N2—C10—C9	124.4 (4)
O4—Pb1—O2	82.50 (9)	N2—C10—H10	117.8
O3—Pb1—O2	132.87 (9)	C9—C10—H10	117.8
O4—Pb1—N2	77.43 (9)	N2—C11—C7	122.0 (3)
O3—Pb1—N2	93.60 (10)	N2—C11—C12	118.7 (3)
O2—Pb1—N2	77.32 (10)	C7—C11—C12	119.3 (3)
O4—Pb1—N3	70.63 (9)	N1—C12—C4	122.4 (4)
O3—Pb1—N3	84.91 (10)	N1—C12—C11	118.8 (3)
O2—Pb1—N3	74.60 (9)	C4—C12—C11	118.8 (4)
N2—Pb1—N3	139.58 (9)	O2—C13—O1'	123.1 (5)
O4—Pb1—N1	115.89 (10)	O2—C13—O1	120.5 (5)
O3—Pb1—N1	84.61 (10)	O2—C13—C14	119.6 (3)
O2—Pb1—N1	125.93 (10)	O1'—C13—C14	115.0 (5)
N2—Pb1—N1	60.37 (9)	O1—C13—C14	117.9 (5)
N3—Pb1—N1	158.10 (10)	C15—C14—C19	114.9 (5)
O4—Pb1—O1'	129.75 (19)	C15—C14—C13	124.3 (5)
O3—Pb1—O1'	175.83 (19)	C19—C14—C13	120.8 (4)
O2—Pb1—O1'	47.27 (19)	FI'—C15—C16	116.7 (9)
N2—Pb1—O1'	90.5 (2)	FI'—C15—C14	116.0 (8)
N3—Pb1—O1'	91.4 (2)	C16—C15—C14	127.2 (8)
N1—Pb1—O1'	98.3 (2)	C16—C15—H15	116.4
O4—Pb1—N4	107.21 (9)	C14—C15—H15	116.4
O3—Pb1—N4	76.18 (10)	C15—C16—C17	115.7 (9)
O2—Pb1—N4	122.03 (9)	C15—C16—H16	122.2
N2—Pb1—N4	160.24 (9)	C17—C16—H16	122.2
N3—Pb1—N4	57.40 (9)	C18—C17—C16	122.0 (9)
N1—Pb1—N4	101.34 (10)	C18—C17—H17	119.0
O1'—Pb1—N4	100.2 (2)	C16—C17—H17	119.0
O4—Pb1—O1	126.25 (16)	C17—C18—C19	120.5 (8)

supplementary materials

O3—Pb1—O1	171.29 (17)	C17—C18—H18	119.8
O2—Pb1—O1	45.79 (16)	C19—C18—H18	119.8
N2—Pb1—O1	77.69 (18)	F1—C19—C18	119.0 (6)
N3—Pb1—O1	101.79 (18)	F1—C19—C14	121.0 (4)
N1—Pb1—O1	90.83 (17)	C18—C19—C14	119.7 (6)
O1'—Pb1—O1	12.8 (2)	C18—C19—H19	120.1
N4—Pb1—O1	112.07 (18)	C14—C19—H19	120.1
O4—Pb1—O7W	176.63 (15)	N3—C21—C22	122.6 (4)
O3—Pb1—O7W	128.7 (2)	N3—C21—H21	118.7
O2—Pb1—O7W	98.0 (2)	C22—C21—H21	118.7
N2—Pb1—O7W	105.94 (17)	C23—C22—C21	119.8 (4)
N3—Pb1—O7W	106.24 (17)	C23—C22—H22	120.1
N1—Pb1—O7W	66.54 (18)	C21—C22—H22	120.1
O1'—Pb1—O7W	50.7 (3)	C22—C23—C24	119.9 (4)
N4—Pb1—O7W	69.71 (18)	C22—C23—H23	120.0
O1—Pb1—O7W	55.1 (2)	C24—C23—H23	120.0
C1—N1—C12	117.0 (4)	C23—C24—C32	118.0 (4)
C1—N1—Pb1	124.3 (3)	C23—C24—C25	123.2 (4)
C12—N1—Pb1	118.5 (2)	C32—C24—C25	118.8 (4)
C10—N2—C11	117.4 (3)	C26—C25—C24	121.2 (4)
C10—N2—Pb1	119.1 (2)	C26—C25—H25	119.4
C11—N2—Pb1	123.4 (2)	C24—C25—H25	119.4
C21—N3—C32	117.8 (3)	C25—C26—C27	121.5 (4)
C21—N3—Pb1	116.7 (3)	C25—C26—H26	119.3
C32—N3—Pb1	125.4 (2)	C27—C26—H26	119.3
C30—N4—C31	116.7 (3)	C31—C27—C28	116.2 (4)
C30—N4—Pb1	122.8 (3)	C31—C27—C26	120.2 (4)
C31—N4—Pb1	120.5 (2)	C28—C27—C26	123.6 (4)
C13—O1—Pb1	84.5 (4)	C29—C28—C27	121.2 (4)
C13—O1'—Pb1	88.6 (5)	C29—C28—H28	119.4
C13—O2—Pb1	101.0 (2)	C27—C28—H28	119.4
C33—O3—Pb1	92.0 (2)	C28—C29—C30	117.8 (4)
C33—O4—Pb1	92.2 (2)	C28—C29—H29	121.1
H5WA—O5W—H5WB	103.4	C30—C29—H29	121.1
H6WA—O6W—H6WB	114.0	N4—C30—C29	125.0 (4)
Pb1—O7W—H7WA	79.7	N4—C30—H30	117.5
Pb1—O7W—H7WB	125.1	C29—C30—H30	117.5
H7WA—O7W—H7WB	117.9	N4—C31—C27	123.2 (4)
F2'—F2—C39	80.5 (15)	N4—C31—C32	118.0 (3)
F2—F2'—C39	74.4 (14)	C27—C31—C32	118.8 (3)
N1—C1—C2	124.8 (5)	N3—C32—C24	121.8 (4)
N1—C1—H1	117.6	N3—C32—C31	118.6 (3)
C2—C1—H1	117.6	C24—C32—C31	119.5 (3)
C3—C2—C1	118.9 (5)	O3—C33—O4	123.2 (3)
C3—C2—H2	120.6	O3—C33—C34	119.0 (3)
C1—C2—H2	120.6	O4—C33—C34	117.8 (3)
C2—C3—C4	120.4 (4)	C39—C34—C35	118.9 (4)
C2—C3—H3	119.8	C39—C34—C33	124.4 (4)
C4—C3—H3	119.8	C35—C34—C33	116.6 (4)

C3—C4—C12	116.5 (4)	C36—C35—C34	113.7 (6)
C3—C4—C5	123.4 (4)	C36—C35—H35	123.2
C12—C4—C5	120.1 (4)	C34—C35—H35	123.2
C6—C5—C4	120.4 (4)	C37—C36—C35	125.0 (7)
C6—C5—H5	119.8	C37—C36—H36	117.5
C4—C5—H5	119.8	C35—C36—H36	117.5
C5—C6—C7	122.5 (4)	C38—C37—C36	121.1 (6)
C5—C6—H6	118.8	C38—C37—H37	119.4
C7—C6—H6	118.8	C36—C37—H37	119.4
C8—C7—C11	117.3 (4)	C37—C38—C39	116.6 (6)
C8—C7—C6	123.8 (4)	C37—C38—H38	121.7
C11—C7—C6	119.0 (4)	C39—C38—H38	121.7
C9—C8—C7	120.8 (4)	F2—C39—C34	119.5 (6)
C9—C8—H8	119.6	F2 ⁱ —C39—C34	116.9 (5)
C7—C8—H8	119.6	F2—C39—C38	113.7 (6)
C8—C9—C10	118.0 (4)	F2 ⁱ —C39—C38	117.8 (6)
C8—C9—H9	121.0	C34—C39—C38	124.6 (5)
C10—C9—H9	121.0		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O5W—H5WA \cdots O2	0.85	1.98	2.796 (4)	162
O5W—H5WB \cdots O6W	0.85	1.97	2.757 (5)	153
O6W—H6WA \cdots O5W ⁱ	0.85	1.98	2.809 (6)	163
O6W—H6WB \cdots O4	0.85	1.97	2.818 (3)	175
O6W—H6WA \cdots O5W ⁱ	0.85	1.98	2.809 (6)	163
O7W—H7WA \cdots O1 ⁱ	0.85	1.98	2.496 (5)	118
O7W—H7WB \cdots O1 ⁱⁱⁱ	0.85	1.99	2.565 (2)	124
C8—H8 \cdots F1 ⁱⁱⁱ	0.93	2.54	3.310 (7)	141
C30—H30 \cdots F1 ⁱⁱⁱ	0.93	2.50	3.032 (12)	115
C29—H29 \cdots O3 ^{iv}	0.93	2.46	3.311 (5)	153

Symmetry codes: (i) $-x+2, -y+2, -z$; (ii) $-x+1, -y+2, -z+1$; (iii) $x+1, y, z$; (iv) $-x+1, -y+1, -z+1$.

Fig. 1




Fig. 2




Fig. 3

