

Bis(2-fluorobenzoato- κ O)bis(pyridin-2-amine- κ N¹)zinc(II)

Jian-Quan Wang, Ya-Wen Zhang and Lin Cheng*

School of Chemistry and Chemical Engineering, Southeast University, Nanjing 211189, People's Republic of China
 Correspondence e-mail: cep02chl@yahoo.com.cn


Received 3 July 2009; accepted 15 July 2009

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.006$ Å; disorder in main residue; R factor = 0.043; wR factor = 0.131; data-to-parameter ratio = 13.9.

In the title compound, $[\text{Zn}(\text{C}_7\text{H}_4\text{FO}_2)_2(\text{C}_5\text{H}_6\text{N}_2)_2]$ or $[\text{Zn}(\text{fa})_2(2\text{-pa})_2]$ (Hfa is 2-fluorobenzoic acid and 2-pa = pyridin-2-amine), the asymmetric unit contains one Zn^{II} cation, two fa ligands and two 2-pa ligands, wherein the Zn^{II} displays a distorted tetrahedral geometry, being surrounded by two monodentate fa ligands with $\text{Zn}-\text{O}$ distances of 1.962 (2) and 1.976 (3) Å, and by two 2-pa ligands with distances involving pyridyl N atoms of 2.069 (2) and 2.056 (2) Å. The F atoms of the fa ligands are equally disordered over two sites, *viz.* the 2- and 6-positions of fa. The mononuclear complex molecules are joined by $\text{N}-\text{H}\cdots\text{O}$ and $\text{N}-\text{H}\cdots\text{F}$ hydrogen bonds into a two-dimensional layer, which is further constructed into a three-dimensional supra-molecular network by weak $\text{C}-\text{H}\cdots\text{F}$ interactions and effective $\pi-\pi$ stacking [centroid-centroid separation of 3.74 (3) Å] between the interlayer aromatic rings and adjacent heterocycles.

Related literature

For related structures, see: Darensbourg *et al.* (2002). For crystal engineering, see: Fyfe & Stoddart (1997).



Experimental

Crystal data

$[\text{Zn}(\text{C}_7\text{H}_4\text{FO}_2)_2(\text{C}_5\text{H}_6\text{N}_2)_2]$
 $M_r = 531.83$
 Monoclinic, $P2_1/c$
 $a = 9.1259$ (7) Å
 $b = 11.1020$ (9) Å
 $c = 24.5707$ (17) Å
 $\beta = 108.048$ (2)°

$V = 2366.9$ (3) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 1.09$ mm⁻¹
 $T = 293$ K
 $0.25 \times 0.20 \times 0.18$ mm

Data collection

Bruker SMART APEX CCD diffractometer
 Absorption correction: multi-scan (SADABS; Sheldrick, 2000)
 $T_{\text{min}} = 0.772$, $T_{\text{max}} = 0.828$

13150 measured reflections
 4645 independent reflections
 3838 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.016$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.043$
 $wR(F^2) = 0.131$
 $S = 1.04$
 4645 reflections

334 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.50$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.28$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N2}-\text{H2A}\cdots\text{O3}$	0.86	2.06	2.881 (3)	159
$\text{N2}-\text{H2B}\cdots\text{O1}^{\text{i}}$	0.86	2.03	2.864 (3)	162
$\text{N4}-\text{H4B}\cdots\text{O2}$	0.86	2.02	2.838 (4)	159
$\text{N4}-\text{H4C}\cdots\text{O4}^{\text{ii}}$	0.86	2.03	2.884 (4)	175
$\text{N4}-\text{H4B}\cdots\text{F2}$	0.86	2.52	3.143 (5)	130
$\text{C5}-\text{H5A}\cdots\text{F4}^{\text{iii}}$	0.93	2.34	3.147 (6)	144
$\text{C20}-\text{H20A}\cdots\text{F3}^{\text{iv}}$	0.93	2.47	3.244 (10)	141

Symmetry codes: (i) $-x + 2, -y, -z + 1$; (ii) $-x + 1, y - \frac{1}{2}, -z + \frac{1}{2}$; (iii) $-x + 2, y - \frac{1}{2}, -z + \frac{1}{2}$; (iv) $-x + 1, -y, -z + 1$.

Data collection: SMART (Bruker, 2000); cell refinement: SAINT (Bruker, 2000); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

The authors thank the Program for Young Excellent Talents in Southeast University for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: PV2177).

References

- Bruker (2000). SMART and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
 Darensbourg, D. J., Wildeson, J. R. & Yarbrough, J. C. (2002). *Inorg. Chem.* **41**, 973–980.
 Fyfe, M. C. T. & Stoddart, J. F. (1997). *Acc. Chem. Res.* **30**, 393–401.
 Sheldrick, G. M. (2000). SADABS. University of Göttingen, Germany.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supplementary materials

Acta Cryst. (2009). E65, m950 [doi:10.1107/S1600536809027779]

Bis(2-fluorobenzoato- κO)bis(pyridin-2-amine- κN^1)zinc(II)

J.-Q. Wang, Y.-W. Zhang and L. Cheng

Comment

Recently, coordination compounds of d_{10} monovalent ions of the coinage metals have attracted much attention because of their interesting photophysical properties. In order to construct many novel and interesting metal-organic frameworks carboxylates have been used widely. Moreover, supramolecular interactions such as hydrogen bonding and π - π stacking interactions have important effects on crystal engineering (Fyfe & Stoddart, 1997). In this paper, we present the synthesis and crystal structure of a new coordination compound of d_{10} monovalent ion $Zn(fa)_2(2-pa)_2$ (Hfa = 2-fluorobenzoic acid; 2-pa = pyridin-2-amine), (I).

The asymmetric unit of the title compound, contains a Zn^{II} cation, two fa and two 2-pa ligands (Fig. 1). The Zn^{II} ion in (I) is surrounded by two monodentate fa ligands with Zn—O coordinating distances 1.962 (2) and 1.976 (3) Å and two 2-pa ligands with distances involving pyridyl N atoms and Zn being 2.069 (2) and 2.056 (2) Å; the Zn^{II} displays a distorted tetrahedral geometry with angles around Zn in the range 101.17 (10) - 137.12 (10)°. The other O-atoms of the fa ligands are at significantly longer distances from the Zn^{II} ion (Zn1—O4 2.551 (3) and Zn1—O1 (2.781 (3)Å). The mononuclear complex (I) is joined into a two-dimensional layer by N—H \cdots O type hydrogen-bonds; details have been provided in Table 1. The layers are further constructed into a three-dimensional supramolecular network by rather weak C—H \cdots F type interactions (C5 \cdots F4^c 3.158 (2) and C20 \cdots F3^d 3.256 (3) Å; symmetry codes: ^c, 1 - x, 1/2 + y, 1/2 - z and ^d, -x, -y, 1 - z) and effective π - π stacking between the interlayer adjacent benzene rings and pyridyl rings with the centroid-centroid separation of 3.74 (3) Å. The F atoms of fa ligands in (I) are disordered over two sites, the 2- and 6-position of fa with equal site occupancy factors.

The crystal structures of a few Zn^{II} benzoate complexes have been reported by Darensbourg *et al.*, (2002).

Experimental

A mixture of 2-fluorobenzoic acid (0.0280 g, 0.2 mmol), pyridin-2-amine(0.0188 g, 0.2 mmol), $ZnSO_4 \cdot 7H_2O$ (0.0285 g, 0.1 mmol) and H_2O (8 ml) was heated in a 15-ml Teflon-lined autoclave at 393 K for 5 days, followed by slow cooling (5 ° h⁻¹) to room temperature. The resulting mixture was washed with water, and colorless block crystals were collected and dried in air [yield, 62% (32.4 mg) based on Zn^{II}].

Refinement

All the H atoms were positioned geometrically and refined using a riding model with C—H = 0.93 Å and N—H = 0.86 Å with $U_{iso}(H) = 1.2U_{iso}(C/N)$. The site occupancy factors of all F-atoms are given as 0.5 because of disordered position over two sites, the 2- and 6-position of fa.

Figures




Fig. 1. Molecular structure of the title compound with 15% thermal ellipsoids.




Fig. 2. The three-dimensional supramolecular network of the title compound.

Bis(2-fluorobenzoato- κO)bis(pyridin-2-amine- κN^1)zinc(II)

Crystal data

$[\text{Zn}(\text{C}_7\text{H}_4\text{FO}_2)_2(\text{C}_5\text{H}_6\text{N}_2)_2]$

$M_r = 531.83$

Monoclinic, $P2_1/c$

Hall symbol: $-P\ 2_1/c$

$a = 9.1259\ (7)\ \text{\AA}$

$b = 11.1020\ (9)\ \text{\AA}$

$c = 24.5707\ (17)\ \text{\AA}$

$\beta = 108.048\ (2)^\circ$

$V = 2366.9\ (3)\ \text{\AA}^3$

$Z = 4$

$F_{000} = 1088$

$D_x = 1.492\ \text{Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 785 reflections

$\theta = 2.4\text{--}28.0^\circ$

$\mu = 1.09\ \text{mm}^{-1}$

$T = 293\ \text{K}$

Block, colorless

$0.25 \times 0.20 \times 0.18\ \text{mm}$

Data collection

Bruker SMART APEX CCD
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 293\ \text{K}$

φ and ω scans

Absorption correction: multi-scan
(SADABS; Sheldrick, 2000)

$T_{\min} = 0.772$, $T_{\max} = 0.828$

13150 measured reflections

4645 independent reflections

3838 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.016$

$\theta_{\max} = 26.0^\circ$

$\theta_{\min} = 2.0^\circ$

$h = -10 \rightarrow 11$

$k = -13 \rightarrow 11$

$l = -30 \rightarrow 26$

Refinement

Refinement on F^2

Secondary atom site location: difference Fourier map

Least-squares matrix: full

$$R[F^2 > 2\sigma(F^2)] = 0.043$$

$$wR(F^2) = 0.131$$

$$S = 1.04$$

4645 reflections

334 parameters

Primary atom site location: structure-invariant direct methods

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.077P)^2 + 0.6445P]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} = 0.001$$

$$\Delta\rho_{\max} = 0.50 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.28 \text{ e } \text{\AA}^{-3}$$

Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Zn1	0.69105 (4)	-0.07190 (3)	0.378555 (11)	0.06568 (15)	
O1	0.9562 (3)	0.0314 (3)	0.36588 (9)	0.1045 (8)	
O2	0.7767 (3)	-0.0875 (2)	0.31487 (9)	0.0835 (6)	
O3	0.6376 (3)	0.06015 (18)	0.42292 (10)	0.0884 (7)	
O4	0.5834 (4)	0.1323 (3)	0.33750 (11)	0.1169 (9)	
N1	0.8367 (2)	-0.1820 (2)	0.43970 (9)	0.0634 (5)	
N2	0.8527 (3)	-0.0615 (2)	0.51811 (10)	0.0716 (6)	
H2A	0.7811	-0.0165	0.4967	0.086*	
H2B	0.8928	-0.0447	0.5538	0.086*	
N3	0.4880 (2)	-0.1672 (2)	0.35345 (9)	0.0638 (5)	
N4	0.5189 (4)	-0.2413 (3)	0.27024 (11)	0.0985 (9)	
H4B	0.6070	-0.2060	0.2786	0.118*	
H4C	0.4860	-0.2826	0.2392	0.118*	
C1	0.9007 (3)	-0.0286 (3)	0.32286 (11)	0.0658 (6)	
C2	0.9760 (3)	-0.0354 (3)	0.27689 (12)	0.0663 (7)	
C3	1.1117 (4)	0.0251 (3)	0.28201 (18)	0.0943 (10)	
H3A	1.1558	0.0718	0.3144	0.113*	0.50
F1	1.1788 (7)	0.0903 (6)	0.3380 (3)	0.155 (2)	0.50
C4	1.1843 (5)	0.0177 (5)	0.2398 (3)	0.1265 (18)	
H4A	1.2753	0.0594	0.2435	0.152*	
C5	1.1190 (7)	-0.0514 (6)	0.1936 (3)	0.136 (2)	
H5A	1.1673	-0.0575	0.1655	0.163*	

supplementary materials

C6	0.9885 (6)	-0.1112 (5)	0.18654 (18)	0.1150 (14)	
H6A	0.9461	-0.1573	0.1538	0.138*	
C7	0.9163 (4)	-0.1044 (3)	0.22799 (13)	0.0822 (9)	
H7A	0.8255	-0.1471	0.2231	0.099*	0.50
F2	0.7836 (4)	-0.1724 (4)	0.21912 (14)	0.0941 (11)	0.50
C8	0.5867 (3)	0.1415 (3)	0.38668 (13)	0.0768 (8)	
C9	0.5172 (3)	0.2517 (3)	0.40796 (12)	0.0707 (7)	
C10	0.4409 (5)	0.2449 (4)	0.44744 (17)	0.1011 (11)	
H10A	0.4357	0.1708	0.4644	0.121*	0.50
F3	0.4654 (14)	0.1351 (9)	0.4726 (4)	0.224 (4)	0.50
C11	0.3718 (6)	0.3414 (5)	0.4633 (2)	0.1287 (17)	
H11A	0.3202	0.3320	0.4902	0.154*	
C12	0.3779 (5)	0.4498 (5)	0.4402 (2)	0.1193 (15)	
H12A	0.3288	0.5154	0.4505	0.143*	
C13	0.4567 (5)	0.4637 (4)	0.4015 (2)	0.1106 (13)	
H13A	0.4655	0.5390	0.3862	0.133*	
C14	0.5223 (4)	0.3637 (3)	0.38576 (17)	0.0946 (10)	
H14A	0.5730	0.3728	0.3585	0.114*	0.50
F4	0.6067 (6)	0.3817 (4)	0.3530 (2)	0.1135 (15)	0.50
C15	0.8819 (4)	-0.2827 (3)	0.41838 (13)	0.0803 (8)	
H15A	0.8346	-0.3002	0.3799	0.096*	
C16	0.9906 (4)	-0.3591 (3)	0.44900 (17)	0.0921 (9)	
H16A	1.0179	-0.4272	0.4323	0.111*	
C17	1.0612 (4)	-0.3324 (3)	0.50709 (16)	0.0888 (9)	
H17A	1.1377	-0.3827	0.5296	0.107*	
C18	1.0180 (3)	-0.2340 (3)	0.53020 (13)	0.0768 (8)	
H18A	1.0644	-0.2162	0.5687	0.092*	
C19	0.9016 (3)	-0.1578 (2)	0.49576 (10)	0.0606 (6)	
C20	0.4016 (4)	-0.1603 (3)	0.38926 (13)	0.0765 (7)	
H20A	0.4384	-0.1141	0.4223	0.092*	
C21	0.2633 (4)	-0.2177 (3)	0.37954 (17)	0.0867 (9)	
H21A	0.2080	-0.2120	0.4055	0.104*	
C22	0.2084 (4)	-0.2842 (3)	0.32992 (18)	0.0908 (10)	
H22A	0.1143	-0.3238	0.3218	0.109*	
C23	0.2914 (4)	-0.2918 (3)	0.29318 (14)	0.0833 (9)	
H23A	0.2541	-0.3366	0.2598	0.100*	
C24	0.4341 (3)	-0.2322 (2)	0.30510 (12)	0.0692 (7)	

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Zn1	0.0722 (2)	0.0778 (2)	0.04154 (19)	-0.01443 (14)	0.00970 (14)	-0.00021 (12)
O1	0.1209 (19)	0.1082 (17)	0.0622 (13)	-0.0016 (15)	-0.0042 (12)	-0.0207 (12)
O2	0.0868 (14)	0.1085 (16)	0.0585 (11)	-0.0235 (12)	0.0275 (10)	-0.0056 (10)
O3	0.1029 (16)	0.0703 (13)	0.0712 (13)	-0.0056 (10)	-0.0036 (12)	0.0099 (9)
O4	0.143 (2)	0.129 (2)	0.0806 (17)	0.0444 (19)	0.0377 (16)	0.0251 (16)
N1	0.0669 (12)	0.0705 (13)	0.0498 (11)	-0.0109 (10)	0.0136 (9)	0.0004 (9)
N2	0.0803 (15)	0.0780 (15)	0.0471 (11)	-0.0004 (11)	0.0059 (11)	-0.0008 (10)

N3	0.0636 (12)	0.0679 (13)	0.0518 (11)	-0.0068 (10)	0.0061 (9)	0.0010 (9)
N4	0.0970 (19)	0.125 (2)	0.0697 (16)	-0.0311 (17)	0.0199 (14)	-0.0357 (16)
C1	0.0700 (16)	0.0686 (15)	0.0487 (13)	0.0060 (13)	0.0035 (11)	0.0076 (12)
C2	0.0594 (14)	0.0699 (15)	0.0645 (15)	0.0093 (12)	0.0116 (12)	0.0204 (13)
C3	0.0734 (19)	0.088 (2)	0.115 (3)	0.0021 (17)	0.0206 (19)	0.031 (2)
F1	0.119 (4)	0.137 (5)	0.195 (7)	-0.042 (3)	0.026 (4)	-0.008 (4)
C4	0.077 (2)	0.139 (4)	0.174 (5)	0.014 (3)	0.054 (3)	0.070 (4)
C5	0.120 (4)	0.180 (6)	0.133 (4)	0.058 (4)	0.074 (4)	0.066 (4)
C6	0.110 (3)	0.167 (4)	0.078 (2)	0.039 (3)	0.043 (2)	0.016 (3)
C7	0.0791 (19)	0.108 (2)	0.0595 (16)	0.0155 (17)	0.0223 (14)	0.0096 (16)
F2	0.085 (2)	0.134 (3)	0.0648 (19)	-0.030 (2)	0.0249 (17)	-0.040 (2)
C8	0.0685 (16)	0.0784 (19)	0.0685 (17)	-0.0052 (14)	-0.0005 (13)	0.0200 (15)
C9	0.0593 (14)	0.0695 (17)	0.0711 (16)	-0.0056 (12)	0.0024 (12)	0.0139 (13)
C10	0.131 (3)	0.086 (2)	0.091 (2)	-0.030 (2)	0.042 (2)	0.0014 (19)
F3	0.339 (13)	0.194 (8)	0.187 (7)	-0.059 (8)	0.150 (8)	-0.001 (6)
C11	0.136 (4)	0.135 (4)	0.136 (4)	-0.039 (3)	0.073 (3)	-0.033 (3)
C12	0.094 (3)	0.118 (4)	0.144 (4)	0.003 (2)	0.035 (3)	-0.021 (3)
C13	0.113 (3)	0.076 (2)	0.129 (3)	0.018 (2)	0.017 (3)	0.025 (2)
C14	0.089 (2)	0.089 (2)	0.102 (2)	0.0056 (18)	0.0255 (19)	0.029 (2)
F4	0.172 (4)	0.090 (3)	0.121 (3)	-0.015 (3)	0.106 (3)	0.015 (2)
C15	0.093 (2)	0.081 (2)	0.0661 (17)	-0.0059 (16)	0.0231 (15)	-0.0037 (15)
C16	0.092 (2)	0.080 (2)	0.106 (3)	0.0051 (18)	0.034 (2)	-0.0042 (19)
C17	0.0722 (18)	0.080 (2)	0.104 (2)	0.0020 (15)	0.0115 (17)	0.0141 (19)
C18	0.0678 (16)	0.0799 (19)	0.0685 (16)	-0.0092 (14)	0.0003 (13)	0.0102 (14)
C19	0.0559 (13)	0.0696 (15)	0.0516 (12)	-0.0148 (11)	0.0097 (10)	0.0071 (11)
C20	0.0832 (19)	0.0736 (18)	0.0721 (17)	-0.0086 (14)	0.0235 (15)	-0.0016 (14)
C21	0.0767 (19)	0.079 (2)	0.109 (3)	-0.0037 (16)	0.0346 (18)	0.0059 (18)
C22	0.0663 (17)	0.0722 (19)	0.122 (3)	-0.0099 (14)	0.0109 (19)	0.0084 (19)
C23	0.0765 (18)	0.0679 (17)	0.085 (2)	-0.0096 (14)	-0.0043 (16)	-0.0052 (15)
C24	0.0690 (15)	0.0647 (15)	0.0618 (15)	-0.0022 (12)	0.0025 (12)	-0.0003 (12)

Geometric parameters (Å, °)

Zn1—O2	1.962 (2)	C7—F2	1.386 (5)
Zn1—O3	1.976 (3)	C7—H7A	0.9300
Zn1—N3	2.056 (2)	C8—C9	1.542 (5)
Zn1—N1	2.069 (2)	C9—C10	1.360 (5)
Zn1—O4	2.551 (3)	C9—C14	1.365 (4)
Zn1—O1	2.781 (3)	C10—F3	1.353 (10)
O1—C1	1.219 (3)	C10—C11	1.359 (6)
O2—C1	1.268 (3)	C10—H10A	0.9300
O3—C8	1.252 (3)	C11—C12	1.339 (7)
O4—C8	1.204 (4)	C11—H11A	0.9300
N1—C19	1.347 (3)	C12—C13	1.368 (7)
N1—C15	1.352 (4)	C12—H12A	0.9300
N2—C19	1.340 (4)	C13—C14	1.372 (6)
N2—H2A	0.8599	C13—H13A	0.9300
N2—H2B	0.8603	C14—F4	1.291 (5)
N3—C24	1.346 (3)	C14—H14A	0.9300

supplementary materials

N3—C20	1.353 (4)	C15—C16	1.344 (5)
N4—C24	1.324 (4)	C15—H15A	0.9300
N4—H4B	0.8601	C16—C17	1.404 (5)
N4—H4C	0.8601	C16—H16A	0.9300
C1—C2	1.495 (4)	C17—C18	1.345 (5)
C2—C3	1.380 (4)	C17—H17A	0.9300
C2—C7	1.386 (5)	C18—C19	1.416 (4)
C3—C4	1.395 (6)	C18—H18A	0.9300
C3—F1	1.506 (8)	C20—C21	1.367 (4)
C3—H3A	0.9300	C20—H20A	0.9300
C4—C5	1.347 (8)	C21—C22	1.380 (5)
C4—H4A	0.9300	C21—H21A	0.9300
C5—C6	1.327 (8)	C22—C23	1.349 (5)
C5—H5A	0.9300	C22—H22A	0.9300
C6—C7	1.375 (5)	C23—C24	1.408 (4)
C6—H6A	0.9300	C23—H23A	0.9300
O2—Zn1—O3	137.12 (10)	C10—C9—C14	115.1 (3)
O2—Zn1—N3	105.09 (9)	C10—C9—C8	123.6 (3)
O3—Zn1—N3	101.17 (10)	C14—C9—C8	121.3 (3)
O2—Zn1—N1	101.66 (10)	F3—C10—C11	127.2 (5)
O3—Zn1—N1	104.50 (9)	F3—C10—C9	109.1 (5)
N3—Zn1—N1	103.33 (8)	C11—C10—C9	123.1 (4)
O2—Zn1—O4	87.86 (9)	C11—C10—H10A	118.5
O3—Zn1—O4	55.12 (9)	C9—C10—H10A	118.5
N3—Zn1—O4	97.91 (10)	C12—C11—C10	120.3 (4)
N1—Zn1—O4	153.41 (10)	C12—C11—H11A	119.9
C1—O2—Zn1	112.85 (18)	C10—C11—H11A	119.9
C8—O3—Zn1	104.1 (2)	C11—C12—C13	119.7 (5)
C19—N1—C15	118.1 (2)	C11—C12—H12A	120.2
C19—N1—Zn1	127.21 (19)	C13—C12—H12A	120.2
C15—N1—Zn1	114.31 (18)	C12—C13—C14	118.3 (4)
C19—N2—H2A	120.0	C12—C13—H13A	120.8
C19—N2—H2B	120.0	C14—C13—H13A	120.8
H2A—N2—H2B	120.0	F4—C14—C9	119.4 (4)
C24—N3—C20	118.5 (2)	F4—C14—C13	116.6 (4)
C24—N3—Zn1	126.11 (19)	C9—C14—C13	123.5 (4)
C20—N3—Zn1	115.36 (18)	C9—C14—H14A	118.2
C24—N4—H4B	120.0	C13—C14—H14A	118.2
C24—N4—H4C	120.0	C16—C15—N1	124.5 (3)
H4B—N4—H4C	120.0	C16—C15—H15A	117.8
O1—C1—O2	121.8 (3)	N1—C15—H15A	117.8
O1—C1—C2	121.2 (3)	C15—C16—C17	117.5 (3)
O2—C1—C2	117.0 (2)	C15—C16—H16A	121.2
C3—C2—C7	116.6 (3)	C17—C16—H16A	121.2
C3—C2—C1	121.1 (3)	C18—C17—C16	119.9 (3)
C7—C2—C1	122.3 (3)	C18—C17—H17A	120.1
C2—C3—C4	121.6 (4)	C16—C17—H17A	120.1
C2—C3—F1	114.6 (4)	C17—C18—C19	119.8 (3)
C4—C3—F1	123.6 (5)	C17—C18—H18A	120.1

C2—C3—H3A	119.2	C19—C18—H18A	120.1
C4—C3—H3A	119.2	N2—C19—N1	118.8 (2)
C5—C4—C3	118.2 (4)	N2—C19—C18	121.0 (2)
C5—C4—H4A	120.9	N1—C19—C18	120.2 (3)
C3—C4—H4A	120.9	N3—C20—C21	123.5 (3)
C6—C5—C4	122.7 (5)	N3—C20—H20A	118.2
C6—C5—H5A	118.7	C21—C20—H20A	118.2
C4—C5—H5A	118.7	C20—C21—C22	117.8 (3)
C5—C6—C7	119.5 (5)	C20—C21—H21A	121.1
C5—C6—H6A	120.3	C22—C21—H21A	121.1
C7—C6—H6A	120.3	C23—C22—C21	120.0 (3)
C6—C7—F2	116.7 (4)	C23—C22—H22A	120.0
C6—C7—C2	121.5 (4)	C21—C22—H22A	120.0
F2—C7—C2	121.8 (3)	C22—C23—C24	120.4 (3)
C6—C7—H7A	119.3	C22—C23—H23A	119.8
C2—C7—H7A	119.3	C24—C23—H23A	119.8
O4—C8—O3	122.8 (3)	N4—C24—N3	119.0 (3)
O4—C8—C9	121.5 (3)	N4—C24—C23	121.2 (3)
O3—C8—C9	115.6 (3)	N3—C24—C23	119.8 (3)
O3—Zn1—O2—C1	51.4 (3)	C3—C2—C7—F2	177.7 (3)
N3—Zn1—O2—C1	177.2 (2)	C1—C2—C7—F2	-0.7 (5)
N1—Zn1—O2—C1	-75.3 (2)	Zn1—O4—C8—O3	2.2 (3)
O4—Zn1—O2—C1	79.6 (2)	Zn1—O4—C8—C9	-173.4 (3)
O2—Zn1—O3—C8	36.5 (3)	Zn1—O3—C8—O4	-2.9 (4)
N3—Zn1—O3—C8	-90.5 (2)	Zn1—O3—C8—C9	173.00 (19)
N1—Zn1—O3—C8	162.39 (19)	O4—C8—C9—C10	142.4 (4)
O4—Zn1—O3—C8	1.38 (19)	O3—C8—C9—C10	-33.5 (4)
O2—Zn1—O4—C8	-158.3 (2)	O4—C8—C9—C14	-34.3 (5)
O3—Zn1—O4—C8	-1.4 (2)	O3—C8—C9—C14	149.8 (3)
N3—Zn1—O4—C8	96.7 (2)	C14—C9—C10—F3	-170.2 (6)
N1—Zn1—O4—C8	-46.2 (3)	C8—C9—C10—F3	12.9 (7)
O2—Zn1—N1—C19	135.2 (2)	C14—C9—C10—C11	1.1 (6)
O3—Zn1—N1—C19	-10.5 (2)	C8—C9—C10—C11	-175.8 (4)
N3—Zn1—N1—C19	-116.0 (2)	F3—C10—C11—C12	169.1 (8)
O4—Zn1—N1—C19	26.1 (3)	C9—C10—C11—C12	-0.6 (7)
O2—Zn1—N1—C15	-37.3 (2)	C10—C11—C12—C13	-1.3 (8)
O3—Zn1—N1—C15	177.0 (2)	C11—C12—C13—C14	2.6 (7)
N3—Zn1—N1—C15	71.5 (2)	C10—C9—C14—F4	172.2 (4)
O4—Zn1—N1—C15	-146.4 (2)	C8—C9—C14—F4	-10.8 (6)
O2—Zn1—N3—C24	4.1 (2)	C10—C9—C14—C13	0.2 (5)
O3—Zn1—N3—C24	149.9 (2)	C8—C9—C14—C13	177.2 (3)
N1—Zn1—N3—C24	-102.1 (2)	C12—C13—C14—F4	-174.2 (5)
O4—Zn1—N3—C24	94.0 (2)	C12—C13—C14—C9	-2.0 (6)
O2—Zn1—N3—C20	-174.6 (2)	C19—N1—C15—C16	-1.8 (4)
O3—Zn1—N3—C20	-28.8 (2)	Zn1—N1—C15—C16	171.4 (3)
N1—Zn1—N3—C20	79.2 (2)	N1—C15—C16—C17	0.2 (5)
O4—Zn1—N3—C20	-84.7 (2)	C15—C16—C17—C18	0.7 (5)
Zn1—O2—C1—O1	-0.8 (4)	C16—C17—C18—C19	-0.1 (5)
Zn1—O2—C1—C2	179.51 (18)	C15—N1—C19—N2	-176.8 (2)

supplementary materials

O1—C1—C2—C3	0.7 (4)	Zn1—N1—C19—N2	10.9 (3)
O2—C1—C2—C3	-179.6 (3)	C15—N1—C19—C18	2.4 (4)
O1—C1—C2—C7	179.1 (3)	Zn1—N1—C19—C18	-169.80 (19)
O2—C1—C2—C7	-1.3 (4)	C17—C18—C19—N2	177.7 (3)
C7—C2—C3—C4	0.5 (5)	C17—C18—C19—N1	-1.6 (4)
C1—C2—C3—C4	178.9 (3)	C24—N3—C20—C21	1.2 (4)
C7—C2—C3—F1	-175.1 (4)	Zn1—N3—C20—C21	180.0 (3)
C1—C2—C3—F1	3.3 (5)	N3—C20—C21—C22	-1.2 (5)
C2—C3—C4—C5	-0.5 (6)	C20—C21—C22—C23	0.6 (5)
F1—C3—C4—C5	174.7 (5)	C21—C22—C23—C24	0.1 (5)
C3—C4—C5—C6	0.7 (8)	C20—N3—C24—N4	-178.8 (3)
C4—C5—C6—C7	-0.8 (8)	Zn1—N3—C24—N4	2.5 (4)
C5—C6—C7—F2	-177.7 (4)	C20—N3—C24—C23	-0.5 (4)
C5—C6—C7—C2	0.7 (6)	Zn1—N3—C24—C23	-179.2 (2)
C3—C2—C7—C6	-0.5 (5)	C22—C23—C24—N4	178.2 (3)
C1—C2—C7—C6	-178.9 (3)	C22—C23—C24—N3	-0.1 (5)

Hydrogen-bond geometry (\AA , $^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
N2—H2A \cdots O3	0.86	2.06	2.881 (3)	159
N2—H2B \cdots O1 ⁱ	0.86	2.03	2.864 (3)	162
N4—H4B \cdots O2	0.86	2.02	2.838 (4)	159
N4—H4B \cdots F2	0.86	2.52	3.143 (5)	130
N4—H4C \cdots O4 ⁱⁱ	0.86	2.03	2.884 (4)	175
C5—H5A \cdots F4 ⁱⁱⁱ	0.93	2.34	3.147 (6)	144
C20—H20A \cdots F3 ^{iv}	0.93	2.47	3.244 (10)	141

Symmetry codes: (i) $-x+2, -y, -z+1$; (ii) $-x+1, y-1/2, -z+1/2$; (iii) $-x+2, y-1/2, -z+1/2$; (iv) $-x+1, -y, -z+1$.

Fig. 1




Fig. 2

