

Bis(acetone 4-phenylthiosemicarbazoneato- $\kappa^2 N^1, S$)zinc(II)

Kong Wai Tan,^a Chew Hee Ng,^b Mohd Jamil Maah^{a*} and Seik Weng Ng^a

^aDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia, and ^bFaculty of Engineering and Science, Universiti Tunku Abdul Rahman, 53300 Kuala Lumpur, Malaysia

Correspondence e-mail: mjamil@um.edu.my


Received 16 July 2009; accepted 17 July 2009

Key indicators: single-crystal X-ray study; $T = 153\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.005\text{ \AA}$; R factor = 0.039; wR factor = 0.103; data-to-parameter ratio = 18.4.

The Zn^{II} atom in the title compound, $[\text{Zn}(\text{C}_{10}\text{H}_{12}\text{N}_3\text{S})_2]$, is N, S -chelated by the deprotonated Schiff base in a tetrahedral environment. The metal atom lies on a twofold rotation axis that relates one anion to the other. The amino H atom forms an intermolecular $\text{N}-\text{H}\cdots\pi$ interaction to an aromatic ring.

Related literature

For the two modifications of acetone 4-phenylthiosemicarbazone, see: Jian *et al.* (2005); Venkatraman *et al.* (2005).



Experimental

Crystal data

$[\text{Zn}(\text{C}_{10}\text{H}_{12}\text{N}_3\text{S})_2]$
 $M_r = 477.94$
Monoclinic, $C2/c$

$a = 23.5203(5)\text{ \AA}$
 $b = 7.2938(2)\text{ \AA}$
 $c = 15.1134(5)\text{ \AA}$

$\beta = 122.761(2)^\circ$
 $V = 2180.3(1)\text{ \AA}^3$
 $Z = 4$
Mo $K\alpha$ radiation

$\mu = 1.34\text{ mm}^{-1}$
 $T = 153\text{ K}$
 $0.30 \times 0.20 \times 0.05\text{ mm}$

Data collection

Bruker SMART APEX diffractometer
Absorption correction: multi-scan (*SADABS*; Sheldrick, 1996)
 $T_{\min} = 0.690$, $T_{\max} = 0.936$

6983 measured reflections
2488 independent reflections
1896 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.031$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.039$
 $wR(F^2) = 0.103$
 $S = 1.10$
2488 reflections

135 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.40\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.45\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N3}-\text{H3}\cdots Cg^{\dagger}$	0.86	2.86	3.671 (3)	157

Symmetry code: (i) $-x + \frac{3}{2}, y - \frac{1}{2}, -z + \frac{1}{2}$. Cg is the centroid of the aromatic ring.

Data collection: *APEX2* (Bruker, 2008); cell refinement: *SAINT* (Bruker, 2008); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2009).

We thank the University of Malaya (PJP FS316/2008 C) and MOSTI (ESc 02-02-11-SF0033) for supporting this study; KWT thanks the Ministry of Higher Education for a SLAI scholarship.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT5011).

References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
- Bruker (2008). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Jian, F.-F., Bai, Z.-S., Xiao, H.-L. & Li, K. (2005). *Acta Cryst. E* **61**, o653–o654.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Venkatraman, R., Swesi, A. T. & Yamin, B. M. (2005). *Acta Cryst. E* **61**, o3914–o3916.
- Westrip, S. P. (2009). *publCIF*. In preparation.

supplementary materials

Acta Cryst. (2009). E65, m969 [doi:10.1107/S1600536809028244]

Bis(acetone 4-phenylthiosemicarbazonoato- $\kappa^2 N^1, S$)zinc(II)

K. W. Tan, C. H. Ng, M. J. Maah and S. W. Ng

Experimental

Zinc acetate monohydrate (0.22 g, 1 mmol) and 2,4-dihydroxybenzaldehyde 4-phenylthiosemicarbazone (0.29 g, 1 mmol) were heated in ethanol (50 ml) to form [1-(4-hydroxy-2-oxidobenzylidene)-4-phenylthiosemicarbazonoato]zinc. The product was recrystallized from acetone, cleaved part of the anion.

Refinement

Carbon-bound H-atoms were placed in calculated positions (C—H 0.95–0.98 Å) and were included in the refinement in the riding model approximation, with $U(H)$ set to 1.2–1.5 $U_{\text{eq}}(\text{C})$. The amino H-atom was similarly treated (N—H 0.88 Å).

Figures




Fig. 1. Anisotropic displacement ellipsoid (Barbour, 2001) plot of $\text{Zn}(\text{C}_{10}\text{H}_{12}\text{N}_3\text{S})_2$ at the 70% probability level. Hydrogen atoms are drawn as spheres of arbitrary radii.

Bis(acetone 4-phenylthiosemicarbazonoato- $\kappa^2 N^1, S$)zinc(II)

Crystal data

$[\text{Zn}(\text{C}_{10}\text{H}_{12}\text{N}_3\text{S})_2]$	$F_{000} = 992$
$M_r = 477.94$	$D_x = 1.456 \text{ Mg m}^{-3}$
Monoclinic, $C2/c$	$\text{Mo } K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -C 2yc	Cell parameters from 2390 reflections
$a = 23.5203 (5) \text{ \AA}$	$\theta = 3.8\text{--}28.0^\circ$
$b = 7.2938 (2) \text{ \AA}$	$\mu = 1.34 \text{ mm}^{-1}$
$c = 15.1134 (5) \text{ \AA}$	$T = 153 \text{ K}$
$\beta = 122.761 (2)^\circ$	Wedge, brown
$V = 2180.3 (1) \text{ \AA}^3$	$0.30 \times 0.20 \times 0.05 \text{ mm}$
$Z = 4$	

Data collection

Bruker SMART APEX diffractometer	2488 independent reflections
Radiation source: fine-focus sealed tube	1896 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.031$

supplementary materials

$T = 153\text{ K}$	$\theta_{\max} = 27.5^\circ$
ω scans	$\theta_{\min} = 2.1^\circ$
Absorption correction: Multi-scan (SADABS; Sheldrick, 1996)	$h = -30 \rightarrow 30$
$T_{\min} = 0.690$, $T_{\max} = 0.936$	$k = -9 \rightarrow 9$
6983 measured reflections	$l = -19 \rightarrow 19$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.039$	H-atom parameters constrained
$wR(F^2) = 0.103$	$w = 1/[\sigma^2(F_o^2) + (0.0387P)^2 + 4.48P]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.10$	$(\Delta/\sigma)_{\max} = 0.001$
2488 reflections	$\Delta\rho_{\max} = 0.40\text{ e \AA}^{-3}$
135 parameters	$\Delta\rho_{\min} = -0.45\text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Zn1	0.5000	0.30141 (7)	0.2500	0.03064 (17)
S1	0.55393 (4)	0.14771 (13)	0.18550 (8)	0.0501 (3)
N1	0.58756 (10)	0.4440 (3)	0.34189 (19)	0.0245 (5)
N2	0.64335 (10)	0.3888 (3)	0.33788 (19)	0.0253 (5)
N3	0.68276 (11)	0.1943 (4)	0.2634 (2)	0.0336 (6)
H3	0.6712	0.1130	0.2154	0.040*
C1	0.75216 (12)	0.2404 (4)	0.3216 (2)	0.0255 (6)
C2	0.78857 (13)	0.1872 (4)	0.2775 (2)	0.0296 (6)
H2	0.7662	0.1268	0.2113	0.035*
C3	0.85719 (14)	0.2225 (4)	0.3300 (3)	0.0348 (7)
H3A	0.8816	0.1866	0.2993	0.042*
C4	0.89046 (13)	0.3090 (4)	0.4263 (3)	0.0332 (7)
H4	0.9376	0.3330	0.4620	0.040*
C5	0.85427 (13)	0.3606 (4)	0.4703 (2)	0.0289 (6)
H5	0.8770	0.4194	0.5370	0.035*
C6	0.78518 (13)	0.3275 (4)	0.4181 (2)	0.0283 (6)
H6	0.7608	0.3646	0.4486	0.034*
C7	0.63093 (13)	0.2588 (4)	0.2715 (2)	0.0275 (6)
C8	0.59964 (13)	0.5716 (4)	0.4081 (2)	0.0302 (7)
C9	0.66790 (15)	0.6551 (5)	0.4773 (3)	0.0456 (9)
H9A	0.7007	0.5590	0.5196	0.068*
H9B	0.6663	0.7453	0.5241	0.068*
H9C	0.6815	0.7159	0.4338	0.068*
C10	0.54313 (16)	0.6384 (5)	0.4182 (3)	0.0451 (9)

H10A	0.5039	0.5575	0.3784	0.068*
H10B	0.5307	0.7633	0.3903	0.068*
H10C	0.5578	0.6380	0.4925	0.068*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Zn1	0.0144 (2)	0.0296 (3)	0.0436 (3)	0.000	0.0129 (2)	0.000
S1	0.0171 (3)	0.0539 (6)	0.0665 (7)	-0.0064 (3)	0.0143 (4)	-0.0345 (5)
N1	0.0156 (9)	0.0257 (12)	0.0320 (14)	0.0020 (9)	0.0128 (10)	0.0034 (11)
N2	0.0148 (10)	0.0287 (12)	0.0321 (14)	0.0015 (9)	0.0124 (10)	-0.0009 (11)
N3	0.0196 (11)	0.0373 (14)	0.0408 (16)	-0.0017 (10)	0.0143 (11)	-0.0158 (13)
C1	0.0168 (11)	0.0235 (14)	0.0328 (16)	0.0030 (10)	0.0113 (12)	0.0006 (12)
C2	0.0245 (13)	0.0303 (15)	0.0332 (17)	0.0041 (12)	0.0152 (13)	-0.0016 (14)
C3	0.0241 (14)	0.0429 (18)	0.0432 (19)	0.0065 (13)	0.0221 (14)	0.0041 (16)
C4	0.0189 (12)	0.0373 (17)	0.0398 (18)	0.0023 (12)	0.0136 (13)	0.0086 (15)
C5	0.0216 (13)	0.0264 (14)	0.0310 (17)	0.0007 (11)	0.0092 (12)	0.0040 (13)
C6	0.0219 (13)	0.0307 (16)	0.0328 (17)	0.0034 (11)	0.0152 (13)	0.0004 (13)
C7	0.0171 (12)	0.0299 (15)	0.0328 (17)	-0.0001 (11)	0.0116 (12)	-0.0030 (13)
C8	0.0228 (13)	0.0258 (15)	0.0408 (18)	0.0015 (11)	0.0165 (13)	-0.0037 (14)
C9	0.0297 (15)	0.0361 (19)	0.063 (2)	-0.0023 (13)	0.0197 (17)	-0.0213 (17)
C10	0.0321 (16)	0.046 (2)	0.062 (2)	0.0046 (14)	0.0284 (17)	-0.0102 (18)

Geometric parameters (\AA , $^\circ$)

Zn1—N1	2.039 (2)	C3—C4	1.376 (5)
Zn1—N1 ⁱ	2.039 (2)	C3—H3A	0.9500
Zn1—S1	2.2702 (8)	C4—C5	1.386 (4)
Zn1—S1 ⁱ	2.2702 (8)	C4—H4	0.9500
S1—C7	1.754 (3)	C5—C6	1.390 (4)
N1—C8	1.281 (4)	C5—H5	0.9500
N1—N2	1.405 (3)	C6—H6	0.9500
N2—C7	1.294 (4)	C8—C9	1.492 (4)
N3—C7	1.372 (3)	C8—C10	1.499 (4)
N3—C1	1.413 (3)	C9—H9A	0.9800
N3—H3	0.8600	C9—H9B	0.9800
C1—C6	1.381 (4)	C9—H9C	0.9800
C1—C2	1.395 (4)	C10—H10A	0.9800
C2—C3	1.384 (4)	C10—H10B	0.9800
C2—H2	0.9500	C10—H10C	0.9800
N1—Zn1—N1 ⁱ	118.66 (13)	C5—C4—H4	120.5
N1—Zn1—S1	87.20 (6)	C4—C5—C6	120.9 (3)
N1 ⁱ —Zn1—S1	123.54 (7)	C4—C5—H5	119.6
N1—Zn1—S1 ⁱ	123.54 (7)	C6—C5—H5	119.6
N1 ⁱ —Zn1—S1 ⁱ	87.20 (6)	C1—C6—C5	119.7 (3)
S1—Zn1—S1 ⁱ	120.82 (6)	C1—C6—H6	120.1
C7—S1—Zn1	92.67 (9)	C5—C6—H6	120.1

supplementary materials

C8—N1—N2	115.1 (2)	N2—C7—N3	118.8 (2)
C8—N1—Zn1	128.17 (17)	N2—C7—S1	128.47 (19)
N2—N1—Zn1	116.57 (17)	N3—C7—S1	112.8 (2)
C7—N2—N1	114.8 (2)	N1—C8—C9	123.0 (2)
C7—N3—C1	130.2 (3)	N1—C8—C10	118.8 (3)
C7—N3—H3	114.9	C9—C8—C10	118.2 (3)
C1—N3—H3	114.9	C8—C9—H9A	109.5
C6—C1—C2	119.5 (2)	C8—C9—H9B	109.5
C6—C1—N3	124.5 (2)	H9A—C9—H9B	109.5
C2—C1—N3	116.0 (3)	C8—C9—H9C	109.5
C3—C2—C1	120.1 (3)	H9A—C9—H9C	109.5
C3—C2—H2	120.0	H9B—C9—H9C	109.5
C1—C2—H2	120.0	C8—C10—H10A	109.5
C4—C3—C2	120.8 (3)	C8—C10—H10B	109.5
C4—C3—H3A	119.6	H10A—C10—H10B	109.5
C2—C3—H3A	119.6	C8—C10—H10C	109.5
C3—C4—C5	119.1 (2)	H10A—C10—H10C	109.5
C3—C4—H4	120.5	H10B—C10—H10C	109.5
N1—Zn1—S1—C7	4.02 (12)	C2—C3—C4—C5	0.0 (5)
N1 ⁱ —Zn1—S1—C7	126.96 (13)	C3—C4—C5—C6	-0.6 (4)
S1 ⁱ —Zn1—S1—C7	-123.93 (11)	C2—C1—C6—C5	-0.3 (4)
N1 ⁱ —Zn1—N1—C8	52.6 (2)	N3—C1—C6—C5	177.8 (3)
S1—Zn1—N1—C8	179.7 (3)	C4—C5—C6—C1	0.7 (4)
S1 ⁱ —Zn1—N1—C8	-54.6 (3)	N1—N2—C7—N3	-179.0 (2)
N1 ⁱ —Zn1—N1—N2	-132.0 (2)	N1—N2—C7—S1	1.3 (4)
S1—Zn1—N1—N2	-4.85 (18)	C1—N3—C7—N2	3.8 (5)
S1 ⁱ —Zn1—N1—N2	120.81 (17)	C1—N3—C7—S1	-176.5 (3)
C8—N1—N2—C7	179.4 (3)	Zn1—S1—C7—N2	-4.3 (3)
Zn1—N1—N2—C7	3.4 (3)	Zn1—S1—C7—N3	176.0 (2)
C7—N3—C1—C6	18.0 (5)	N2—N1—C8—C9	-0.7 (4)
C7—N3—C1—C2	-163.9 (3)	Zn1—N1—C8—C9	174.8 (2)
C6—C1—C2—C3	-0.3 (4)	N2—N1—C8—C10	-179.7 (3)
N3—C1—C2—C3	-178.5 (3)	Zn1—N1—C8—C10	-4.1 (4)
C1—C2—C3—C4	0.4 (5)		

Symmetry codes: (i) $-x+1, y, -z+1/2$.

Hydrogen-bond geometry (\AA , $^\circ$)

$D\text{—H}\cdots A$	$D\text{—H}$	$H\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
N3—H3 \cdots Cg ⁱⁱ	0.86	2.86	3.671 (3)	157

Symmetry codes: (ii) $-x+3/2, y-1/2, -z+1/2$.

Fig. 1

