

1,4-Diazoniabicyclo[2.2.2]octane tetra-bromidocadmate(II) monohydrate

Kong Mun Lo and Seik Weng Ng*

Department of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia
Correspondence e-mail: seikweng@um.edu.my


Received 16 July 2009; accepted 17 July 2009

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.023$ Å; R factor = 0.057; wR factor = 0.187; data-to-parameter ratio = 19.1.

The metal atom in the anion of the title salt, $(\text{C}_6\text{H}_{14}\text{N}_2)\text{[CdBr}_4\text{]}\cdot\text{H}_2\text{O}$, shows a slightly distorted tetrahedral coordination. The water molecule is involved in three hydrogen bonds, *viz.* one $\text{N}-\text{H}\cdots\text{O}$ and two $\text{O}-\text{H}\cdots\text{Br}$, and an $\text{N}-\text{H}\cdots\text{Br}$ interaction consolidates the three-dimensional network.

Related literature

For other ammonium tetrabromidocadmates, see: Al-Far & Ali (2008); Battaglia *et al.* (1991); Chen *et al.* (2006); Geselle & Fuess (1994); Hatano *et al.* (2008); Ishihara *et al.* (2002, 2006); Ravikumar *et al.* (1995); Waskowska (1994); Zhang & Fang (2005).



Experimental

Crystal data

$(\text{C}_6\text{H}_{14}\text{N}_2)\text{[CdBr}_4\text{]}\cdot\text{H}_2\text{O}$
 $M_r = 564.25$
Orthorhombic, $P2_12_12_1$
 $a = 8.6323$ (1) Å
 $b = 11.8736$ (2) Å
 $c = 13.5619$ (2) Å

$V = 1390.05$ (4) Å³
 $Z = 4$
Mo $K\alpha$ radiation
 $\mu = 13.04$ mm⁻¹
 $T = 296$ K
 $0.30 \times 0.15 \times 0.05$ mm

Data collection

Bruker SMART APEX diffractometer
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)
 $T_{\min} = 0.111$, $T_{\max} = 0.562$
(expected range = 0.103–0.521)

10779 measured reflections
2451 independent reflections
2267 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.034$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.057$
 $wR(F^2) = 0.187$
 $S = 1.36$
2451 reflections
128 parameters
H-atom parameters constrained

$\Delta\rho_{\text{max}} = 2.21$ e Å⁻³
 $\Delta\rho_{\text{min}} = -2.09$ e Å⁻³
Absolute structure: Flack (1983),
1021 Friedel pairs
Flack parameter: 0.14 (3)

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{O1w}-\text{H11}\cdots\text{Br4}^{\text{i}}$	0.84	2.72	3.15 (3)	113
$\text{O1w}-\text{H12}\cdots\text{Br1}^{\text{ii}}$	0.84	2.83	3.65 (3)	167
$\text{N1}-\text{H1}\cdots\text{Br1}$	0.86	2.92	3.568 (11)	134
$\text{N2}-\text{H2}\cdots\text{O1w}$	0.86	2.04	2.80 (2)	146

Symmetry codes: (i) $-x + 1, y - \frac{1}{2}, -z + \frac{3}{2}$; (ii) $x + 1, y, z$.

Data collection: APEX2 (Bruker, 2008); cell refinement: SAINT (Bruker, 2008); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: X-SEED (Barbour, 2001); software used to prepare material for publication: publCIF (Westrip, 2009).

We thank the University of Malaya (RG020/09AFR) for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: TK2504).

References

- Al-Far, R. & Ali, B. F. (2008). *J. Chem. Crystallogr.* **37**, 333–341.
Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
Battaglia, L. P., Corradiab, B., Cariatif, K. & Koman, M. (1991). *Inorg. Chim. Acta*, **187**, 141–147.
Bruker (2008). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
Chen, W.-T., Zeng, X.-R., Fang, X.-N., Li, X.-F. & Kuang, H.-M. (2006). *Acta Cryst.* **C62**, m571–m573.
Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
Geselle, M. & Fuess, H. (1994). *Acta Cryst.* **C50**, 1582–1585.
Hatano, N., Nakashima, M., Horiuchi, K., Terao, H. & Ishihara, H. (2008). *Z. Naturforsch. Teil B*, **63**, 1181–1186.
Ishihara, H., Horiuchi, K., Gesing, T. M., Dou, S.-Q., Buhl, J. C. & Erk, P. (2002). *Z. Naturforsch.* **57**, 503–508.
Ishihara, H., Koriuchi, K., Svoboda, I., Fuess, H., Gesing, T. M., Buhl, J. C. & Terao, H. (2006). *Z. Naturforsch. Teil B*, **61**, 69–72.
Ravikumar, K., Venkata Lakshmi, N., Swamy, G. Y. S. K. & Chandra Mohan, K. (1995). *Acta Cryst.* **C51**, 1556–1558.
Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Waskowska, A. (1994). *Z. Kristallogr.* **209**, 750–754.
Westrip, S. P. (2009). publCIF. In preparation.
Zhang, H. & Fang, L. (2005). *Acta Cryst.* **E61**, m101–m102.

supplementary materials

Acta Cryst. (2009). E65, m973 [doi:10.1107/S160053680902813X]

1,4-Diazoniabicyclo[2.2.2]octane tetrabromidocadm(II) monohydrate

K. M. Lo and S. W. Ng

Experimental

Triethylenediammonium dibromide was prepared from the reaction of triethylenediamine (1 g, 1.68 mmol) with bromine (1:2) in the presence of excess hydrobromic acid. To this was added cadmium chloride hemipentahydrate (0.38 g, 1.68 mmol) in ethanol (50 ml). The mixture was heated for an hour. The filtered solution when allow to evaporate slowly yielded colorless crystals.

Refinement

C- and N-bound H atoms were placed at calculated positions (C–H 0.97 Å and N–H 0.86 Å) and were treated as riding on their parent atoms with $U(H)$ set to $1.2U_{eq}(C, N)$. The water-bound H atoms were placed in chemically sensible positions on the basis of hydrogen bonding interactions but were not refined.

The final difference Fourier map had a peak 0.2 Å from Cd1 and a hole 0.4 Å from Br4.

Figures




Fig. 1. Thermal ellipsoid plot (Barbour, 2001) of $[C_6H_{14}N_2][CdBr_4] \cdot H_2O$ at the 50% probability level. Hydrogen atoms are drawn as spheres of arbitrary radius.

1,4-Diazoniabicyclo[2.2.2]octane tetrabromidocadm(II) monohydrate

Crystal data

$(C_6H_{14}N_2)[CdBr_4] \cdot H_2O$

$M_r = 564.25$

Orthorhombic, $P2_12_12_1$

Hall symbol: P 2ac 2ab

$a = 8.6323$ (1) Å

$b = 11.8736$ (2) Å

$c = 13.5619$ (2) Å

$V = 1390.05$ (4) Å³

$Z = 4$

$F_{000} = 1048$

$D_x = 2.696$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 6547 reflections

$\theta = 2.8$ – 28.2°

$\mu = 13.04$ mm⁻¹

$T = 296$ K

Block, colorless

$0.30 \times 0.15 \times 0.05$ mm

supplementary materials

Data collection

Bruker SMART APEX diffractometer	2451 independent reflections
Radiation source: fine-focus sealed tube	2267 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.034$
$T = 296$ K	$\theta_{\text{max}} = 25.0^\circ$
ω scans	$\theta_{\text{min}} = 2.3^\circ$
Absorption correction: Multi-scan (SADABS; Sheldrick, 1996)	$h = -10 \rightarrow 10$
$T_{\text{min}} = 0.111$, $T_{\text{max}} = 0.562$	$k = -14 \rightarrow 14$
10779 measured reflections	$l = -16 \rightarrow 16$

Refinement

Refinement on F^2	Hydrogen site location: inferred from neighbouring sites
Least-squares matrix: full	H-atom parameters constrained
$R[F^2 > 2\sigma(F^2)] = 0.057$	$w = 1/[\sigma^2(F_o^2) + (0.1P)^2 + 5P]$
$wR(F^2) = 0.187$	where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.36$	$(\Delta/\sigma)_{\text{max}} = 0.001$
2451 reflections	$\Delta\rho_{\text{max}} = 2.21 \text{ e } \text{\AA}^{-3}$
128 parameters	$\Delta\rho_{\text{min}} = -2.09 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none
Secondary atom site location: difference Fourier map	Absolute structure: Flack (1983), 1021 Friedel pairs
	Flack parameter: 0.14 (3)

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Cd1	0.25482 (11)	0.47127 (8)	1.00019 (6)	0.0410 (3)
Br1	0.3099 (2)	0.28309 (13)	0.91553 (12)	0.0596 (5)
Br2	0.4870 (2)	0.59802 (15)	0.95680 (13)	0.0579 (5)
Br3	0.2467 (2)	0.46741 (14)	1.19150 (10)	0.0552 (4)
Br4	0.0032 (2)	0.56945 (19)	0.95508 (17)	0.0741 (6)
O1W	1.151 (3)	0.265 (2)	0.6666 (18)	0.144 (9)
H11	1.1965	0.2150	0.6336	0.216*
H12	1.2016	0.2746	0.7189	0.216*
N1	0.6148 (13)	0.4081 (10)	0.7757 (7)	0.040 (3)
H1	0.5272	0.4215	0.8034	0.048*
N2	0.8660 (14)	0.3708 (12)	0.6978 (13)	0.063 (4)
H2	0.9523	0.3577	0.6682	0.076*
C1	0.725 (2)	0.3684 (14)	0.8530 (12)	0.056 (4)
H1A	0.6860	0.3008	0.8846	0.067*
H1B	0.7396	0.4260	0.9029	0.067*
C2	0.882 (2)	0.3435 (18)	0.7975 (16)	0.086 (8)

H2A	0.9647	0.3877	0.8264	0.104*
H2B	0.9083	0.2645	0.8042	0.104*
C3	0.5912 (16)	0.3214 (11)	0.7016 (12)	0.045 (3)
H3A	0.5121	0.3449	0.6551	0.054*
H3B	0.5577	0.2519	0.7326	0.054*
C4	0.7409 (17)	0.3035 (13)	0.6494 (9)	0.047 (3)
H4A	0.7308	0.3262	0.5810	0.056*
H4B	0.7679	0.2242	0.6510	0.056*
C5	0.6697 (19)	0.5139 (11)	0.7291 (11)	0.044 (3)
H5A	0.6781	0.5729	0.7783	0.052*
H5B	0.5968	0.5382	0.6790	0.052*
C6	0.824 (2)	0.4924 (13)	0.6836 (17)	0.066 (5)
H6A	0.9016	0.5400	0.7143	0.079*
H6B	0.8206	0.5100	0.6138	0.079*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cd1	0.0398 (5)	0.0445 (6)	0.0387 (5)	0.0007 (4)	0.0034 (5)	-0.0059 (4)
Br1	0.0813 (11)	0.0441 (8)	0.0535 (8)	0.0002 (8)	0.0191 (7)	-0.0081 (7)
Br2	0.0570 (9)	0.0650 (10)	0.0515 (8)	-0.0032 (8)	-0.0029 (7)	0.0044 (7)
Br3	0.0546 (8)	0.0682 (10)	0.0428 (7)	0.0061 (9)	0.0040 (7)	0.0035 (6)
Br4	0.0606 (11)	0.0862 (13)	0.0756 (12)	0.0076 (10)	-0.0051 (9)	-0.0244 (10)
O1W	0.150 (18)	0.153 (19)	0.129 (16)	0.073 (17)	0.022 (14)	0.012 (15)
N1	0.037 (5)	0.063 (7)	0.020 (5)	-0.003 (5)	0.003 (4)	-0.009 (5)
N2	0.029 (6)	0.057 (8)	0.104 (12)	-0.002 (5)	0.019 (7)	-0.038 (8)
C1	0.067 (11)	0.051 (9)	0.051 (8)	0.005 (8)	-0.023 (8)	0.003 (7)
C2	0.089 (14)	0.071 (12)	0.099 (15)	0.039 (11)	-0.064 (13)	-0.051 (11)
C3	0.036 (6)	0.031 (7)	0.069 (9)	0.000 (5)	-0.002 (7)	0.000 (6)
C4	0.045 (7)	0.056 (8)	0.040 (6)	-0.008 (7)	-0.005 (7)	-0.014 (6)
C5	0.059 (9)	0.026 (6)	0.045 (7)	0.002 (6)	0.007 (7)	0.000 (6)
C6	0.066 (11)	0.040 (9)	0.091 (12)	-0.011 (7)	0.035 (10)	-0.017 (8)

Geometric parameters (\AA , $^\circ$)

Cd1—Br4	2.540 (2)	C1—H1A	0.9700
Cd1—Br1	2.557 (2)	C1—H1B	0.9700
Cd1—Br2	2.574 (2)	C2—H2A	0.9700
Cd1—Br3	2.596 (2)	C2—H2B	0.9700
O1W—H11	0.84	C3—C4	1.49 (2)
O1W—H12	0.84	C3—H3A	0.9700
N1—C3	1.453 (18)	C3—H3B	0.9700
N1—C5	1.484 (18)	C4—H4A	0.9700
N1—C1	1.491 (18)	C4—H4B	0.9700
N1—H1	0.8600	C5—C6	1.49 (2)
N2—C2	1.40 (3)	C5—H5A	0.9700
N2—C4	1.495 (18)	C5—H5B	0.9700
N2—C6	1.50 (2)	C6—H6A	0.9700
N2—H2	0.8600	C6—H6B	0.9700

supplementary materials

C1—C2	1.58 (3)		
Br4—Cd1—Br1	116.87 (7)	N2—C2—H2B	109.8
Br4—Cd1—Br2	110.01 (7)	C1—C2—H2B	109.8
Br1—Cd1—Br2	105.29 (7)	H2A—C2—H2B	108.2
Br4—Cd1—Br3	103.05 (7)	N1—C3—C4	107.9 (11)
Br1—Cd1—Br3	116.00 (7)	N1—C3—H3A	110.1
Br2—Cd1—Br3	105.05 (6)	C4—C3—H3A	110.1
H11—O1w—H12	107.7	N1—C3—H3B	110.1
C3—N1—C5	110.5 (10)	C4—C3—H3B	110.1
C3—N1—C1	110.6 (12)	H3A—C3—H3B	108.4
C5—N1—C1	111.3 (12)	C3—C4—N2	110.0 (11)
C3—N1—H1	108.1	C3—C4—H4A	109.7
C5—N1—H1	108.1	N2—C4—H4A	109.7
C1—N1—H1	108.1	C3—C4—H4B	109.7
C2—N2—C4	111.8 (15)	N2—C4—H4B	109.7
C2—N2—C6	111.8 (15)	H4A—C4—H4B	108.2
C4—N2—C6	106.5 (14)	N1—C5—C6	108.5 (11)
C2—N2—H2	108.9	N1—C5—H5A	110.0
C4—N2—H2	108.9	C6—C5—H5A	110.0
C6—N2—H2	108.9	N1—C5—H5B	110.0
N1—C1—C2	105.7 (13)	C6—C5—H5B	110.0
N1—C1—H1A	110.6	H5A—C5—H5B	108.4
C2—C1—H1A	110.6	C5—C6—N2	109.1 (12)
N1—C1—H1B	110.6	C5—C6—H6A	109.9
C2—C1—H1B	110.6	N2—C6—H6A	109.9
H1A—C1—H1B	108.7	C5—C6—H6B	109.9
N2—C2—C1	109.5 (13)	N2—C6—H6B	109.9
N2—C2—H2A	109.8	H6A—C6—H6B	108.3
C1—C2—H2A	109.8		
C3—N1—C1—C2	60.9 (16)	C2—N2—C4—C3	57.5 (17)
C5—N1—C1—C2	-62.3 (16)	C6—N2—C4—C3	-64.8 (17)
C4—N2—C2—C1	-60.2 (17)	C3—N1—C5—C6	-63.7 (16)
C6—N2—C2—C1	59.0 (19)	C1—N1—C5—C6	59.6 (16)
N1—C1—C2—N2	2(2)	N1—C5—C6—N2	2(2)
C5—N1—C3—C4	58.4 (15)	C2—N2—C6—C5	-63 (2)
C1—N1—C3—C4	-65.3 (15)	C4—N2—C6—C5	59.1 (19)
N1—C3—C4—N2	6.0 (17)		

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O1w—H11 \cdots Br4 ⁱ	0.84	2.72	3.15 (3)	113
O1w—H12 \cdots Br1 ⁱⁱ	0.84	2.83	3.65 (3)	167
N1—H1 \cdots Br1	0.86	2.92	3.568 (11)	134
N2—H2 \cdots O1w	0.86	2.04	2.80 (2)	146

Symmetry codes: (i) $-x+1, y-1/2, -z+3/2$; (ii) $x+1, y, z$.

Fig. 1

