

(μ -Pyridine-2-carbaldehyde azine)bis-[bis(triphenylphosphine- κP)copper(I)] bis(tetrafluoridoborate) dichloromethane disolvate

Li Yang,^a Yu Xie,^{a,b*} Jianping Zou,^a Jie Jia^a and Xiaowei Hong^a

^aKey Laboratory of Nondestructive Testing (Ministry of Education), Nanchang Hangkong University, Nanchang 330063, People's Republic of China, and ^bKey Laboratory of Photochemical Conversion and Optoelectronic Materials, TIPC, CAS, Beijing 100190, People's Republic of China
Correspondence e-mail: xieyu_121@163.com


Received 5 July 2009; accepted 28 July 2009

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.013\text{ \AA}$; disorder in solvent or counterion; R factor = 0.081; wR factor = 0.268; data-to-parameter ratio = 13.6.

In the centrosymmetric title complex, $[\text{Cu}_2(\text{C}_{12}\text{H}_{10}\text{N}_4)(\text{C}_{18}\text{H}_{15}\text{P}_4)](\text{BF}_4)_2 \cdot 2\text{CH}_2\text{Cl}_2$, the Cu^I atom adopts a distorted tetrahedral geometry, defined by two P atoms from two triphenylphosphine ligands and two N atoms from a pyridine-2-carbaldehyde azine ligand. The two Cu atoms are bridged by the centrosymmetric pyridine-2-carbaldehyde azine ligand. The F atoms of the tetrafluoridoborate anion are disordered over two sites [occupancy factors = 0.68 (5) and 0.32 (5)]. The dichloromethane solvent molecule is disordered over four sites, with occupancy factors of 0.513 (4), 0.173 (5), 0.141 (5) and 0.173 (5).

Related literature

For general background to the use of neutral pyridine-azines in the construction of di-, tri- and polynuclear complexes, see: Tuna *et al.* (2003); Guo *et al.* (2002); Hamblin *et al.* (2002). For related structures, see: Mo *et al.* (2006); Zhou *et al.* (2006).



Experimental

Crystal data

$[\text{Cu}_2(\text{C}_{12}\text{H}_{10}\text{N}_4)(\text{C}_{18}\text{H}_{15}\text{P}_4)](\text{BF}_4)_2 \cdot 2\text{CH}_2\text{Cl}_2$	$\beta = 115.117(2)^\circ$
$M_r = 1729.89$	$V = 4532.9(9)\text{ \AA}^3$
Monoclinic, $P2_1/n$	$Z = 2$
$a = 13.0932(16)\text{ \AA}$	Mo $K\alpha$ radiation
$b = 27.501(3)\text{ \AA}$	$\mu = 0.72\text{ mm}^{-1}$
$c = 13.9033(18)\text{ \AA}$	$T = 293\text{ K}$
	$0.35 \times 0.32 \times 0.30\text{ mm}$

Data collection

Bruker APEXII CCD diffractometer	24119 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Sheldrick, 1996)	7643 independent reflections
$T_{\min} = 0.787$, $T_{\max} = 0.814$	3919 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.063$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.081$	2 restraints
$wR(F^2) = 0.268$	H-atom parameters constrained
$S = 0.95$	$\Delta\rho_{\text{max}} = 1.44\text{ e \AA}^{-3}$
7643 reflections	$\Delta\rho_{\text{min}} = -0.62\text{ e \AA}^{-3}$
564 parameters	

Table 1
Selected bond lengths (Å).

Cu1—N1	2.061 (6)	Cu1—P1	2.2741 (19)
Cu1—N2	2.175 (5)	Cu1—P2	2.292 (2)

Data collection: *APEX2* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The authors thank the Key Laboratory of Photochemical Conversion and Optoelectronic Materials, TIPC, CAS, and Jiangxi Provincial Department of Education for financial support of this work. The Materials Chemistry Department of Nanchang Hangkong University is also acknowledged.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HY2211).

References

- Bruker (2007). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Guo, D., Duan, C., Fang, C. & Meng, Q. (2002). *J. Chem. Soc. Dalton Trans.* pp. 834–836.
- Hamblin, J., Jackson, A., Alcock, N. W. & Hannon, M. J. (2002). *J. Chem. Soc. Dalton Trans.* pp. 1635–1641.
- Mo, J., Qian, H.-Y., Du, X.-D. & Chen, W. (2006). *Acta Cryst. E62*, m726–m727.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Tuna, F., Hamblin, J., Jackson, A., Clarkson, G., Alcock, N. W. & Hannon, M. J. (2003). *Dalton Trans.* pp. 2141–2148.
- Zhou, X., Wu, T. & Li, D. (2006). *Inorg. Chim. Acta*, **359**, 1442–1448.

supplementary materials

Acta Cryst. (2009). E65, m1017 [doi:10.1107/S160053680902995X]

(μ -Pyridine-2-carbaldehyde azine)bis[bis(triphenylphosphine- κP)copper(I)] bis(tetrafluoridoborate) dichloromethane disolvate

L. Yang, Y. Xie, J. Zou, J. Jia and X. Hong

Comment

Neutral pyridine-azines are excellent bridging ligands in coordination chemistry. They are often used to construct some di-, tri- and polynuclear complexes (Tuna *et al.*, 2003; Guo *et al.*, 2002; Hamblin *et al.*, 2002; Zhou *et al.*, 2006). We describe here the synthesis and structure of a new copper(I) compound with a pyridine-2-carbaldehyde azine ligand.

The molecular structure of the title compound is depicted in Fig. 1. The complex is a centrosymmetric dimer with two Cu^I atoms bridged by a pyridine-2-carbaldehyde azine ligand. The Cu^I atom has a distorted tetrahedral geometry with two P atoms from two triphenylphosphine ligands and two N atoms from the bridging pyridine-2-carbaldehyde azine ligand. The bond angles around the Cu atom are in the range of 77.5 (2) $^\circ$ (N1—Cu1—N2) to 124.02 (7) (P1—Cu1—P2) $^\circ$. The Cu—P [2.2741 (19) and 2.292 (2) Å] and Cu—N [2.061 (6) and 2.175 (5) Å] bond distances (Table 1) are within the normal ranges for analogous complexes (Mo *et al.*, 2006; Zhou *et al.*, 2006).

Experimental

Pyridine-2-carbaldehyde and CuBF₄·4CH₃CN were prepared by literature method (Mo *et al.*, 2006; Zhou *et al.*, 2006). The title compound was prepared by reacting pyridine-2-carbaldehyde (0.021 g, 0.1 mmol), CuBF₄·4CH₃CN (0.031 g, 0.1 mmol) and triphenylphosphine (0.052 g, 0.2 mmol) in 30 ml dry dichloromethane under N₂ atmosphere. Brown needle crystals suitable for X-ray analysis were obtained by vapor diffusion of diethyl ether into the solution of the title compound in dichloromethane (yield 83%).

Refinement

H atoms were positioned geometrically and treated as riding, with C—H = 0.93 (aromatic) and 0.97 (dichloromethane) Å and with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$. The tetrafluoroborate anion and dichloromethane solvent molecule are disordered. The tetrafluoroborate anion is splitted into two components with site occupancy factor (SOF) values of 0.68 (5) and 0.32 (5). The dichloromethane molecule is splitted into four components with SOF values of 0.513 (4), 0.173 (5), 0.141 (5) and 0.173 (5), respectively. The disorder of the anion and solvent molecule may cause high value of the weighted *R* factor for this structure. The highest peak and deepest hole in the final difference map were associated with atom C45 (at distances of 0.78 Å).

supplementary materials

Figures




Fig. 1. The molecular structure of the title compound. Displacement ellipsoids are drawn at the 30% probability level. H atoms, tetrafluoroborate anion and dichloromethane solvent molecule are omitted for clarity. [Symmetry code: (i) $1 - x, 1 - y, 1 - z$.]

(μ -Pyridine-2-carbaldehyde azine)bis[bis(triphenylphosphine- κP)copper(II)] bis(tetrafluoroborate) dichloromethane disolvate

Crystal data

$[\text{Cu}_2(\text{C}_{12}\text{H}_{10}\text{N}_4)(\text{C}_{18}\text{H}_{15}\text{P})_4](\text{BF}_4)_2 \cdot 2\text{CH}_2\text{Cl}_2$	$F_{000} = 1772$
$M_r = 1729.89$	$D_x = 1.267 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/n$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2yn	Cell parameters from 24119 reflections
$a = 13.0932 (16) \text{ \AA}$	$\theta = 2.2\text{--}25.0^\circ$
$b = 27.501 (3) \text{ \AA}$	$\mu = 0.72 \text{ mm}^{-1}$
$c = 13.9033 (18) \text{ \AA}$	$T = 293 \text{ K}$
$\beta = 115.117 (2)^\circ$	Block, brown
$V = 4532.9 (9) \text{ \AA}^3$	$0.35 \times 0.32 \times 0.30 \text{ mm}$
$Z = 2$	

Data collection

Bruker APEXII CCD diffractometer	7643 independent reflections
Radiation source: fine-focus sealed tube	3919 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.063$
$T = 293 \text{ K}$	$\theta_{\text{max}} = 25.0^\circ$
φ and ω scans	$\theta_{\text{min}} = 1.5^\circ$
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	$h = 0 \rightarrow 15$
$T_{\text{min}} = 0.787, T_{\text{max}} = 0.814$	$k = 0 \rightarrow 32$
24119 measured reflections	$l = -16 \rightarrow 14$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.081$	H-atom parameters constrained
$wR(F^2) = 0.268$	$w = 1/[\sigma^2(F_o^2) + (0.1662P)^2]$
$S = 0.95$	where $P = (F_o^2 + 2F_c^2)/3$
	$(\Delta/\sigma)_{\text{max}} = 0.001$

7643 reflections $\Delta\rho_{\max} = 1.44 \text{ e Å}^{-3}$
 564 parameters $\Delta\rho_{\min} = -0.62 \text{ e Å}^{-3}$
 2 restraints Extinction correction: none
 Primary atom site location: structure-invariant direct methods

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Cu1	0.39337 (7)	0.56453 (3)	0.34029 (6)	0.0439 (3)	
F1	0.9059 (12)	0.6834 (11)	0.7796 (13)	0.130 (8)	0.68 (5)
F2	0.919 (3)	0.7544 (5)	0.7121 (13)	0.151 (10)	0.68 (5)
F3	0.956 (3)	0.6852 (17)	0.647 (4)	0.152 (10)	0.68 (5)
F4	1.077 (2)	0.7109 (11)	0.806 (2)	0.152 (8)	0.68 (5)
F1'	1.011 (6)	0.7535 (15)	0.713 (3)	0.17 (2)	0.32 (5)
F2'	1.065 (5)	0.684 (2)	0.799 (5)	0.145 (16)	0.32 (5)
F3'	0.906 (3)	0.721 (2)	0.789 (2)	0.129 (17)	0.32 (5)
F4'	0.910 (5)	0.685 (3)	0.645 (8)	0.123 (15)	0.32 (5)
N1	0.4999 (5)	0.6178 (2)	0.4340 (4)	0.0469 (14)	
N2	0.4973 (5)	0.52483 (19)	0.4841 (4)	0.0437 (13)	
Cl1	0.7434 (6)	0.1422 (2)	0.1283 (7)	0.140 (3)	0.513 (4)
Cl2	0.6074 (8)	0.2242 (3)	0.1389 (9)	0.163 (3)	0.513 (4)
Cl3	0.4482 (17)	0.2880 (6)	0.2405 (19)	0.140 (3)	0.173 (5)
Cl4	0.303 (3)	0.2584 (14)	0.038 (4)	0.163 (3)	0.173 (5)
Cl5	0.687 (2)	0.2339 (8)	0.229 (3)	0.140 (3)	0.141 (5)
Cl6	0.846 (3)	0.2284 (12)	0.451 (3)	0.163 (3)	0.141 (5)
Cl7	0.2701 (18)	0.2418 (7)	0.105 (2)	0.140 (3)	0.173 (5)
Cl8	0.473 (2)	0.2573 (9)	0.090 (2)	0.163 (3)	0.173 (5)
P1	0.44194 (15)	0.54008 (7)	0.20904 (13)	0.0438 (5)	
P2	0.21113 (15)	0.57584 (7)	0.31766 (14)	0.0456 (5)	
B1	0.9692 (15)	0.7102 (6)	0.7357 (12)	0.103 (4)	
C1	0.5039 (7)	0.6647 (3)	0.4100 (7)	0.064 (2)	
H1	0.4538	0.6759	0.3435	0.077*	
C2	0.5824 (8)	0.6983 (3)	0.4827 (8)	0.082 (3)	
H2	0.5860	0.7302	0.4621	0.098*	
C3	0.6508 (8)	0.6832 (3)	0.5813 (8)	0.088 (3)	
H3	0.6985	0.7050	0.6314	0.106*	
C4	0.6489 (7)	0.6349 (3)	0.6066 (7)	0.074 (2)	
H4	0.6985	0.6235	0.6731	0.089*	
C5	0.5730 (6)	0.6032 (3)	0.5329 (6)	0.0514 (18)	
C6	0.5683 (6)	0.5522 (2)	0.5556 (5)	0.0482 (17)	
H6	0.6163	0.5395	0.6213	0.058*	
C7	0.3910 (6)	0.5791 (3)	0.0920 (5)	0.0484 (17)	
C8	0.4119 (8)	0.6284 (3)	0.1071 (6)	0.067 (2)	
H8	0.4525	0.6402	0.1756	0.080*	
C9	0.3739 (8)	0.6607 (3)	0.0231 (7)	0.081 (3)	
H9	0.3866	0.6939	0.0347	0.097*	
C10	0.3171 (8)	0.6428 (3)	-0.0782 (7)	0.075 (2)	

supplementary materials

H10	0.2902	0.6640	-0.1356	0.090*
C11	0.3000 (7)	0.5942 (3)	-0.0948 (6)	0.067 (2)
H11	0.2636	0.5824	-0.1638	0.080*
C12	0.3363 (6)	0.5620 (3)	-0.0101 (5)	0.0536 (18)
H12	0.3237	0.5289	-0.0224	0.064*
C13	0.3871 (6)	0.4803 (2)	0.1559 (5)	0.0468 (17)
C14	0.4509 (7)	0.4451 (3)	0.1386 (7)	0.065 (2)
H14	0.5260	0.4513	0.1535	0.079*
C15	0.4029 (9)	0.3994 (3)	0.0981 (8)	0.085 (3)
H15	0.4456	0.3756	0.0845	0.101*
C16	0.2935 (9)	0.3901 (3)	0.0789 (8)	0.083 (3)
H16	0.2636	0.3595	0.0546	0.100*
C17	0.2244 (8)	0.4257 (3)	0.0947 (7)	0.073 (2)
H17	0.1492	0.4196	0.0792	0.087*
C18	0.2741 (7)	0.4699 (3)	0.1344 (6)	0.060 (2)
H18	0.2313	0.4938	0.1475	0.072*
C19	0.5921 (6)	0.5342 (3)	0.2423 (5)	0.0485 (17)
C20	0.6401 (7)	0.5499 (3)	0.1748 (6)	0.060 (2)
H20	0.5952	0.5659	0.1120	0.072*
C21	0.7526 (7)	0.5422 (3)	0.1992 (7)	0.068 (2)
H21	0.7828	0.5533	0.1536	0.081*
C22	0.8193 (7)	0.5184 (3)	0.2904 (7)	0.064 (2)
H22	0.8946	0.5123	0.3060	0.077*
C23	0.7752 (7)	0.5031 (3)	0.3599 (7)	0.074 (2)
H23	0.8209	0.4876	0.4231	0.089*
C24	0.6624 (7)	0.5111 (3)	0.3344 (6)	0.063 (2)
H24	0.6331	0.5006	0.3812	0.076*
C25	0.1225 (6)	0.5213 (3)	0.2757 (6)	0.0512 (18)
C26	0.1505 (7)	0.4826 (3)	0.3480 (6)	0.061 (2)
H26	0.2064	0.4864	0.4167	0.073*
C27	0.0942 (7)	0.4386 (3)	0.3166 (8)	0.071 (2)
H27	0.1140	0.4128	0.3641	0.085*
C28	0.0077 (7)	0.4326 (4)	0.2137 (8)	0.077 (3)
H28	-0.0296	0.4030	0.1927	0.093*
C29	-0.0202 (7)	0.4710 (3)	0.1457 (8)	0.072 (2)
H29	-0.0781	0.4678	0.0779	0.086*
C30	0.0366 (6)	0.5151 (3)	0.1763 (6)	0.0582 (19)
H30	0.0160	0.5409	0.1284	0.070*
C31	0.1930 (6)	0.5960 (3)	0.4352 (6)	0.0567 (19)
C32	0.0874 (7)	0.5932 (3)	0.4398 (7)	0.070 (2)
H32	0.0263	0.5787	0.3847	0.084*
C33	0.0749 (9)	0.6121 (4)	0.5264 (7)	0.079 (3)
H33	0.0043	0.6124	0.5273	0.094*
C34	0.1650 (8)	0.6302 (4)	0.6095 (7)	0.080 (3)
H34	0.1563	0.6413	0.6688	0.096*
C35	0.2679 (9)	0.6326 (4)	0.6087 (7)	0.089 (3)
H35	0.3288	0.6454	0.6667	0.107*
C36	0.2826 (7)	0.6156 (3)	0.5201 (6)	0.070 (2)
H36	0.3529	0.6177	0.5189	0.083*

C37	0.1303 (6)	0.6189 (3)	0.2139 (6)	0.0516 (18)	
C38	0.0403 (8)	0.6452 (3)	0.2120 (7)	0.079 (3)	
H38	0.0229	0.6437	0.2702	0.095*	
C39	-0.0260 (8)	0.6740 (4)	0.1252 (8)	0.091 (3)	
H39	-0.0890	0.6901	0.1241	0.109*	
C40	0.0028 (9)	0.6785 (3)	0.0414 (7)	0.082 (3)	
H40	-0.0394	0.6984	-0.0158	0.098*	
C41	0.0906 (9)	0.6544 (4)	0.0423 (7)	0.086 (3)	
H41	0.1109	0.6574	-0.0139	0.104*	
C42	0.1533 (8)	0.6240 (3)	0.1295 (7)	0.075 (3)	
H42	0.2137	0.6067	0.1281	0.090*	
C43	0.709 (2)	0.2027 (8)	0.109 (2)	0.131 (7)	0.513 (4)
H43A	0.7768	0.2211	0.1497	0.157*	0.513 (4)
H43B	0.6878	0.2098	0.0346	0.157*	0.513 (4)
C44	0.325 (7)	0.255 (3)	0.169 (7)	0.131 (7)	0.173 (5)
H44A	0.3341	0.2214	0.1924	0.157*	0.173 (5)
H44B	0.2618	0.2690	0.1780	0.157*	0.173 (5)
C45	0.798 (7)	0.204 (3)	0.328 (8)	0.131 (7)	0.141 (5)
H45A	0.7746	0.1705	0.3323	0.157*	0.141 (5)
H45B	0.8602	0.2021	0.3081	0.157*	0.141 (5)
C46	0.330 (9)	0.260 (4)	0.020 (11)	0.131 (7)	0.173 (5)
H46A	0.3051	0.2395	-0.0415	0.157*	0.173 (5)
H46B	0.3070	0.2935	-0.0035	0.157*	0.173 (5)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cu1	0.0449 (5)	0.0480 (5)	0.0388 (5)	0.0011 (4)	0.0178 (4)	-0.0006 (4)
F1	0.133 (10)	0.127 (17)	0.133 (11)	0.000 (9)	0.059 (8)	0.021 (10)
F2	0.16 (2)	0.080 (8)	0.176 (13)	0.035 (10)	0.036 (12)	0.016 (7)
F3	0.15 (3)	0.182 (16)	0.139 (15)	0.01 (2)	0.07 (2)	-0.059 (11)
F4	0.095 (10)	0.135 (18)	0.157 (12)	-0.007 (13)	-0.014 (8)	-0.005 (15)
F1'	0.16 (5)	0.14 (3)	0.18 (3)	-0.01 (3)	0.03 (3)	0.03 (2)
F2'	0.12 (3)	0.12 (3)	0.16 (3)	0.02 (3)	0.02 (2)	0.01 (3)
F3'	0.13 (2)	0.13 (4)	0.13 (2)	0.00 (2)	0.059 (16)	-0.01 (2)
F4'	0.10 (3)	0.13 (3)	0.14 (3)	0.00 (3)	0.05 (3)	-0.039 (19)
N1	0.052 (4)	0.038 (3)	0.050 (3)	0.001 (3)	0.021 (3)	0.000 (3)
N2	0.047 (3)	0.044 (3)	0.044 (3)	0.006 (3)	0.023 (3)	0.000 (3)
Cl1	0.115 (4)	0.079 (3)	0.200 (6)	0.004 (3)	0.042 (4)	0.054 (4)
Cl2	0.141 (6)	0.136 (6)	0.219 (8)	0.004 (5)	0.083 (6)	0.030 (5)
Cl3	0.115 (4)	0.079 (3)	0.200 (6)	0.004 (3)	0.042 (4)	0.054 (4)
Cl4	0.141 (6)	0.136 (6)	0.219 (8)	0.004 (5)	0.083 (6)	0.030 (5)
Cl5	0.115 (4)	0.079 (3)	0.200 (6)	0.004 (3)	0.042 (4)	0.054 (4)
Cl6	0.141 (6)	0.136 (6)	0.219 (8)	0.004 (5)	0.083 (6)	0.030 (5)
Cl7	0.115 (4)	0.079 (3)	0.200 (6)	0.004 (3)	0.042 (4)	0.054 (4)
Cl8	0.141 (6)	0.136 (6)	0.219 (8)	0.004 (5)	0.083 (6)	0.030 (5)
P1	0.0456 (11)	0.0486 (11)	0.0392 (9)	0.0020 (8)	0.0200 (8)	0.0000 (8)
P2	0.0431 (10)	0.0539 (11)	0.0409 (10)	0.0029 (8)	0.0190 (8)	0.0010 (8)

supplementary materials

B1	0.124 (13)	0.080 (10)	0.093 (10)	0.018 (10)	0.033 (10)	-0.015 (8)
C1	0.068 (6)	0.052 (5)	0.066 (5)	0.001 (4)	0.022 (4)	-0.005 (4)
C2	0.082 (7)	0.052 (5)	0.092 (7)	-0.011 (5)	0.019 (5)	0.001 (5)
C3	0.081 (7)	0.064 (6)	0.091 (7)	-0.014 (5)	0.009 (5)	-0.014 (5)
C4	0.067 (6)	0.058 (5)	0.071 (5)	-0.004 (4)	0.005 (4)	-0.007 (4)
C5	0.049 (4)	0.049 (4)	0.056 (4)	0.003 (3)	0.022 (4)	-0.002 (3)
C6	0.050 (4)	0.051 (4)	0.042 (4)	0.005 (3)	0.018 (3)	0.000 (3)
C7	0.049 (4)	0.056 (5)	0.046 (4)	0.005 (3)	0.025 (3)	0.002 (3)
C8	0.085 (6)	0.062 (5)	0.050 (4)	-0.004 (5)	0.025 (4)	0.001 (4)
C9	0.101 (8)	0.062 (6)	0.072 (6)	0.000 (5)	0.029 (5)	0.006 (5)
C10	0.088 (7)	0.075 (6)	0.061 (5)	-0.002 (5)	0.030 (5)	0.017 (5)
C11	0.076 (6)	0.074 (6)	0.052 (5)	-0.001 (5)	0.028 (4)	0.001 (4)
C12	0.058 (5)	0.054 (4)	0.047 (4)	0.006 (4)	0.021 (3)	-0.001 (3)
C13	0.052 (4)	0.047 (4)	0.050 (4)	0.003 (3)	0.030 (3)	0.001 (3)
C14	0.063 (5)	0.057 (5)	0.082 (6)	-0.005 (4)	0.036 (5)	-0.010 (4)
C15	0.083 (7)	0.066 (6)	0.107 (8)	0.002 (5)	0.043 (6)	-0.021 (5)
C16	0.081 (7)	0.065 (6)	0.104 (7)	-0.019 (5)	0.039 (6)	-0.016 (5)
C17	0.070 (6)	0.074 (6)	0.075 (6)	-0.016 (5)	0.032 (5)	-0.016 (5)
C18	0.063 (5)	0.060 (5)	0.062 (5)	0.005 (4)	0.032 (4)	-0.007 (4)
C19	0.051 (4)	0.055 (4)	0.043 (4)	0.000 (3)	0.023 (3)	-0.001 (3)
C20	0.052 (5)	0.072 (5)	0.057 (5)	0.003 (4)	0.024 (4)	0.007 (4)
C21	0.056 (5)	0.086 (6)	0.073 (6)	-0.009 (5)	0.038 (5)	-0.003 (5)
C22	0.050 (5)	0.074 (6)	0.077 (6)	0.003 (4)	0.035 (4)	-0.007 (5)
C23	0.052 (5)	0.095 (7)	0.073 (6)	0.021 (5)	0.023 (4)	0.009 (5)
C24	0.057 (5)	0.077 (6)	0.063 (5)	0.011 (4)	0.033 (4)	0.009 (4)
C25	0.049 (4)	0.055 (4)	0.057 (4)	0.004 (4)	0.029 (4)	0.003 (3)
C26	0.062 (5)	0.061 (5)	0.064 (5)	0.004 (4)	0.031 (4)	0.006 (4)
C27	0.071 (6)	0.066 (6)	0.090 (6)	0.003 (5)	0.048 (5)	0.013 (5)
C28	0.061 (6)	0.074 (6)	0.099 (7)	-0.006 (5)	0.036 (5)	-0.003 (5)
C29	0.060 (6)	0.077 (6)	0.077 (6)	-0.011 (5)	0.028 (4)	-0.009 (5)
C30	0.055 (5)	0.062 (5)	0.059 (5)	-0.005 (4)	0.026 (4)	0.002 (4)
C31	0.054 (5)	0.071 (5)	0.049 (4)	0.011 (4)	0.026 (4)	0.003 (4)
C32	0.068 (6)	0.087 (6)	0.059 (5)	0.005 (5)	0.030 (4)	0.001 (4)
C33	0.080 (7)	0.106 (7)	0.070 (6)	0.016 (6)	0.051 (5)	0.005 (5)
C34	0.079 (7)	0.114 (8)	0.059 (5)	0.015 (6)	0.041 (5)	-0.005 (5)
C35	0.085 (7)	0.115 (8)	0.058 (5)	0.007 (6)	0.022 (5)	-0.021 (5)
C36	0.061 (5)	0.094 (7)	0.055 (5)	0.000 (5)	0.025 (4)	-0.010 (4)
C37	0.052 (5)	0.059 (5)	0.052 (4)	0.004 (4)	0.030 (4)	0.003 (3)
C38	0.080 (6)	0.095 (7)	0.067 (6)	0.025 (6)	0.037 (5)	0.022 (5)
C39	0.085 (7)	0.101 (8)	0.085 (7)	0.047 (6)	0.035 (6)	0.026 (6)
C40	0.083 (7)	0.081 (7)	0.070 (6)	0.017 (5)	0.021 (5)	0.023 (5)
C41	0.095 (7)	0.102 (8)	0.070 (6)	0.026 (6)	0.042 (5)	0.032 (5)
C42	0.075 (6)	0.089 (7)	0.068 (5)	0.025 (5)	0.037 (5)	0.019 (5)
C43	0.114 (17)	0.079 (13)	0.18 (2)	-0.002 (11)	0.048 (14)	0.025 (14)
C44	0.114 (17)	0.079 (13)	0.18 (2)	-0.002 (11)	0.048 (14)	0.025 (14)
C45	0.114 (17)	0.079 (13)	0.18 (2)	-0.002 (11)	0.048 (14)	0.025 (14)
C46	0.114 (17)	0.079 (13)	0.18 (2)	-0.002 (11)	0.048 (14)	0.025 (14)

Geometric parameters (Å, °)

Cu1—N1	2.061 (6)	C16—C17	1.412 (12)
Cu1—N2	2.175 (5)	C16—H16	0.9300
Cu1—P1	2.2741 (19)	C17—C18	1.379 (11)
Cu1—P2	2.292 (2)	C17—H17	0.9300
F1—B1	1.43 (2)	C18—H18	0.9300
F2—B1	1.35 (2)	C19—C24	1.374 (10)
F3—B1	1.36 (4)	C19—C20	1.402 (10)
F4—B1	1.33 (3)	C20—C21	1.379 (11)
F1'—B1	1.40 (4)	C20—H20	0.9300
F2'—B1	1.39 (5)	C21—C22	1.364 (12)
F3'—B1	1.36 (3)	C21—H21	0.9300
F4'—B1	1.36 (9)	C22—C23	1.385 (12)
N1—C1	1.338 (9)	C22—H22	0.9300
N1—C5	1.361 (9)	C23—C24	1.382 (11)
N2—C6	1.278 (8)	C23—H23	0.9300
N2—N2 ⁱ	1.428 (10)	C24—H24	0.9300
Cl1—C43	1.72 (2)	C25—C30	1.373 (10)
Cl2—C43	1.66 (3)	C25—C26	1.402 (10)
Cl3—C44	1.75 (8)	C26—C27	1.386 (11)
Cl4—C44	1.72 (9)	C26—H26	0.9300
Cl5—C45	1.74 (9)	C27—C28	1.407 (13)
Cl6—C45	1.69 (10)	C27—H27	0.9300
Cl7—C46	1.75 (12)	C28—C29	1.361 (12)
Cl8—C46	1.71 (11)	C28—H28	0.9300
P1—C13	1.820 (7)	C29—C30	1.391 (11)
P1—C7	1.823 (7)	C29—H29	0.9300
P1—C19	1.825 (7)	C30—H30	0.9300
P2—C37	1.819 (7)	C31—C36	1.373 (11)
P2—C31	1.833 (7)	C31—C32	1.414 (11)
P2—C25	1.833 (8)	C32—C33	1.384 (11)
C1—C2	1.432 (11)	C32—H32	0.9300
C1—H1	0.9300	C33—C34	1.347 (13)
C2—C3	1.346 (12)	C33—H33	0.9300
C2—H2	0.9300	C34—C35	1.354 (13)
C3—C4	1.378 (12)	C34—H34	0.9300
C3—H3	0.9300	C35—C36	1.404 (11)
C4—C5	1.391 (10)	C35—H35	0.9300
C4—H4	0.9300	C36—H36	0.9300
C5—C6	1.444 (10)	C37—C42	1.338 (11)
C6—H6	0.9300	C37—C38	1.372 (11)
C7—C12	1.374 (9)	C38—C39	1.396 (12)
C7—C8	1.381 (10)	C38—H38	0.9300
C8—C9	1.380 (11)	C39—C40	1.374 (13)
C8—H8	0.9300	C39—H39	0.9300
C9—C10	1.374 (12)	C40—C41	1.322 (13)
C9—H9	0.9300	C40—H40	0.9300

supplementary materials

C10—C11	1.358 (11)	C41—C42	1.413 (11)
C10—H10	0.9300	C41—H41	0.9300
C11—C12	1.386 (10)	C42—H42	0.9300
C11—H11	0.9300	C43—H43A	0.9700
C12—H12	0.9300	C43—H43B	0.9700
C13—C14	1.365 (10)	C44—H44A	0.9700
C13—C18	1.408 (10)	C44—H44B	0.9700
C14—C15	1.412 (11)	C45—H45A	0.9700
C14—H14	0.9300	C45—H45B	0.9700
C15—C16	1.364 (13)	C46—H46A	0.9700
C15—H15	0.9300	C46—H46B	0.9700
N1—Cu1—N2	77.5 (2)	C21—C20—C19	121.6 (7)
N1—Cu1—P1	111.87 (17)	C21—C20—H20	119.2
N2—Cu1—P1	108.09 (15)	C19—C20—H20	119.2
N1—Cu1—P2	112.10 (17)	C22—C21—C20	119.9 (8)
N2—Cu1—P2	113.75 (15)	C22—C21—H21	120.1
P1—Cu1—P2	124.02 (7)	C20—C21—H21	120.1
C1—N1—C5	116.9 (6)	C21—C22—C23	120.1 (8)
C1—N1—Cu1	128.0 (5)	C21—C22—H22	119.9
C5—N1—Cu1	115.2 (5)	C23—C22—H22	119.9
C6—N2—N2 ⁱ	113.5 (7)	C24—C23—C22	119.3 (8)
C6—N2—Cu1	112.5 (4)	C24—C23—H23	120.4
N2 ⁱ —N2—Cu1	133.9 (6)	C22—C23—H23	120.4
C13—P1—C7	103.8 (3)	C19—C24—C23	122.2 (8)
C13—P1—C19	102.7 (3)	C19—C24—H24	118.9
C7—P1—C19	103.0 (3)	C23—C24—H24	118.9
C13—P1—Cu1	113.3 (2)	C30—C25—C26	118.7 (7)
C7—P1—Cu1	114.9 (2)	C30—C25—P2	124.0 (6)
C19—P1—Cu1	117.4 (2)	C26—C25—P2	117.1 (6)
C37—P2—C31	104.9 (3)	C27—C26—C25	119.6 (8)
C37—P2—C25	101.8 (3)	C27—C26—H26	120.2
C31—P2—C25	103.1 (3)	C25—C26—H26	120.2
C37—P2—Cu1	114.8 (2)	C26—C27—C28	121.0 (8)
C31—P2—Cu1	116.1 (3)	C26—C27—H27	119.5
C25—P2—Cu1	114.5 (2)	C28—C27—H27	119.5
F4—B1—F2	114.7 (17)	C29—C28—C27	118.5 (9)
F4'—B1—F3'	113 (3)	C29—C28—H28	120.8
F4—B1—F3	111 (2)	C27—C28—H28	120.8
F2—B1—F3	112 (2)	C28—C29—C30	120.9 (9)
F4'—B1—F2'	109 (4)	C28—C29—H29	119.5
F3'—B1—F2'	111 (4)	C30—C29—H29	119.5
F4'—B1—F1'	110 (4)	C25—C30—C29	121.3 (8)
F3'—B1—F1'	110 (2)	C25—C30—H30	119.4
F2'—B1—F1'	104 (3)	C29—C30—H30	119.4
F4—B1—F1	108.5 (19)	C36—C31—C32	118.5 (7)
F2—B1—F1	105.0 (17)	C36—C31—P2	119.9 (6)
F3—B1—F1	105 (2)	C32—C31—P2	121.5 (6)
N1—C1—C2	122.5 (8)	C33—C32—C31	119.9 (8)

N1—C1—H1	118.7	C33—C32—H32	120.1
C2—C1—H1	118.7	C31—C32—H32	120.1
C3—C2—C1	119.2 (8)	C34—C33—C32	120.0 (9)
C3—C2—H2	120.4	C34—C33—H33	120.0
C1—C2—H2	120.4	C32—C33—H33	120.0
C2—C3—C4	118.8 (8)	C33—C34—C35	121.6 (8)
C2—C3—H3	120.6	C33—C34—H34	119.2
C4—C3—H3	120.6	C35—C34—H34	119.2
C3—C4—C5	120.1 (8)	C34—C35—C36	119.9 (9)
C3—C4—H4	120.0	C34—C35—H35	120.1
C5—C4—H4	120.0	C36—C35—H35	120.1
N1—C5—C4	122.4 (7)	C31—C36—C35	120.0 (8)
N1—C5—C6	115.5 (6)	C31—C36—H36	120.0
C4—C5—C6	122.1 (7)	C35—C36—H36	120.0
N2—C6—C5	119.1 (6)	C42—C37—C38	116.0 (7)
N2—C6—H6	120.4	C42—C37—P2	119.4 (6)
C5—C6—H6	120.4	C38—C37—P2	124.4 (6)
C12—C7—C8	118.4 (7)	C37—C38—C39	122.0 (8)
C12—C7—P1	123.6 (6)	C37—C38—H38	119.0
C8—C7—P1	118.0 (5)	C39—C38—H38	119.0
C9—C8—C7	121.6 (8)	C40—C39—C38	119.3 (9)
C9—C8—H8	119.2	C40—C39—H39	120.4
C7—C8—H8	119.2	C38—C39—H39	120.4
C10—C9—C8	118.8 (8)	C41—C40—C39	119.9 (8)
C10—C9—H9	120.6	C41—C40—H40	120.1
C8—C9—H9	120.6	C39—C40—H40	120.1
C11—C10—C9	120.3 (8)	C40—C41—C42	119.4 (9)
C11—C10—H10	119.8	C40—C41—H41	120.3
C9—C10—H10	119.8	C42—C41—H41	120.3
C10—C11—C12	120.7 (8)	C37—C42—C41	123.3 (8)
C10—C11—H11	119.6	C37—C42—H42	118.4
C12—C11—H11	119.6	C41—C42—H42	118.4
C7—C12—C11	120.0 (7)	C12—C43—C11	119.5 (15)
C7—C12—H12	120.0	C12—C43—H43A	107.4
C11—C12—H12	120.0	C11—C43—H43A	107.4
C14—C13—C18	119.0 (7)	C12—C43—H43B	107.4
C14—C13—P1	123.1 (6)	C11—C43—H43B	107.4
C18—C13—P1	117.8 (5)	H43A—C43—H43B	107.0
C13—C14—C15	120.0 (8)	C14—C44—C13	106 (5)
C13—C14—H14	120.0	C14—C44—H44A	110.5
C15—C14—H14	120.0	C13—C44—H44A	110.5
C16—C15—C14	119.8 (9)	C14—C44—H44B	110.5
C16—C15—H15	120.1	C13—C44—H44B	110.5
C14—C15—H15	120.1	H44A—C44—H44B	108.7
C15—C16—C17	121.9 (9)	C16—C45—C15	116 (5)
C15—C16—H16	119.0	C16—C45—H45A	108.4
C17—C16—H16	119.0	C15—C45—H45A	108.4
C18—C17—C16	116.7 (8)	C16—C45—H45B	108.4
C18—C17—H17	121.7	C15—C45—H45B	108.4

supplementary materials

C16—C17—H17	121.7	H45A—C45—H45B	107.4
C17—C18—C13	122.6 (7)	C18—C46—Cl7	107 (7)
C17—C18—H18	118.7	C18—C46—H46A	110.2
C13—C18—H18	118.7	Cl7—C46—H46A	110.2
C24—C19—C20	116.9 (7)	Cl8—C46—H46B	110.2
C24—C19—P1	120.1 (5)	Cl7—C46—H46B	109.0
C20—C19—P1	123.0 (6)	H46A—C46—H46B	108.5
N2—Cu1—N1—C1	179.0 (7)	C18—C13—C14—C15	1.0 (12)
P1—Cu1—N1—C1	74.3 (6)	P1—C13—C14—C15	179.4 (7)
P2—Cu1—N1—C1	−70.2 (7)	C13—C14—C15—C16	−1.7 (14)
N2—Cu1—N1—C5	−2.5 (5)	C14—C15—C16—C17	2.3 (16)
P1—Cu1—N1—C5	−107.3 (5)	C15—C16—C17—C18	−2.3 (15)
P2—Cu1—N1—C5	108.3 (5)	C16—C17—C18—C13	1.6 (13)
N1—Cu1—N2—C6	2.1 (5)	C14—C13—C18—C17	−1.1 (12)
P1—Cu1—N2—C6	111.4 (5)	P1—C13—C18—C17	−179.6 (6)
P2—Cu1—N2—C6	−106.8 (5)	C13—P1—C19—C24	−79.3 (7)
N1—Cu1—N2—N2 ⁱ	−178.3 (8)	C7—P1—C19—C24	173.0 (6)
P1—Cu1—N2—N2 ⁱ	−69.1 (7)	Cu1—P1—C19—C24	45.7 (7)
P2—Cu1—N2—N2 ⁱ	72.8 (7)	C13—P1—C19—C20	96.9 (7)
N1—Cu1—P1—C13	159.0 (3)	C7—P1—C19—C20	−10.7 (7)
N2—Cu1—P1—C13	75.5 (3)	Cu1—P1—C19—C20	−138.0 (6)
P2—Cu1—P1—C13	−61.5 (3)	C24—C19—C20—C21	0.5 (12)
N1—Cu1—P1—C7	−81.9 (3)	P1—C19—C20—C21	−175.9 (6)
N2—Cu1—P1—C7	−165.4 (3)	C19—C20—C21—C22	0.7 (13)
P2—Cu1—P1—C7	57.6 (3)	C20—C21—C22—C23	−1.8 (13)
N1—Cu1—P1—C19	39.4 (3)	C21—C22—C23—C24	1.7 (13)
N2—Cu1—P1—C19	−44.0 (3)	C20—C19—C24—C23	−0.6 (12)
P2—Cu1—P1—C19	178.9 (3)	P1—C19—C24—C23	175.9 (7)
N1—Cu1—P2—C37	82.5 (3)	C22—C23—C24—C19	−0.5 (13)
N2—Cu1—P2—C37	168.2 (3)	C37—P2—C25—C30	15.0 (7)
P1—Cu1—P2—C37	−56.9 (3)	C31—P2—C25—C30	123.6 (7)
N1—Cu1—P2—C31	−40.3 (3)	Cu1—P2—C25—C30	−109.4 (6)
N2—Cu1—P2—C31	45.4 (3)	C37—P2—C25—C26	−169.6 (6)
P1—Cu1—P2—C31	−179.7 (3)	C31—P2—C25—C26	−61.0 (6)
N1—Cu1—P2—C25	−160.3 (3)	Cu1—P2—C25—C26	66.0 (6)
N2—Cu1—P2—C25	−74.7 (3)	C30—C25—C26—C27	2.5 (11)
P1—Cu1—P2—C25	60.3 (3)	P2—C25—C26—C27	−173.1 (6)
C5—N1—C1—C2	1.4 (12)	C25—C26—C27—C28	−1.4 (12)
Cu1—N1—C1—C2	179.8 (7)	C26—C27—C28—C29	−0.4 (13)
N1—C1—C2—C3	−3.8 (15)	C27—C28—C29—C30	1.1 (13)
C1—C2—C3—C4	4.7 (16)	C26—C25—C30—C29	−1.9 (11)
C2—C3—C4—C5	−3.4 (16)	P2—C25—C30—C29	173.4 (6)
C1—N1—C5—C4	−0.1 (11)	C28—C29—C30—C25	0.1 (13)
Cu1—N1—C5—C4	−178.7 (6)	C37—P2—C31—C36	−110.6 (7)
C1—N1—C5—C6	−178.7 (7)	C25—P2—C31—C36	143.2 (7)
Cu1—N1—C5—C6	2.6 (8)	Cu1—P2—C31—C36	17.3 (8)
C3—C4—C5—N1	1.1 (14)	C37—P2—C31—C32	67.5 (7)
C3—C4—C5—C6	179.6 (8)	C25—P2—C31—C32	−38.7 (7)

N2 ⁱ —N2—C6—C5	178.9 (6)	Cu1—P2—C31—C32	−164.6 (6)
Cu1—N2—C6—C5	−1.4 (8)	C36—C31—C32—C33	3.0 (13)
N1—C5—C6—N2	−0.7 (10)	P2—C31—C32—C33	−175.1 (7)
C4—C5—C6—N2	−179.4 (7)	C31—C32—C33—C34	−4.6 (14)
C13—P1—C7—C12	−6.9 (7)	C32—C33—C34—C35	3.3 (16)
C19—P1—C7—C12	99.9 (7)	C33—C34—C35—C36	−0.5 (16)
Cu1—P1—C7—C12	−131.2 (6)	C32—C31—C36—C35	−0.2 (13)
C13—P1—C7—C8	175.2 (6)	P2—C31—C36—C35	177.9 (7)
C19—P1—C7—C8	−78.0 (7)	C34—C35—C36—C31	−1.1 (15)
Cu1—P1—C7—C8	50.9 (7)	C31—P2—C37—C42	159.1 (7)
C12—C7—C8—C9	3.5 (13)	C25—P2—C37—C42	−93.7 (7)
P1—C7—C8—C9	−178.4 (7)	Cu1—P2—C37—C42	30.5 (8)
C7—C8—C9—C10	−2.1 (15)	C31—P2—C37—C38	−25.2 (9)
C8—C9—C10—C11	−0.8 (15)	C25—P2—C37—C38	82.0 (8)
C9—C10—C11—C12	2.0 (14)	Cu1—P2—C37—C38	−153.8 (7)
C8—C7—C12—C11	−2.2 (11)	C42—C37—C38—C39	2.4 (14)
P1—C7—C12—C11	179.9 (6)	P2—C37—C38—C39	−173.3 (8)
C10—C11—C12—C7	−0.5 (13)	C37—C38—C39—C40	−3.5 (16)
C7—P1—C13—C14	100.6 (7)	C38—C39—C40—C41	2.0 (16)
C19—P1—C13—C14	−6.4 (7)	C39—C40—C41—C42	0.4 (16)
Cu1—P1—C13—C14	−134.1 (6)	C38—C37—C42—C41	0.1 (14)
C7—P1—C13—C18	−81.0 (6)	P2—C37—C42—C41	176.1 (8)
C19—P1—C13—C18	172.0 (6)	C40—C41—C42—C37	−1.6 (16)
Cu1—P1—C13—C18	44.3 (6)		

Symmetry codes: (i) $-x+1, -y+1, -z+1$.

supplementary materials

Fig. 1

