

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

3-Hydroxy-3-[(2-methylpropanoyl)-methyl]indolin-2-one

Gang Chen,^{a*} Bin Liu,^b Ying Tang^a and Jingfang Xu^a

^aCollege of Chemistry and Chemical Engineering, Xi'an Shiyou University, Dianzi'er Road No. 18 Xi'an 710065, Xi'an, People's Republic of China, and ^bCollege of Environment and Chemical Engineering, Xi'an Polytechnic University, Nouth Jinhua Roud No. 19 Xi'an 710048, Xi'an, People's Republic of China
Correspondence e-mail: gangchen@xsyu.edu.cn


Received 26 November 2008; accepted 26 June 2009

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.045; wR factor = 0.161; data-to-parameter ratio = 16.6.

The title compound, $\text{C}_{13}\text{H}_{15}\text{NO}_3$, was synthesized by the Aldol reaction of isatin and 3-methylbutan-2-one refluxing in methanol catalyzed by dimethylamine. The packing of the molecules in the crystal structure features intermolecular $\text{N}\cdots\text{H}\cdots\text{O}$ and $\text{O}\cdots\text{H}\cdots\text{O}$ hydrogen bonds.

Related literature

For related structures, see: Garden *et al.* (2002); Li, *et al.* (2008). For the bioactivity of derivatives, see: Glover *et al.* (1988); Marti & Carreira (2003); Pandeya *et al.* (2000); Sun *et al.* (1998); Teitz *et al.* (1994).



Experimental

Crystal data

$\text{C}_{13}\text{H}_{15}\text{NO}_3$
 $M_r = 233.26$
Monoclinic, $P2_1/c$
 $a = 11.885$ (2) Å
 $b = 5.9244$ (12) Å
 $c = 16.695$ (3) Å
 $\beta = 98.60$ (3)°

$V = 1162.3$ (5) Å³
 $Z = 4$
Mo $K\alpha$ radiation
 $\mu = 0.10$ mm⁻¹
 $T = 293$ K
0.23 × 0.18 × 0.15 mm

Data collection

Bruker SMART CCD area-detector diffractometer
Absorption correction: multi-scan (SADABS; Sheldrick, 2005)
 $T_{\min} = 0.945$, $T_{\max} = 0.985$
8623 measured reflections
2577 independent reflections
1902 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.028$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.045$
 $wR(F^2) = 0.161$
 $S = 1.09$
2577 reflections
155 parameters
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.24$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.22$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D\cdots H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N1}-\text{H1A}\cdots\text{O2}^{\text{i}}$	0.86	2.22	3.0049 (18)	151
$\text{O2}-\text{H2A}\cdots\text{O1}^{\text{ii}}$	0.82	2.00	2.8220 (17)	175

Symmetry codes: (i) $x, y + 1, z$; (ii) $-x + 1, -y + 1, -z + 1$.

Data collection: SMART (Bruker, 2002); cell refinement: SAINT (Bruker, 2002); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 (Farrugia, 1997) and DIAMOND (Brandenburg, 1999); software used to prepare material for publication: WinGX (Farrugia, 1999).

This work was supported financially by grants from the National Science Foundation of China (No. 50874092) and the Scientific Research Plan Projects of Shaanxi Education Department (08 J K 413).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: WW2134).

References

- Brandenburg, K. (1999). DIAMOND. Crystal Impact GbR, Bonn, Germany.
Bruker (2002). SMART and SAINT for WNT/2000. Bruker AXS Inc., Madison, Wisconsin, USA.
Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
Garden, S. J., da Silva, R. B. & Pinto, A. C. (2002). *Tetrahedron*, **58**, 8399–8412.
Glover, V., Halket, J. M., Watkins, P. J., Clow, A., Goodwin, B. L. & Sandler, M. (1988). *J. Neurochem.* **51**, 656–659.
Li, Y. M., Zhang, Z. K., Jia, Y. T., Shen, Y. M., He, H. P., Fang, R. X., Chen, X. Y. & Hao, X. J. (2008). *Plant Biotechnol. J.* **6**, 301–308.
Marti, C. & Carreira, E. M. (2003). *Eur. J. Org. Chem.* pp. 2209–2219.
Pandeya, S. N., Sriram, D., Nath, G. & DeClercq, E. (2000). *Eur. J. Med. Chem.* **35**, 249–255.
Sheldrick, G. M. (2005). SADABS. University of Göttingen, Germany.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Sun, L., Tran, N., Tang, F., App, H., Hirth, P., McMahon, G. & Tang, C. (1998). *J. Med. Chem.* **41**, 2588–2603.
Teitz, Y., Ronen, D., Vansover, A., Stematsky, T. & Riggs, J. L. (1994). *Antivir. Res.* **24**, 305–314.

supplementary materials

Acta Cryst. (2009). E65, o1723 [doi:10.1107/S1600536809024611]

3-Hydroxy-3-[(2-methylpropanoyl)methyl]indolin-2-one

G. Chen, B. Liu, Y. Tang and J. Xu

Comment

Isatin, an endogenous compound in mammalian tissues and body fluids (Glover, *et al.*, 1988), has caught great attention of many researchers as a versatile lead molecule for designing of potential drugs (Pandeya, *et al.*, 2000; Sun, *et al.*, 1998; Teitz, *et al.*, 1994). In our previous study, 3-hydroxy-3-(2-oxo-propyl)-1,3-dihydro-indol-2-one, yielded from the Aldol reaction of isatin with acetone, was found to be a new chemical activator against tobacco mosaic virus (TMV) infection (Li, 2008). In tobacco plant, external application of 3-hydroxy-3-(2-oxo-propyl)-1,3-dihydro-indol-2-one results in restriction of TMV multiplication and spread, accumulation of salicylic acid level expression of PR-1 gene, and activation and increase of phenylalanine ammonia-lyase (PAL) activity. With these findings, some analogs need to be synthesized for structure activity relationship research to find more potent molecules.

One of these analogs, 3-hydroxy-3-(3-methyl-2-oxo-butyl)-1,3-dihydro-indol-2-one (compound I), was synthesized by the Aldol reaction of isatin with 3-methyl-butan-2-one. In order to provide the structural information of compound I, we studied its crystal structure. The title compound was synthesized in a one step Aldol reaction of isatin (0.01 mmol) with 3-methyl-butan-2-one, according to the reported method (Garden *et al.*, 2002). The molar ratio of isatin: 3-methyl-butan-2-one = 1:3 refluxed in methanol gave compound I in 72% yield, and colorless crystals of compound I were obtained in ethanol by recrystallization. The values of the geometric parameters of compound I are within normal ranges and experimental errors.

The molecular structure of compound I is illustrated in Fig. 1. In the molecule, the 2-oxo-indole ring is planar, and the angle between hydroxyl group and 3-methyl-2-oxo-butyl group is 105.10 (12)°. The intermolecular interactions are primarily responsible for the formation of the crystal structure of compound I. Each molecule is fixed by four hydrogen bond of other three molecules. N1–H1 acts as a hydrogen bond donor, and O1 is a hydrogen bond acceptor. To O2–H2, it acts as both a hydrogen bond donor and a hydrogen bond acceptor, which connects with two molecules by O–H···O and O···H–N with the angle of 119.21 (9)°. There is no anticipated intramolecular hydrogen bond between O2–H2 and O3, and O3 is not involved in any hydrogen bond (Fig. 2).

Experimental

A mixture of isatin (0.01 mmol) and 3-methyl-butan-2-one (0.03 mmol) was refluxed in methanol (60 ml), catalyzed by a drop of dimethylamine, until the disappearance of the starting material, as evidenced by thin-layer chromatography. The solvent was removed *in vacuo* and the residue was separated by column chromatography (silica gel, petroleum ether/ethyl acetate = 5:1), giving the title compound I 0.168 g, yield 72%. ¹H-NMR (CDCl₃, 400 MHz): 8.5 (1H, s), 7.32 (1H, d, J = 7.2 Hz), 7.24 (1H, t, J = 7.6 Hz), 7.03 (1H, t, J = 7.2 Hz), 6.88 (1H, d, J = 16.8 Hz), 4.81 (1H, s), 3.25 (1H, d, J = 16.8 Hz), 3.17 (1H, d, J = 16.8 Hz), 2.56 (1H, q, J = 6.8 Hz), 1.05 (6H, dd, J = 29.2, 6.8 Hz); ¹³C-NMR (CDCl₃, 100 MHz): 213.9, 178.5, 140.6, 130.3, 129.9, 124.1, 123.1, 110.5, 74.7, 45.6, 41.9, 17.6; MS (EI) *m/z*: 219 (*M*⁺). 30 mg of compound I was dissolved in 30 ml methanol and the solution was kept at room temperature for 4 d, natural evaporation gave colorless single crystals of compound I suitable for X-ray analysis.

Refinement

All H atoms were positioned geometrically, with C–H = 0.93–0.98 Å, and refined using riding model, with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{carrier})$.

Figures




Fig. 1. An ORTEP-3 drawing of compound I, with the atom-numbering scheme and 30% probability displacement ellipsoids.




Fig. 2. Packing diagram of the title compound. Dashed lines indicate hydrogen bonds.

3-Hydroxy-3-[(2-methylpropanoyl)methyl]indolin-2-one

Crystal data

$\text{C}_{13}\text{H}_{15}\text{NO}_3$

$M_r = 233.26$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

$a = 11.885(2) \text{ \AA}$

$b = 5.9244(12) \text{ \AA}$

$c = 16.695(3) \text{ \AA}$

$\beta = 98.60(3)^\circ$

$V = 1162.3(5) \text{ \AA}^3$

$Z = 4$

$F_{000} = 496$

$D_x = 1.333 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 3113 reflections

$\theta = 1.9\text{--}27.4^\circ$

$\mu = 0.10 \text{ mm}^{-1}$

$T = 293 \text{ K}$

Block, colorless

$0.23 \times 0.18 \times 0.15 \text{ mm}$

Data collection

Bruker SMART CCD area-detector
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 293 \text{ K}$

2577 independent reflections

1902 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.028$

$\theta_{\text{max}} = 27.5^\circ$

φ and ω scans	$\theta_{\min} = 1.7^\circ$
Absorption correction: multi-scan (SADABS; Sheldrick, 2005)	$h = -15 \rightarrow 14$
$T_{\min} = 0.945$, $T_{\max} = 0.985$	$k = -7 \rightarrow 7$
8623 measured reflections	$l = -21 \rightarrow 15$

Refinement

Refinement on F^2	Hydrogen site location: inferred from neighbouring sites
Least-squares matrix: full	H-atom parameters constrained
$R[F^2 > 2\sigma(F^2)] = 0.045$	$w = 1/[\sigma^2(F_o^2) + (0.0562P)^2 + 0.2707P]$
$wR(F^2) = 0.161$	where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.09$	$(\Delta/\sigma)_{\max} < 0.001$
2577 reflections	$\Delta\rho_{\max} = 0.24 \text{ e } \text{\AA}^{-3}$
155 parameters	$\Delta\rho_{\min} = -0.22 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: SHELXL, $F_c^* = kFc[1 + 0.001xFc^2\lambda^3/\sin(2\theta)]^{-1/4}$
Secondary atom site location: difference Fourier map	Extinction coefficient: 0.018 (4)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O2	0.57929 (9)	0.14669 (16)	0.55228 (6)	0.0335 (3)
H2A	0.5183	0.2128	0.5411	0.050*
O1	0.62325 (9)	0.59992 (19)	0.48205 (6)	0.0368 (3)
N1	0.60709 (11)	0.6720 (2)	0.61556 (8)	0.0324 (3)
H1A	0.5830	0.8088	0.6103	0.039*
O3	0.87069 (11)	0.5445 (2)	0.61665 (9)	0.0542 (4)
C1	0.63221 (12)	0.5435 (2)	0.55343 (9)	0.0292 (4)
C8	0.66420 (12)	0.3373 (2)	0.67659 (9)	0.0278 (4)
C9	0.77543 (13)	0.2149 (3)	0.56337 (10)	0.0320 (4)
H9A	0.7868	0.0611	0.5829	0.038*
H9B	0.7679	0.2105	0.5047	0.038*

supplementary materials

C10	0.87990 (14)	0.3519 (3)	0.59554 (10)	0.0347 (4)
C7	0.62536 (13)	0.5539 (2)	0.69000 (9)	0.0289 (4)
C2	0.66509 (12)	0.3034 (2)	0.58702 (9)	0.0276 (4)
C3	0.69101 (13)	0.1899 (3)	0.74041 (9)	0.0322 (4)
H3A	0.7185	0.0462	0.7320	0.039*
C6	0.61006 (14)	0.6260 (3)	0.76617 (10)	0.0351 (4)
H6A	0.5832	0.7702	0.7746	0.042*
C4	0.67614 (14)	0.2602 (3)	0.81790 (10)	0.0383 (4)
H4A	0.6931	0.1621	0.8615	0.046*
C5	0.63660 (14)	0.4736 (3)	0.83018 (10)	0.0381 (4)
H5A	0.6273	0.5176	0.8823	0.046*
C11	0.99369 (15)	0.2365 (3)	0.59805 (12)	0.0500 (5)
H11A	0.9932	0.1539	0.5471	0.060*
C12	1.0100 (2)	0.0663 (4)	0.66754 (18)	0.0843 (9)
H12A	0.9472	-0.0375	0.6614	0.126*
H12B	1.0133	0.1446	0.7182	0.126*
H12C	1.0796	-0.0153	0.6667	0.126*
C13	1.09104 (19)	0.4058 (5)	0.60623 (16)	0.0752 (7)
H13A	1.0797	0.5089	0.5614	0.113*
H13B	1.1617	0.3271	0.6065	0.113*
H13C	1.0931	0.4880	0.6560	0.113*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O2	0.0286 (6)	0.0262 (6)	0.0431 (7)	-0.0024 (4)	-0.0035 (5)	-0.0012 (4)
O1	0.0390 (7)	0.0385 (7)	0.0323 (6)	0.0007 (5)	0.0031 (5)	0.0073 (5)
N1	0.0381 (8)	0.0242 (6)	0.0349 (8)	0.0044 (5)	0.0053 (6)	0.0016 (5)
O3	0.0401 (8)	0.0400 (8)	0.0817 (10)	-0.0055 (5)	0.0068 (7)	-0.0124 (7)
C1	0.0247 (8)	0.0269 (8)	0.0353 (9)	-0.0018 (6)	0.0021 (6)	0.0021 (6)
C8	0.0247 (8)	0.0271 (8)	0.0317 (8)	-0.0010 (6)	0.0044 (6)	0.0004 (6)
C9	0.0290 (8)	0.0312 (8)	0.0352 (8)	0.0019 (6)	0.0025 (6)	-0.0032 (6)
C10	0.0325 (9)	0.0390 (9)	0.0334 (9)	-0.0020 (7)	0.0074 (7)	-0.0007 (7)
C7	0.0252 (8)	0.0283 (8)	0.0332 (8)	-0.0003 (6)	0.0041 (6)	0.0025 (6)
C2	0.0263 (8)	0.0253 (8)	0.0300 (8)	-0.0016 (6)	0.0002 (6)	-0.0015 (6)
C3	0.0324 (9)	0.0277 (8)	0.0360 (9)	0.0022 (6)	0.0034 (7)	0.0036 (6)
C6	0.0346 (9)	0.0301 (9)	0.0415 (9)	0.0009 (6)	0.0085 (7)	-0.0054 (7)
C4	0.0377 (10)	0.0429 (10)	0.0333 (9)	-0.0001 (7)	0.0022 (7)	0.0066 (7)
C5	0.0340 (9)	0.0512 (11)	0.0298 (8)	-0.0023 (7)	0.0071 (6)	-0.0045 (7)
C11	0.0313 (10)	0.0610 (12)	0.0566 (12)	0.0022 (8)	0.0033 (8)	-0.0117 (10)
C12	0.0569 (15)	0.0593 (14)	0.127 (2)	0.0053 (11)	-0.0164 (14)	0.0254 (15)
C13	0.0335 (12)	0.1033 (19)	0.0886 (18)	-0.0095 (12)	0.0084 (11)	0.0082 (15)

Geometric parameters (\AA , $^\circ$)

O2—C2	1.4356 (16)	C3—C4	1.395 (2)
O2—H2A	0.8200	C3—H3A	0.9300
O1—C1	1.2270 (18)	C6—C5	1.399 (2)
N1—C1	1.355 (2)	C6—H6A	0.9300

N1—C7	1.4144 (19)	C4—C5	1.375 (3)
N1—H1A	0.8600	C4—H4A	0.9300
O3—C10	1.204 (2)	C5—H5A	0.9300
C1—C2	1.557 (2)	C11—C13	1.522 (3)
C8—C3	1.378 (2)	C11—C12	1.527 (3)
C8—C7	1.393 (2)	C11—H11A	0.9800
C8—C2	1.510 (2)	C12—H12A	0.9600
C9—C10	1.513 (2)	C12—H12B	0.9600
C9—C2	1.518 (2)	C12—H12C	0.9600
C9—H9A	0.9700	C13—H13A	0.9600
C9—H9B	0.9700	C13—H13B	0.9600
C10—C11	1.511 (2)	C13—H13C	0.9600
C7—C6	1.379 (2)		
C2—O2—H2A	109.5	C8—C3—H3A	120.6
C1—N1—C7	111.85 (12)	C4—C3—H3A	120.6
C1—N1—H1A	124.1	C7—C6—C5	117.29 (15)
C7—N1—H1A	124.1	C7—C6—H6A	121.4
O1—C1—N1	126.39 (14)	C5—C6—H6A	121.4
O1—C1—C2	125.40 (14)	C5—C4—C3	120.42 (15)
N1—C1—C2	108.04 (13)	C5—C4—H4A	119.8
C3—C8—C7	120.23 (14)	C3—C4—H4A	119.8
C3—C8—C2	130.26 (14)	C4—C5—C6	121.53 (15)
C7—C8—C2	109.47 (12)	C4—C5—H5A	119.2
C10—C9—C2	114.65 (13)	C6—C5—H5A	119.2
C10—C9—H9A	108.6	C13—C11—C10	111.69 (16)
C2—C9—H9A	108.6	C13—C11—C12	111.02 (18)
C10—C9—H9B	108.6	C10—C11—C12	109.35 (17)
C2—C9—H9B	108.6	C13—C11—H11A	108.2
H9A—C9—H9B	107.6	C10—C11—H11A	108.2
O3—C10—C11	122.77 (15)	C12—C11—H11A	108.2
O3—C10—C9	120.45 (15)	C11—C12—H12A	109.5
C11—C10—C9	116.77 (14)	C11—C12—H12B	109.5
C6—C7—C8	121.75 (14)	H12A—C12—H12B	109.5
C6—C7—N1	129.26 (14)	C11—C12—H12C	109.5
C8—C7—N1	108.99 (13)	H12A—C12—H12C	109.5
O2—C2—C8	112.10 (12)	H12B—C12—H12C	109.5
O2—C2—C9	105.10 (11)	C11—C13—H13A	109.5
C8—C2—C9	115.95 (12)	C11—C13—H13B	109.5
O2—C2—C1	108.68 (11)	H13A—C13—H13B	109.5
C8—C2—C1	101.33 (11)	C11—C13—H13C	109.5
C9—C2—C1	113.69 (13)	H13A—C13—H13C	109.5
C8—C3—C4	118.76 (15)	H13B—C13—H13C	109.5

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N1—H1A \cdots O2 ⁱ	0.86	2.22	3.0049 (18)	151
O2—H2A \cdots O1 ⁱⁱ	0.82	2.00	2.8220 (17)	175

supplementary materials

Symmetry codes: (i) $x, y+1, z$; (ii) $-x+1, -y+1, -z+1$.

Fig. 1




Fig. 2

