

## t-3-Ethyl-r-2,c-6-bis(4-methoxyphenyl)-1-nitrosopiperidin-4-one

T. Kavitha,<sup>a</sup> M. Thenmozhi,<sup>a</sup> S. Ponnuswamy,<sup>b</sup>  
P. Sakthivel<sup>b</sup> and M. N. Ponnuswamy<sup>a\*</sup>

<sup>a</sup>Centre of Advanced Study in Crystallography and Biophysics, University of Madras, Guindy Campus, Chennai 600 025, India, and <sup>b</sup>Department of Chemistry, Government Arts College (Autonomous), Coimbatore 641 018, Tamilnadu, India  
Correspondence e-mail: mnpsey2004@yahoo.com


Received 20 May 2009; accepted 28 June 2009

Key indicators: single-crystal X-ray study;  $T = 293\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$ ; R factor = 0.040; wR factor = 0.107; data-to-parameter ratio = 13.7.

In the title molecule,  $\text{C}_{21}\text{H}_{24}\text{N}_2\text{O}_4$ , the piperidine ring adopts a distorted boat conformation with the ethyl substituent in the axial position. The dihedral angle between the two benzene rings is  $70.25(9)^\circ$ . An intramolecular C—H···O interaction is observed. In the crystal, molecules are linked into a chain along the *c* axis by C—H···O hydrogen bonds and the chains are linked via weak C—H···π interactions.

### Related literature

For general background to 4-piperidones, see: Wang *et al.* (1992); Grishina *et al.* (1994). For ring conformational analysis, see: Cremer & Pople (1975).



### Experimental

#### Crystal data



$M_r = 368.42$

Orthorhombic,  $P2_12_12_1$   
*a* = 7.2742 (4) Å  
*b* = 15.8459 (7) Å  
*c* = 16.4800 (7) Å  
*V* = 1899.59 (16) Å<sup>3</sup>

*Z* = 4  
Mo  $K\alpha$  radiation  
 $\mu = 0.09\text{ mm}^{-1}$   
*T* = 293 K  
 $0.25 \times 0.20 \times 0.20\text{ mm}$

#### Data collection

Bruker Kappa APEXII CCD area-detector diffractometer  
Absorption correction: multi-scan (*SADABS*; Sheldrick, 2001)  
 $R_{\min} = 0.978$ ,  $T_{\max} = 0.982$

15051 measured reflections  
3334 independent reflections  
2550 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.026$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.040$   
 $wR(F^2) = 0.107$   
 $S = 1.02$   
3334 reflections

244 parameters  
H-atom parameters constrained  
 $\Delta\rho_{\max} = 0.20\text{ e \AA}^{-3}$   
 $\Delta\rho_{\min} = -0.17\text{ e \AA}^{-3}$

**Table 1**  
Hydrogen-bond geometry (Å, °).

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
C15—H15C···O1	0.96	2.56	3.163 (4)	121
C18—H18···O1 <sup>i</sup>	0.93	2.56	3.472 (3)	167
C23—H23B···Cg1 <sup>ii</sup>	0.96	2.89	3.718 (2)	144

Symmetry codes: (i)  $-x + \frac{5}{2}, -y, z + \frac{1}{2}$ ; (ii)  $x - \frac{1}{2}, -y + \frac{1}{2}, -z + 1$ . Cg1 is the centroid of the C16—C21 ring.

Data collection: *APEX2* (Bruker, 2004); cell refinement: *SAINT* (Bruker, 2004); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997); software used to prepare material for publication: *SHELXL97* and *PLATON* (Spek, 2009).

TK thanks Dr Babu Varghese, SAIF, IIT-Madras, Chennai, India, for his help with the data collection. SP thanks the UGC, India, for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CI2810).

### References

- Bruker (2004). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Cremer, D. & Pople, J. A. (1975). *J. Am. Chem. Soc.* **97**, 1354–1358.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Grishina, G. V., Gaidarov, E. L. & Zefirov, N. S. (1994). *Chem. Heterocycl. Compd.* **30**, 1401–1426.
- Sheldrick, G. M. (2001). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.
- Wang, C.-L. & Wuorola, M. A. (1992). *Org. Prep. Proceed. Int.* **24**, 585–621.

## **supplementary materials**

*Acta Cryst.* (2009). E65, o1765 [doi:10.1107/S1600536809024878]

### ***t*-3-Ethyl-*r*-2,*c*-6-bis(4-methoxyphenyl)-1-nitrosopiperidin-4-one**

**T. Kavitha, M. Thenmozhi, S. Ponnuswamy, P. Sakthivel and M. N. Ponnuswamy**

#### **Comment**

Piperidine derivatives, namely 4-piperidones, are synthetic intermediates in the preparation of various alkaloids and pharmaceutical products (Wang *et al.*, 1992; Grishina *et al.*, 1994).

The piperidine ring adopts a distorted boat conformation, with the ethyl substituent at C3 position in the axial orientation. The puckering parameters for the piperidine ring are  $q_2 = 0.591(2)$  Å,  $q_3 = 0.097(2)$  Å,  $Q_T = 0.599(2)$  Å and  $\phi_2 = 73.2(2)^\circ$  (Cremer & Pople, 1975). The dihedral angle between the two benzene rings is  $70.25(9)^\circ$ . The sum of the bond angles around N1 ( $359^\circ$ ) indicates  $sp^2$  hybridization. An intramolecular C—H···O interaction is observed.

The crystal structure is stabilized by intermolecular C—H···O hydrogen bonds (Table 1) which link the molecules into a chain along the *c* axis. The chains are linked via C—H···π interactions involving the C16-C21 ring.

#### **Experimental**

To a solution of *t*-3-ethyl-*r*-2,*c*-6-bis(4-methoxyphenyl)piperidin-4-one (1.69 g, 5 mmol) in chloroform (10 ml) was added conc. HCl (1.5 ml) and water (1.5 ml) and while stirring, solid NaNO<sub>2</sub> (0.84 g, 12 mmol) was added in portions during 0.5 h. The solution was stirred at room temperature for another 0.5 h. The organic layer was washed with water, saturated aqueous NaHCO<sub>3</sub> and dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>. The resulting solution was concentrated and the residue was crystallized from ethanol.

#### **Refinement**

H atoms were positioned geometrically (C—H = 0.93–0.98 Å) and allowed to ride on their parent atoms, with  $U_{\text{iso}}(\text{H}) = 1.2\text{--}1.5U_{\text{eq}}(\text{C})$ . In the absence of significant anomalous scattering effects, the Friedel pairs were averaged.

#### **Figures**




Fig. 1. Molecular structure of the title compound. Displacement ellipsoids are drawn at the 30% probability level. H atoms have been omitted for clarity.

## supplementary materials

---




Fig. 2. The molecular packing of the title compound, viewed approximately along the  $a$  axis. Dashed lines indicate hydrogen bonds. H atoms not involved in hydrogen bonding have been omitted.

### t-3-Ethyl-*r*-2,6-bis(4-methoxyphenyl)-1-nitrosopiperidin-4-one

#### Crystal data

C <sub>21</sub> H <sub>24</sub> N <sub>2</sub> O <sub>4</sub>	$F_{000} = 784$
$M_r = 368.42$	$D_x = 1.288 \text{ Mg m}^{-3}$
Orthorhombic, $P2_12_12_1$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: P 2ac 2ab	Cell parameters from 3334 reflections
$a = 7.2742 (4) \text{ \AA}$	$\theta = 2.5\text{--}31.8^\circ$
$b = 15.8459 (7) \text{ \AA}$	$\mu = 0.09 \text{ mm}^{-1}$
$c = 16.4800 (7) \text{ \AA}$	$T = 293 \text{ K}$
$V = 1899.59 (16) \text{ \AA}^3$	Block, colourless
$Z = 4$	$0.25 \times 0.20 \times 0.20 \text{ mm}$

#### Data collection

Bruker Kappa APEXII CCD area-detector diffractometer	3334 independent reflections
Radiation source: fine-focus sealed tube	2550 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.026$
$T = 293 \text{ K}$	$\theta_{\text{max}} = 31.8^\circ$
$\omega$ scans	$\theta_{\text{min}} = 2.5^\circ$
Absorption correction: multi-scan (SADABS; Sheldrick, 2001)	$h = -10 \rightarrow 10$
$T_{\text{min}} = 0.978$ , $T_{\text{max}} = 0.982$	$k = -23 \rightarrow 16$
15051 measured reflections	$l = -15 \rightarrow 24$

#### Refinement

Refinement on $F^2$	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.040$	H-atom parameters constrained
$wR(F^2) = 0.107$	$w = 1/[\sigma^2(F_o^2) + (0.0555P)^2 + 0.1347P]$ where $P = (F_o^2 + 2F_c^2)/3$

$S = 1.02$	$(\Delta/\sigma)_{\max} = 0.001$
3334 reflections	$\Delta\rho_{\max} = 0.20 \text{ e \AA}^{-3}$
244 parameters	$\Delta\rho_{\min} = -0.16 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

### Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	1.2631 (2)	0.06968 (12)	0.09708 (10)	0.0722 (5)
O2	0.56881 (19)	0.08201 (12)	0.27581 (11)	0.0648 (4)
O3	1.2829 (2)	-0.26039 (11)	0.42094 (11)	0.0677 (5)
O4	0.96286 (19)	0.17870 (9)	0.59490 (8)	0.0480 (3)
N1	0.82220 (18)	0.02169 (10)	0.24151 (9)	0.0379 (3)
C2	0.9060 (2)	-0.04979 (11)	0.19832 (11)	0.0394 (4)
H2	0.8044	-0.0852	0.1792	0.047*
C3	1.0038 (3)	-0.01651 (12)	0.12220 (12)	0.0431 (4)
H3	1.0693	-0.0633	0.0962	0.052*
C4	1.1403 (3)	0.05176 (12)	0.14271 (12)	0.0439 (4)
C5	1.1176 (2)	0.09628 (12)	0.22290 (12)	0.0406 (4)
H5A	1.1624	0.1536	0.2166	0.049*
H5B	1.1967	0.0686	0.2621	0.049*
C6	0.9234 (2)	0.10068 (11)	0.25901 (11)	0.0361 (4)
H6	0.8583	0.1469	0.2319	0.043*
N7	0.6402 (2)	0.01792 (13)	0.24877 (11)	0.0523 (4)
C8	1.0192 (2)	-0.10403 (10)	0.25549 (12)	0.0383 (4)
C9	1.2088 (3)	-0.11160 (12)	0.25259 (13)	0.0452 (4)
H9	1.2747	-0.0822	0.2134	0.054*
C10	1.3015 (3)	-0.16248 (13)	0.30744 (13)	0.0496 (5)
H10	1.4290	-0.1664	0.3051	0.060*
C11	1.2061 (3)	-0.20718 (12)	0.36542 (13)	0.0484 (5)
C12	1.0155 (3)	-0.20026 (13)	0.36894 (14)	0.0516 (5)
H12	0.9496	-0.2303	0.4077	0.062*
C13	0.9252 (3)	-0.14900 (12)	0.31510 (13)	0.0473 (5)
H13	0.7980	-0.1441	0.3184	0.057*
C16	0.9304 (2)	0.12116 (10)	0.34877 (11)	0.0351 (4)

## supplementary materials

---

C17	1.0182 (2)	0.06863 (11)	0.40442 (12)	0.0407 (4)
H17	1.0712	0.0186	0.3865	0.049*
C18	1.0279 (2)	0.08954 (12)	0.48551 (11)	0.0411 (4)
H18	1.0871	0.0537	0.5218	0.049*
C19	0.9491 (2)	0.16428 (11)	0.51316 (11)	0.0373 (4)
C20	0.8647 (3)	0.21793 (12)	0.45857 (11)	0.0415 (4)
H20	0.8142	0.2686	0.4762	0.050*
C21	0.8557 (2)	0.19574 (11)	0.37750 (12)	0.0402 (4)
H21	0.7977	0.2320	0.3412	0.048*
C22	1.4718 (4)	-0.28172 (19)	0.41023 (18)	0.0810 (8)
H22A	1.5095	-0.3196	0.4525	0.122*
H22B	1.5451	-0.2314	0.4125	0.122*
H22C	1.4881	-0.3085	0.3585	0.122*
C23	0.8815 (3)	0.25378 (15)	0.62544 (14)	0.0602 (6)
H23A	0.8998	0.2567	0.6831	0.090*
H23B	0.7521	0.2535	0.6139	0.090*
H23C	0.9375	0.3019	0.6000	0.090*
C14	0.8639 (4)	0.02004 (17)	0.06142 (14)	0.0644 (6)
H14A	0.7562	-0.0160	0.0604	0.077*
H14B	0.8255	0.0753	0.0802	0.077*
C15	0.9381 (5)	0.0281 (2)	-0.02394 (16)	0.0977 (10)
H15A	0.8446	0.0512	-0.0586	0.147*
H15B	0.9737	-0.0265	-0.0436	0.147*
H15C	1.0430	0.0649	-0.0238	0.147*

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
O1	0.0685 (10)	0.0868 (12)	0.0614 (10)	-0.0296 (9)	0.0256 (9)	-0.0182 (9)
O2	0.0365 (7)	0.0822 (10)	0.0758 (11)	0.0125 (7)	-0.0027 (7)	-0.0142 (10)
O3	0.0698 (10)	0.0673 (10)	0.0660 (10)	0.0246 (8)	0.0030 (9)	0.0130 (9)
O4	0.0508 (8)	0.0563 (8)	0.0369 (7)	0.0020 (6)	-0.0037 (6)	-0.0045 (6)
N1	0.0286 (6)	0.0435 (7)	0.0416 (8)	-0.0018 (6)	-0.0024 (6)	-0.0043 (7)
C2	0.0366 (8)	0.0379 (8)	0.0436 (10)	-0.0058 (7)	-0.0018 (7)	-0.0076 (8)
C3	0.0489 (10)	0.0419 (8)	0.0386 (10)	-0.0050 (8)	0.0007 (8)	-0.0061 (8)
C4	0.0426 (9)	0.0461 (9)	0.0429 (10)	-0.0062 (8)	0.0033 (8)	-0.0022 (9)
C5	0.0368 (8)	0.0413 (9)	0.0437 (10)	-0.0063 (7)	0.0010 (7)	-0.0054 (8)
C6	0.0340 (7)	0.0357 (8)	0.0384 (10)	0.0006 (6)	-0.0017 (7)	-0.0023 (8)
N7	0.0310 (7)	0.0675 (11)	0.0585 (11)	-0.0007 (8)	-0.0035 (7)	-0.0043 (9)
C8	0.0381 (8)	0.0340 (8)	0.0427 (10)	-0.0032 (6)	0.0031 (7)	-0.0075 (8)
C9	0.0404 (9)	0.0447 (10)	0.0505 (11)	-0.0009 (7)	0.0062 (8)	-0.0027 (9)
C10	0.0394 (9)	0.0521 (10)	0.0574 (13)	0.0091 (8)	0.0038 (9)	-0.0057 (10)
C11	0.0527 (11)	0.0431 (9)	0.0495 (12)	0.0111 (8)	-0.0008 (9)	-0.0020 (10)
C12	0.0532 (11)	0.0494 (10)	0.0521 (12)	-0.0015 (9)	0.0085 (10)	0.0030 (10)
C13	0.0387 (9)	0.0458 (10)	0.0573 (13)	-0.0034 (8)	0.0039 (9)	-0.0017 (10)
C16	0.0296 (7)	0.0377 (8)	0.0381 (9)	0.0008 (6)	-0.0025 (6)	-0.0011 (8)
C17	0.0403 (9)	0.0364 (8)	0.0455 (10)	0.0067 (7)	-0.0038 (8)	-0.0025 (8)
C18	0.0395 (9)	0.0412 (9)	0.0426 (10)	0.0027 (7)	-0.0063 (7)	0.0049 (8)

C19	0.0309 (7)	0.0441 (9)	0.0369 (9)	-0.0032 (7)	-0.0008 (7)	-0.0033 (8)
C20	0.0413 (9)	0.0393 (8)	0.0439 (10)	0.0085 (7)	-0.0012 (8)	-0.0052 (8)
C21	0.0381 (9)	0.0403 (8)	0.0422 (10)	0.0093 (7)	-0.0046 (7)	0.0011 (8)
C22	0.0746 (17)	0.0949 (19)	0.0736 (17)	0.0407 (15)	-0.0056 (14)	0.0020 (16)
C23	0.0586 (12)	0.0756 (14)	0.0465 (12)	0.0077 (11)	0.0004 (10)	-0.0187 (12)
C14	0.0659 (13)	0.0755 (15)	0.0517 (13)	-0.0138 (12)	-0.0142 (11)	0.0046 (12)
C15	0.117 (3)	0.129 (3)	0.0469 (15)	-0.005 (2)	-0.0102 (16)	0.0193 (18)

*Geometric parameters ( $\text{\AA}$ ,  $^\circ$ )*

O1—C4	1.202 (2)	C11—C12	1.392 (3)
O2—N7	1.225 (2)	C12—C13	1.370 (3)
O3—C11	1.364 (3)	C12—H12	0.93
O3—C22	1.426 (3)	C13—H13	0.93
O4—C19	1.370 (2)	C16—C21	1.384 (2)
O4—C23	1.421 (2)	C16—C17	1.394 (2)
N1—N7	1.331 (2)	C17—C18	1.379 (3)
N1—C2	1.470 (2)	C17—H17	0.93
N1—C6	1.481 (2)	C18—C19	1.392 (3)
C2—C8	1.518 (3)	C18—H18	0.93
C2—C3	1.536 (3)	C19—C20	1.382 (3)
C2—H2	0.98	C20—C21	1.383 (3)
C3—C4	1.506 (3)	C20—H20	0.93
C3—C14	1.541 (3)	C21—H21	0.93
C3—H3	0.98	C22—H22A	0.96
C4—C5	1.507 (3)	C22—H22B	0.96
C5—C6	1.535 (2)	C22—H22C	0.96
C5—H5A	0.97	C23—H23A	0.96
C5—H5B	0.97	C23—H23B	0.96
C6—C16	1.515 (3)	C23—H23C	0.96
C6—H6	0.98	C14—C15	1.512 (4)
C8—C9	1.385 (3)	C14—H14A	0.97
C8—C13	1.393 (3)	C14—H14B	0.97
C9—C10	1.386 (3)	C15—H15A	0.96
C9—H9	0.93	C15—H15B	0.96
C10—C11	1.377 (3)	C15—H15C	0.96
C10—H10	0.93		
C11—O3—C22	117.3 (2)	C11—C12—H12	120.1
C19—O4—C23	117.23 (16)	C12—C13—C8	121.65 (18)
N7—N1—C2	114.93 (15)	C12—C13—H13	119.2
N7—N1—C6	121.01 (16)	C8—C13—H13	119.2
C2—N1—C6	122.65 (13)	C21—C16—C17	117.70 (17)
N1—C2—C8	111.15 (14)	C21—C16—C6	120.23 (15)
N1—C2—C3	108.85 (14)	C17—C16—C6	122.00 (15)
C8—C2—C3	116.74 (15)	C18—C17—C16	121.19 (16)
N1—C2—H2	106.5	C18—C17—H17	119.4
C8—C2—H2	106.5	C16—C17—H17	119.4
C3—C2—H2	106.5	C17—C18—C19	120.04 (17)
C4—C3—C2	111.62 (15)	C17—C18—H18	120.0

## supplementary materials

---

C4—C3—C14	108.13 (17)	C19—C18—H18	120.0
C2—C3—C14	110.74 (17)	O4—C19—C20	124.75 (17)
C4—C3—H3	108.8	O4—C19—C18	115.70 (16)
C2—C3—H3	108.8	C20—C19—C18	119.55 (17)
C14—C3—H3	108.8	C19—C20—C21	119.58 (16)
O1—C4—C3	121.27 (18)	C19—C20—H20	120.2
O1—C4—C5	121.28 (17)	C21—C20—H20	120.2
C3—C4—C5	117.44 (16)	C20—C21—C16	121.92 (17)
C4—C5—C6	117.51 (15)	C20—C21—H21	119.0
C4—C5—H5A	107.9	C16—C21—H21	119.0
C6—C5—H5A	107.9	O3—C22—H22A	109.5
C4—C5—H5B	107.9	O3—C22—H22B	109.5
C6—C5—H5B	107.9	H22A—C22—H22B	109.5
H5A—C5—H5B	107.2	O3—C22—H22C	109.5
N1—C6—C16	112.83 (15)	H22A—C22—H22C	109.5
N1—C6—C5	110.11 (14)	H22B—C22—H22C	109.5
C16—C6—C5	110.94 (14)	O4—C23—H23A	109.5
N1—C6—H6	107.6	O4—C23—H23B	109.5
C16—C6—H6	107.6	H23A—C23—H23B	109.5
C5—C6—H6	107.6	O4—C23—H23C	109.5
O2—N7—N1	114.66 (17)	H23A—C23—H23C	109.5
C9—C8—C13	117.96 (18)	H23B—C23—H23C	109.5
C9—C8—C2	124.60 (17)	C15—C14—C3	113.7 (2)
C13—C8—C2	117.44 (16)	C15—C14—H14A	108.8
C8—C9—C10	120.83 (19)	C3—C14—H14A	108.8
C8—C9—H9	119.6	C15—C14—H14B	108.8
C10—C9—H9	119.6	C3—C14—H14B	108.8
C11—C10—C9	120.40 (18)	H14A—C14—H14B	107.7
C11—C10—H10	119.8	C14—C15—H15A	109.5
C9—C10—H10	119.8	C14—C15—H15B	109.5
O3—C11—C10	125.24 (18)	H15A—C15—H15B	109.5
O3—C11—C12	115.4 (2)	C14—C15—H15C	109.5
C10—C11—C12	119.37 (19)	H15A—C15—H15C	109.5
C13—C12—C11	119.8 (2)	H15B—C15—H15C	109.5
C13—C12—H12	120.1		
N7—N1—C2—C8	-111.25 (18)	C8—C9—C10—C11	0.7 (3)
C6—N1—C2—C8	82.2 (2)	C22—O3—C11—C10	-10.4 (3)
N7—N1—C2—C3	118.82 (18)	C22—O3—C11—C12	168.6 (2)
C6—N1—C2—C3	-47.7 (2)	C9—C10—C11—O3	178.42 (19)
N1—C2—C3—C4	54.76 (19)	C9—C10—C11—C12	-0.6 (3)
C8—C2—C3—C4	-72.04 (19)	O3—C11—C12—C13	-179.33 (18)
N1—C2—C3—C14	-65.78 (19)	C10—C11—C12—C13	-0.2 (3)
C8—C2—C3—C14	167.43 (16)	C11—C12—C13—C8	1.0 (3)
C2—C3—C4—O1	159.7 (2)	C9—C8—C13—C12	-0.9 (3)
C14—C3—C4—O1	-78.2 (3)	C2—C8—C13—C12	179.36 (17)
C2—C3—C4—C5	-19.8 (2)	N1—C6—C16—C21	-119.97 (17)
C14—C3—C4—C5	102.2 (2)	C5—C6—C16—C21	115.94 (18)
O1—C4—C5—C6	153.2 (2)	N1—C6—C16—C17	63.0 (2)
C3—C4—C5—C6	-27.2 (2)	C5—C6—C16—C17	-61.1 (2)

N7—N1—C6—C16	72.0 (2)	C21—C16—C17—C18	1.0 (3)
C2—N1—C6—C16	-122.28 (17)	C6—C16—C17—C18	178.05 (17)
N7—N1—C6—C5	-163.49 (17)	C16—C17—C18—C19	0.0 (3)
C2—N1—C6—C5	2.3 (2)	C23—O4—C19—C20	1.0 (3)
C4—C5—C6—N1	36.3 (2)	C23—O4—C19—C18	-179.06 (16)
C4—C5—C6—C16	161.94 (15)	C17—C18—C19—O4	178.80 (17)
C2—N1—N7—O2	-171.53 (17)	C17—C18—C19—C20	-1.3 (3)
C6—N1—N7—O2	-4.7 (3)	O4—C19—C20—C21	-178.57 (18)
N1—C2—C8—C9	-112.31 (19)	C18—C19—C20—C21	1.5 (3)
C3—C2—C8—C9	13.3 (3)	C19—C20—C21—C16	-0.5 (3)
N1—C2—C8—C13	67.46 (19)	C17—C16—C21—C20	-0.7 (3)
C3—C2—C8—C13	-166.89 (16)	C6—C16—C21—C20	-177.86 (17)
C13—C8—C9—C10	0.0 (3)	C4—C3—C14—C15	75.9 (3)
C2—C8—C9—C10	179.80 (17)	C2—C3—C14—C15	-161.5 (2)

*Hydrogen-bond geometry (Å, °)*

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
C15—H15C···O1	0.96	2.56	3.163 (4)	121
C18—H18···O1 <sup>i</sup>	0.93	2.56	3.472 (3)	167
C23—H23B···Cg1 <sup>ii</sup>	0.96	2.89	3.718 (2)	144

Symmetry codes: (i)  $-x+5/2, -y, z+1/2$ ; (ii)  $x-1/2, -y+1/2, -z+1$ .

## supplementary materials

---

Fig. 1




Fig. 2

