

2,3-Dimethyl-6-nitro-2H-indazole

Yan Chen,^a Zheng Fang^{b*} and Ping Wei^a

^aCollege of Biotechnology and Pharmaceutical Engineering, Nanjing University of Technology, Xinmofan Road No. 5 Nanjing, Nanjing 210009, People's Republic of China, and ^bSchool of Pharmaceutical Sciences, Nanjing University of Technology, Xinmofan Road No. 5 Nanjing, Nanjing 210009, People's Republic of China
Correspondence e-mail: fzcpu@163.com


Received 26 June 2009; accepted 1 July 2009

Key indicators: single-crystal X-ray study; $T = 294\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.054; wR factor = 0.154; data-to-parameter ratio = 12.4.

In the molecule of the title compound, $\text{C}_9\text{H}_9\text{N}_3\text{O}_2$, the indazole ring system is almost planar [maximum deviation = 0.019 (3) Å for the C atom bearing the nitro group]. In the crystal structure, intermolecular C–H···O interactions link the molecules into centrosymmetric dimers, forming $R_2^2(18)$ ring motifs. Aromatic π – π contacts between indazole rings [centroid–centroid distances = 3.632 (1) and 3.705 (1) Å] may further stabilize the structure.

Related literature

For a related structure, see: Xu *et al.* (1999). For bond-length data, see: Allen *et al.* (1987). For ring-motifs, see: Bernstein *et al.* (1995).



Experimental

Crystal data

$\text{C}_9\text{H}_9\text{N}_3\text{O}_2$
 $M_r = 191.19$
Triclinic, $P\bar{1}$
 $a = 6.5800 (13)\text{ \AA}$

$b = 7.2050 (14)\text{ \AA}$
 $c = 10.752 (2)\text{ \AA}$
 $\alpha = 75.07 (3)^\circ$
 $\beta = 74.67 (3)^\circ$

$\gamma = 66.73 (3)^\circ$
 $V = 444.81 (19)\text{ \AA}^3$
 $Z = 2$
Mo $K\alpha$ radiation

$\mu = 0.11\text{ mm}^{-1}$
 $T = 294\text{ K}$
 $0.30 \times 0.20 \times 0.10\text{ mm}$

Data collection

Enraf–Nonius CAD-4 diffractometer
Absorption correction: ψ scan (North *et al.*, 1968)
 $T_{\min} = 0.969$, $T_{\max} = 0.990$
1756 measured reflections

1606 independent reflections
1292 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.031$
3 standard reflections
frequency: 120 min
intensity decay: 1%

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.054$
 $wR(F^2) = 0.154$
 $S = 1.00$
1606 reflections

129 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.32\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.25\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C1–H1A···O2 ⁱ	0.96	2.58	3.533 (4)	171

Symmetry code: (i) $-x - 1, -y + 2, -z$.

Data collection: *CAD-4 Software* (Enraf–Nonius, 1989); cell refinement: *CAD-4 Software*; data reduction: *XCAD4* (Harms & Wocadlo, 1995); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997) and *PLATON* (Spek, 2009); software used to prepare material for publication: *SHELXL97* and *PLATON*.

The authors thank the Center of Testing and Analysis, Nanjing University, for support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HK2724).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Enraf–Nonius (1989). *CAD-4 Software*. Enraf–Nonius, Delft, The Netherlands.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Harms, K. & Wocadlo, S. (1995). *XCAD4*. University of Marburg, Germany.
- North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst. A* **24**, 351–359.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.
- Xu, B.-C., Deng, F. & Wang, H.-Z. (1999). *Speciality Petro. Chem.* pp. 18–20.

supplementary materials

Acta Cryst. (2009). E65, o1775 [doi:10.1107/S1600536809025410]

2,3-Dimethyl-6-nitro-2H-indazole

Y. Chen, Z. Fang and P. Wei

Comment

Some derivatives of indazole are important chemical materials. We report herein the crystal structure of the title compound.

In the molecule of the title compound (Fig 1), the bond lengths (Allen *et al.*, 1987) and angles are within normal ranges. Rings A (N1/N2/C3/C4/C9) and B (C4-C9) are, of course, planar and the dihedral angle between them is A/B = 0.80 (3) $^{\circ}$. The indazole ring system is planar with a maximum deviation of -0.019 (3) Å for atom C6. Atoms O1, O2, N3, C1 and C2 are 0.024 (3), -0.124 (3), -0.038 (3), 0.003 (3) and -0.056 (3) Å away from the plane of the indazole ring system, respectively.

In the crystal structure, weak intermolecular C-H···O interactions (Table 1) link the molecules into centrosymmetric dimers forming $R_2^2(18)$ ring motifs (Bernstein *et al.*, 1995) (Fig. 2), in which they may be effective in the stabilization of the structure. The π – π contacts between the indazole rings, Cg1—Cg2ⁱ and Cg2—Cg2ⁱⁱ [symmetry codes: (i) 2 - x, 2 - y, -z, (ii) 2 - x, 1 - y, -z, where Cg1 and Cg2 are centroids of the rings A (N1/N2/C3/C4/C9) and B (C4-C9), respectively] may further stabilize the structure, with centroid-centroid distances of 3.632 (1) and 3.705 (1) Å, respectively.

Experimental

For the preparation of the title compound, metallic sodium (3.22 g) was dissolved in regurgitant 2-propanol (140 ml). Then, the solution was added to 3-methyl-6-nitro-1*H*-indazole (13 g) and iodomethane (30 g) was added in small portions. The mixture was refluxed for 5 h. The suspension was cooled to room temperature, filtered and washed with 2-propanol to give yellow solid (yield; 12 g) (Xu *et al.*, 1999). Crystals suitable for X-ray analysis were obtained by slow evaporation of a methanol solution.

Refinement

H atoms were positioned geometrically, with C-H = 0.93 and 0.96 Å for aromatic and methyl H, respectively, and constrained to ride on their parent atoms, with $U_{\text{iso}}(\text{H}) = xU_{\text{eq}}(\text{C})$, where $x = 1.5$ for methyl H and $x = 1.2$ for aromatic H atoms.

Figures




Fig. 1. The molecular structure of the title molecule, with the atom-numbering scheme. Displacement ellipsoids are drawn at the 30% probability level.

supplementary materials




Fig. 2. A partial packing diagram of the title compound. Hydrogen bonds are shown as dashed lines.

2,3-Dimethyl-6-nitro-2*H*-indazole

Crystal data

C ₉ H ₉ N ₃ O ₂	Z = 2
M _r = 191.19	F ₀₀₀ = 200
Triclinic, P <bar{1}< td=""><td>D_x = 1.427 Mg m⁻³</td></bar{1}<>	D _x = 1.427 Mg m ⁻³
Hall symbol: -P 1	Mo K α radiation, λ = 0.71073 Å
a = 6.5800 (13) Å	Cell parameters from 25 reflections
b = 7.2050 (14) Å	θ = 9–13°
c = 10.752 (2) Å	μ = 0.11 mm ⁻¹
α = 75.07 (3)°	T = 294 K
β = 74.67 (3)°	Block, colorless
γ = 66.73 (3)°	0.30 × 0.20 × 0.10 mm
V = 444.81 (19) Å ³	

Data collection

Enraf–Nonius CAD-4 diffractometer	R _{int} = 0.031
Radiation source: fine-focus sealed tube	$\theta_{\text{max}} = 25.3^\circ$
Monochromator: graphite	$\theta_{\text{min}} = 2.0^\circ$
T = 294 K	$h = 0 \rightarrow 7$
$\omega/2\theta$ scans	$k = -7 \rightarrow 8$
Absorption correction: ψ scan (North <i>et al.</i> , 1968)	$l = -12 \rightarrow 12$
$T_{\text{min}} = 0.969$, $T_{\text{max}} = 0.990$	3 standard reflections
1756 measured reflections	every 120 min
1606 independent reflections	intensity decay: 1%
1292 reflections with $I > 2\sigma(I)$	

Refinement

Refinement on F^2	Hydrogen site location: inferred from neighbouring sites
Least-squares matrix: full	H-atom parameters constrained
$R[F^2 > 2\sigma(F^2)] = 0.054$	$w = 1/[\sigma^2(F_o^2) + (0.08P)^2 + 0.235P]$ where $P = (F_o^2 + 2F_c^2)/3$

$wR(F^2) = 0.154$	$(\Delta/\sigma)_{\max} < 0.001$
$S = 1.00$	$\Delta\rho_{\max} = 0.32 \text{ e \AA}^{-3}$
1606 reflections	$\Delta\rho_{\min} = -0.25 \text{ e \AA}^{-3}$
129 parameters	Extinction correction: SHELXL97 (Sheldrick, 2008), $F_c^* = kF_c[1 + 0.001x F_c^2 \lambda^3 / \sin(2\theta)]^{1/4}$
Primary atom site location: structure-invariant direct methods	Extinction coefficient: 0.059 (12)
Secondary atom site location: difference Fourier map	

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.2155 (5)	0.7495 (5)	-0.3831 (2)	0.1020 (9)
O2	-0.1339 (4)	0.7845 (4)	-0.33747 (19)	0.0803 (7)
N1	-0.2315 (3)	0.7731 (3)	0.25319 (18)	0.0469 (5)
N2	-0.3309 (3)	0.7997 (3)	0.15107 (18)	0.0479 (5)
N3	0.0392 (4)	0.7648 (3)	-0.3051 (2)	0.0611 (6)
C1	-0.3707 (5)	0.7880 (5)	0.3829 (2)	0.0660 (8)
H1A	-0.5144	0.8941	0.3758	0.099*
H1B	-0.2979	0.8202	0.4368	0.099*
H1C	-0.3915	0.6594	0.4218	0.099*
C2	0.1367 (5)	0.6977 (4)	0.3176 (3)	0.0601 (7)
H2B	0.0456	0.7087	0.4032	0.090*
H2C	0.2098	0.7971	0.2929	0.090*
H2D	0.2482	0.5625	0.3186	0.090*
C3	-0.0080 (4)	0.7362 (3)	0.2216 (2)	0.0432 (6)
C4	-0.1582 (3)	0.7777 (3)	0.0489 (2)	0.0390 (5)
C5	-0.1637 (4)	0.7888 (3)	-0.0830 (2)	0.0446 (6)
H5A	-0.2966	0.8160	-0.1108	0.054*
C6	0.0376 (4)	0.7573 (3)	-0.1678 (2)	0.0462 (6)
C7	0.2431 (4)	0.7224 (4)	-0.1315 (2)	0.0517 (6)
H7A	0.3741	0.7057	-0.1943	0.062*
C8	0.2482 (4)	0.7134 (3)	-0.0049 (2)	0.0470 (6)
H8A	0.3819	0.6910	0.0204	0.056*
C9	0.0474 (3)	0.7388 (3)	0.0874 (2)	0.0387 (5)

supplementary materials

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.1021 (19)	0.150 (2)	0.0491 (12)	-0.0512 (17)	0.0174 (12)	-0.0311 (13)
O2	0.1002 (17)	0.0986 (16)	0.0521 (12)	-0.0371 (13)	-0.0217 (11)	-0.0183 (11)
N1	0.0480 (11)	0.0573 (12)	0.0389 (10)	-0.0226 (9)	-0.0063 (8)	-0.0089 (8)
N2	0.0436 (11)	0.0606 (12)	0.0427 (11)	-0.0218 (9)	-0.0069 (8)	-0.0098 (9)
N3	0.0804 (16)	0.0572 (13)	0.0453 (12)	-0.0255 (11)	-0.0051 (12)	-0.0129 (10)
C1	0.0621 (17)	0.095 (2)	0.0422 (14)	-0.0311 (15)	0.0000 (12)	-0.0173 (13)
C2	0.0642 (16)	0.0669 (17)	0.0564 (15)	-0.0235 (13)	-0.0231 (13)	-0.0092 (12)
C3	0.0475 (13)	0.0402 (12)	0.0459 (12)	-0.0177 (9)	-0.0112 (10)	-0.0077 (9)
C4	0.0400 (11)	0.0369 (11)	0.0427 (12)	-0.0162 (9)	-0.0058 (9)	-0.0085 (9)
C5	0.0501 (13)	0.0444 (12)	0.0442 (13)	-0.0207 (10)	-0.0117 (10)	-0.0061 (9)
C6	0.0611 (14)	0.0384 (12)	0.0389 (12)	-0.0202 (10)	-0.0034 (10)	-0.0079 (9)
C7	0.0465 (13)	0.0485 (13)	0.0523 (14)	-0.0148 (10)	0.0048 (10)	-0.0128 (11)
C8	0.0405 (12)	0.0423 (12)	0.0579 (14)	-0.0137 (9)	-0.0079 (10)	-0.0104 (10)
C9	0.0426 (12)	0.0306 (10)	0.0448 (12)	-0.0135 (8)	-0.0098 (9)	-0.0070 (8)

Geometric parameters (\AA , $^\circ$)

O1—N3	1.222 (3)	C2—H2C	0.9600
O2—N3	1.223 (3)	C2—H2D	0.9600
N1—N2	1.357 (3)	C3—C9	1.389 (3)
N1—C1	1.456 (3)	C4—C5	1.409 (3)
N1—C3	1.350 (3)	C4—C9	1.420 (3)
N2—C4	1.346 (3)	C5—C6	1.366 (3)
N3—C6	1.460 (3)	C5—H5A	0.9300
C1—H1A	0.9600	C6—C7	1.416 (4)
C1—H1B	0.9600	C7—C8	1.354 (3)
C1—H1C	0.9600	C7—H7A	0.9300
C2—C3	1.487 (3)	C8—C9	1.405 (3)
C2—H2B	0.9600	C8—H8A	0.9300
N2—N1—C1	118.6 (2)	N1—C3—C2	124.3 (2)
C3—N1—N2	114.84 (19)	C9—C3—C2	130.3 (2)
C3—N1—C1	126.6 (2)	N2—C4—C5	127.8 (2)
C4—N2—N1	102.99 (17)	N2—C4—C9	111.81 (19)
O1—N3—O2	122.6 (2)	C5—C4—C9	120.4 (2)
O1—N3—C6	118.2 (2)	C6—C5—C4	116.1 (2)
O2—N3—C6	119.2 (2)	C6—C5—H5A	121.9
N1—C1—H1A	109.5	C4—C5—H5A	121.9
N1—C1—H1B	109.5	C5—C6—C7	124.2 (2)
H1A—C1—H1B	109.5	C5—C6—N3	117.7 (2)
N1—C1—H1C	109.5	C7—C6—N3	118.1 (2)
H1A—C1—H1C	109.5	C8—C7—C6	119.7 (2)
H1B—C1—H1C	109.5	C8—C7—H7A	120.1
C3—C2—H2B	109.5	C6—C7—H7A	120.1
C3—C2—H2C	109.5	C7—C8—C9	118.6 (2)

H2B—C2—H2C	109.5	C7—C8—H8A	120.7
C3—C2—H2D	109.5	C9—C8—H8A	120.7
H2B—C2—H2D	109.5	C3—C9—C8	134.1 (2)
H2C—C2—H2D	109.5	C3—C9—C4	105.00 (19)
N1—C3—C9	105.36 (19)	C8—C9—C4	120.9 (2)
C1—N1—N2—C4	179.6 (2)	C2—C3—C9—C8	-2.1 (4)
C3—N1—N2—C4	0.1 (2)	N2—C4—C5—C6	-178.9 (2)
N2—N1—C3—C2	-178.6 (2)	C9—C4—C5—C6	0.9 (3)
N2—N1—C3—C9	0.2 (2)	N2—C4—C9—C3	0.4 (2)
C1—N1—C3—C2	1.9 (4)	N2—C4—C9—C8	-179.25 (18)
C1—N1—C3—C9	-179.3 (2)	C5—C4—C9—C3	-179.41 (19)
N1—N2—C4—C5	179.5 (2)	C5—C4—C9—C8	0.9 (3)
N1—N2—C4—C9	-0.3 (2)	C4—C5—C6—C7	-2.2 (3)
O1—N3—C6—C5	175.6 (2)	C4—C5—C6—N3	179.20 (19)
O1—N3—C6—C7	-3.1 (3)	C5—C6—C7—C8	1.8 (4)
O2—N3—C6—C5	-5.0 (3)	N3—C6—C7—C8	-179.7 (2)
O2—N3—C6—C7	176.4 (2)	C6—C7—C8—C9	0.2 (3)
N1—C3—C9—C4	-0.4 (2)	C7—C8—C9—C3	179.0 (2)
N1—C3—C9—C8	179.3 (2)	C7—C8—C9—C4	-1.4 (3)
C2—C3—C9—C4	178.3 (2)		

Hydrogen-bond geometry (\AA , $^\circ$)

$D\cdots H$	$D—H$	$H\cdots A$	$D\cdots A$	$D—H\cdots A$
C1—H1A ⁱ —O2 ⁱ	0.96	2.58	3.533 (4)	171

Symmetry codes: (i) $-x-1, -y+2, -z$.

supplementary materials

Fig. 1




Fig. 2

