

## 1-(2-Naphthyl)-3-phenylprop-2-en-1-one

Si-Ping Tang,\* Dai-Zhi Kuang, Yong-Lan Feng, Man-Sheng Chen and Wei Li

Key Laboratory of Functional Organometallic Materials, Hengyang Normal University, Hengyang, Hunan 421008, People's Republic of China

Correspondence e-mail: sptang88@163.com


Received 20 June 2009; accepted 1 July 2009

Key indicators: single-crystal X-ray study;  $T = 295$  K; mean  $\sigma(\text{C}-\text{C}) = 0.002$  Å;  $R$  factor = 0.056;  $wR$  factor = 0.174; data-to-parameter ratio = 14.6.

The title compound,  $\text{C}_{19}\text{H}_{14}\text{O}$ , contains two independent molecules with the same *s-cis* conformation for the ketone unit. Both molecules are non-planar with dihedral angles of  $51.9$  (1) and  $48.0$  (1)° between the benzene ring and the naphthalene ring system. In the crystal, neighboring molecules are stabilized by intermolecular  $\text{C}-\text{H}\cdots\pi$  interactions, giving a two-dimensional supramolecular array parallel to the *ab* plane.

## Related literature

For background to chalcone and its derivatives, see: Agrinskaya *et al.* (1999); Indira *et al.* (2002); Opletalova (2000); Pandey *et al.* (2005). For related structures, see: Moorthi *et al.* (2005); Tang *et al.* (2008).



## Experimental

## Crystal data

$\text{C}_{19}\text{H}_{14}\text{O}$   
 $M_r = 258.30$   
 Triclinic,  $P\bar{1}$   
 $a = 9.5878$  (8) Å  
 $b = 9.6111$  (8) Å  
 $c = 15.5358$  (13) Å  
 $\alpha = 98.746$  (2)°  
 $\beta = 91.222$  (2)°

$\gamma = 105.764$  (1)°  
 $V = 1358.8$  (2) Å<sup>3</sup>  
 $Z = 4$   
 Mo  $K\alpha$  radiation  
 $\mu = 0.08$  mm<sup>-1</sup>  
 $T = 295$  K  
 $0.30 \times 0.28 \times 0.20$  mm

## Data collection

Bruker SMART APEX area-detector diffractometer  
 Absorption correction: multi-scan (SADABS; Sheldrick, 1996)  
 $T_{\min} = 0.972$ ,  $T_{\max} = 0.982$

10623 measured reflections  
 5271 independent reflections  
 3483 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.017$

## Refinement

$R[F^2 > 2\sigma(F^2)] = 0.056$   
 $wR(F^2) = 0.174$   
 $S = 1.03$   
 5271 reflections

361 parameters  
 H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 0.24$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.19$  e Å<sup>-3</sup>

## Table 1

Hydrogen-bond geometry (Å, °).

$\text{Cg}1$ ,  $\text{Cg}2$ ,  $\text{Cg}3$ ,  $\text{Cg}5$  and  $\text{Cg}6$  are the centroids of the  $\text{C}1-\text{C}10$ ,  $\text{C}5-\text{C}10$ ,  $\text{C}14-\text{C}19$ ,  $\text{C}20-\text{C}29$  and  $\text{C}24-\text{C}29$  benzene rings, respectively.

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C}13-\text{H}13\cdots\text{O}1$	0.93	2.49	2.814 (2)	101
$\text{C}32-\text{H}32\cdots\text{O}2$	0.93	2.50	2.820 (2)	101
$\text{C}18-\text{H}18\cdots\text{Cg}1^i$	0.93	2.98	3.644	130
$\text{C}15-\text{H}15\cdots\text{Cg}2^{ii}$	0.93	2.94	3.642	134
$\text{C}37-\text{H}37\cdots\text{Cg}3^{iii}$	0.93	2.96	3.610	128
$\text{C}1-\text{H}1\cdots\text{Cg}5^{ii}$	0.93	2.97	3.611	127
$\text{C}6-\text{H}6\cdots\text{Cg}6^{iii}$	0.93	2.92	3.583	130

Symmetry codes: (i)  $-x+1, -y, -z+1$ ; (ii)  $-x+1, -y+1, -z+1$ ; (iii)  $-x, -y+1, -z+1$ .

Data collection: SMART (Bruker, 2002); cell refinement: SAINT (Bruker, 2002); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

We thank the Construct Program for Key Disciplines in Hunan Province and the Foundation of Hunan Province Education Office (No. 08 C178) for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: AT2828).

## References

- Agrinskaya, N. V., Lukoshkin, V. A., Kudryavtsev, V. V., Nosova, G. I., Solovskaya, N. A. & Yakimanski, A. V. (1999). *Phys. Solid State*, **41**, 1914–1917.
- Bruker (2002). SMART and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
- Indira, J., Prakash Karat, P. & Sarojini, B. K. (2002). *J. Cryst. Growth*, **242**, 209–214.
- Moorthi, S. S., Chinnakali, K., Nanjundan, S., Balaji, R. & Fun, H.-K. (2005). *Acta Cryst.* **E61**, o3885–o3887.
- Opletalova, V. (2000). *Ceska Slov. Farm.* **49**, 278–284.
- Pandey, S., Suryawanshi, S. N., Gupta, S. & Srivastava, V. M. L. (2005). *Eur. J. Med. Chem.* **40**, 751–756.
- Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Tang, S.-P., Kuang, D.-Z., Feng, Y.-L., Li, W. & Chen, Z.-M. (2008). *Acta Cryst.* **E64**, o1123.

**supplementary materials**

*Acta Cryst.* (2009). E65, o1785 [ doi:10.1107/S1600536809025483 ]

## 1-(2-Naphthyl)-3-phenylprop-2-en-1-one

S.-P. Tang, D.-Z. Kuang, Y.-L. Feng, M.-S. Chen and W. Li

### Comment

Chalcone and its analogues have been extensively researched because of their facile synthesis and potential applications as excellent non-linear optical materials (Agrinskaya *et al.*, 1999; Indira *et al.*, 2002) and biological activities (Opletalova, 2000; Pandey *et al.*, 2005). Ongoing our efforts on the research of chalcone compounds (Tang *et al.*, 2008), a new compound was here presented.

As shown in Fig.1, the title molecule contains two independent and isostructural molecules, which are non-planar because of the serious tilts between the benzene and naphthalene rings of 51.9 (1) ° and 48.0 (1) ° dihedral angles, respectively. In the two molecules, the ketone units display the same *s-cis*. conformations with the torsion angles of 14.7 (3) ° and 12.3 (3) °, respectively. Meanwhile, the intramolecular C—H···O hydrogen bonds exist within the ketone units, which are also found in the other similar structures (Moorthi *et al.*, 2005; Tang *et al.*, 2008).

In the crystal structure, as shown in Fig.2, neighboring molecules are stacked into a two-dimensional supramolecular layer by intermolecular C—H··· $\pi$  interactions parallel to the *ab* plane.

### Experimental

A mixture of 2-acetonaphthone (1.70 g, 10.0 mmol) and benzaldehyde (1.069 g, 10.0 mmol), sodium hydroxide (0.40 g, 10.0 mmol), ethanol (20 ml) and water (10 ml) was stirred at room temperature for 10 h. After filtering, the resulting precipitate was washed with water and iced ethanol, and further recrystallized from acetonitrile to afford colourless block crystals of the title compound [yield: 1.22 g (47.2 %)].

### Refinement

All H-atoms were positioned geometrically and refined using a riding model with  $d(\text{C-H}) = 0.93 \text{ \AA}$ ,  $U_{\text{iso}} = 1.2U_{\text{eq}}(\text{C})$ .

### Figures




Fig. 1. The title molecule with displacement ellipsoids drawn at the 30% probability level, and H atoms as spheres of arbitrary radius.




Fig. 2. Packing diagram of the title structure showing the C—H...O and C—H... $\pi$  interactions. The H atoms not involved in hydrogen bonding have been omitted for clarity.

## 1-(2-Naphthyl)-3-phenylprop-2-en-1-one

### Crystal data

$C_{19}H_{14}O$

$M_r = 258.30$

Triclinic,  $P\bar{1}$

Hall symbol: -P 1

$a = 9.5878$  (8) Å

$b = 9.6111$  (8) Å

$c = 15.5358$  (13) Å

$\alpha = 98.746$  (2)°

$\beta = 91.222$  (2)°

$\gamma = 105.7640$  (10)°

$V = 1358.8$  (2) Å<sup>3</sup>

$Z = 4$

$F_{000} = 544$

$D_x = 1.263$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 2584 reflections

$\theta = 2.2$ – $26.9$ °

$\mu = 0.08$  mm<sup>-1</sup>

$T = 295$  K

Block, colourless

$0.30 \times 0.28 \times 0.20$  mm

### Data collection

Bruker SMART APEX area-detector diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 295$  K

$\varphi$  and  $\omega$  scans

Absorption correction: multi-scan (SADABS; Sheldrick, 1996)

$T_{\min} = 0.972$ ,  $T_{\max} = 0.982$

10623 measured reflections

5271 independent reflections

3483 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.017$

$\theta_{\text{max}} = 26.0$ °

$\theta_{\text{min}} = 2.2$ °

$h = -11 \rightarrow 11$

$k = -11 \rightarrow 11$

$l = -16 \rightarrow 19$

### Refinement

Refinement on  $F^2$

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.056$

$wR(F^2) = 0.174$

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.0943P)^2 + 0.07P]$$

where  $P = (F_o^2 + 2F_c^2)/3$

$S = 1.03$   $(\Delta/\sigma)_{\max} < 0.001$   
 5271 reflections  $\Delta\rho_{\max} = 0.24 \text{ e } \text{\AA}^{-3}$   
 361 parameters  $\Delta\rho_{\min} = -0.19 \text{ e } \text{\AA}^{-3}$   
 Primary atom site location: structure-invariant direct methods Extinction correction: none

*Special details*

**Geometry.** All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted R-factor wR and goodness of fit S are based on  $F^2$ , conventional R-factors R are based on F, with F set to zero for negative  $F^2$ . The threshold expression of  $F^2 > 2\text{sigma}(F^2)$  is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on  $F^2$  are statistically about twice as large as those based on F, and R- factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.55561 (16)	0.34700 (16)	0.30964 (10)	0.0513 (4)
H1	0.6529	0.3967	0.3072	0.062*
C2	0.50388 (16)	0.31927 (16)	0.38862 (10)	0.0512 (4)
C3	0.35522 (17)	0.24451 (16)	0.39193 (11)	0.0564 (4)
H3	0.3193	0.2228	0.4449	0.068*
C4	0.26443 (17)	0.20422 (17)	0.31847 (11)	0.0569 (4)
H4	0.1666	0.1581	0.3224	0.068*
C5	0.31541 (16)	0.23088 (15)	0.23681 (10)	0.0492 (4)
C6	0.22432 (18)	0.18909 (17)	0.15896 (12)	0.0617 (4)
H6	0.1260	0.1428	0.1611	0.074*
C7	0.2783 (2)	0.2156 (2)	0.08125 (12)	0.0730 (5)
H7	0.2168	0.1879	0.0308	0.088*
C8	0.4258 (2)	0.2841 (2)	0.07654 (12)	0.0735 (5)
H8	0.4618	0.3014	0.0229	0.088*
C9	0.51710 (19)	0.32576 (18)	0.14943 (11)	0.0634 (5)
H9	0.6152	0.3705	0.1451	0.076*
C10	0.46492 (16)	0.30194 (15)	0.23190 (10)	0.0493 (4)
C11	0.60363 (17)	0.37413 (17)	0.46910 (11)	0.0556 (4)
C12	0.57387 (18)	0.29557 (18)	0.54382 (11)	0.0598 (4)
H12	0.5089	0.2024	0.5359	0.072*
C13	0.63803 (17)	0.35515 (17)	0.62227 (11)	0.0548 (4)
H13	0.6999	0.4496	0.6276	0.066*
C14	0.62209 (15)	0.28857 (16)	0.70127 (11)	0.0510 (4)
C15	0.67187 (17)	0.37583 (18)	0.78179 (11)	0.0581 (4)
H15	0.7170	0.4753	0.7845	0.070*
C16	0.65513 (19)	0.3166 (2)	0.85796 (12)	0.0693 (5)
H16	0.6880	0.3765	0.9114	0.083*

## supplementary materials

---

C17	0.59008 (19)	0.1694 (2)	0.85492 (13)	0.0728 (5)
H17	0.5791	0.1297	0.9062	0.087*
C18	0.54137 (18)	0.08120 (19)	0.77577 (13)	0.0678 (5)
H18	0.4979	-0.0184	0.7737	0.081*
C19	0.55654 (17)	0.13930 (17)	0.69964 (12)	0.0596 (4)
H19	0.5228	0.0785	0.6466	0.071*
C20	0.17274 (16)	0.87506 (16)	0.70822 (10)	0.0511 (4)
H20	0.2253	0.9728	0.7122	0.061*
C21	0.11862 (16)	0.79474 (16)	0.62779 (10)	0.0522 (4)
C22	0.03921 (17)	0.64537 (17)	0.62222 (11)	0.0590 (4)
H22	0.0000	0.5909	0.5682	0.071*
C23	0.01976 (17)	0.58105 (17)	0.69474 (11)	0.0592 (4)
H23	-0.0299	0.4822	0.6894	0.071*
C24	0.07335 (16)	0.66125 (16)	0.77786 (11)	0.0509 (4)
C25	0.05131 (18)	0.59894 (18)	0.85505 (12)	0.0639 (5)
H25	0.0025	0.5001	0.8513	0.077*
C26	0.10026 (19)	0.6810 (2)	0.93437 (12)	0.0701 (5)
H26	0.0844	0.6380	0.9843	0.084*
C27	0.17445 (19)	0.8298 (2)	0.94162 (12)	0.0675 (5)
H27	0.2066	0.8853	0.9964	0.081*
C28	0.19983 (17)	0.89384 (17)	0.86894 (11)	0.0584 (4)
H28	0.2505	0.9925	0.8745	0.070*
C29	0.15019 (15)	0.81224 (16)	0.78502 (10)	0.0489 (4)
C30	0.14645 (18)	0.86629 (18)	0.54882 (11)	0.0586 (4)
C31	0.04008 (18)	0.80888 (19)	0.47213 (11)	0.0641 (5)
H31	-0.0495	0.7444	0.4786	0.077*
C32	0.06982 (18)	0.84735 (17)	0.39480 (11)	0.0570 (4)
H32	0.1619	0.9090	0.3907	0.068*
C33	-0.02592 (17)	0.80371 (16)	0.31523 (10)	0.0519 (4)
C34	0.03151 (18)	0.83379 (17)	0.23635 (11)	0.0590 (4)
H34	0.1296	0.8816	0.2351	0.071*
C35	-0.0556 (2)	0.79344 (19)	0.16030 (12)	0.0668 (5)
H35	-0.0160	0.8138	0.1080	0.080*
C36	-0.2010 (2)	0.72311 (18)	0.16094 (12)	0.0664 (5)
H36	-0.2594	0.6956	0.1092	0.080*
C37	-0.25980 (18)	0.69361 (17)	0.23827 (12)	0.0628 (5)
H37	-0.3582	0.6465	0.2388	0.075*
C38	-0.17429 (17)	0.73317 (17)	0.31472 (11)	0.0581 (4)
H38	-0.2153	0.7129	0.3667	0.070*
O1	0.70779 (13)	0.48236 (13)	0.47157 (8)	0.0757 (4)
O2	0.25405 (14)	0.96832 (14)	0.54733 (8)	0.0820 (4)

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C1	0.0439 (8)	0.0479 (8)	0.0599 (10)	0.0099 (6)	0.0067 (7)	0.0072 (7)
C2	0.0504 (9)	0.0477 (8)	0.0545 (9)	0.0131 (7)	0.0059 (7)	0.0058 (7)
C3	0.0554 (9)	0.0568 (9)	0.0559 (10)	0.0126 (7)	0.0119 (8)	0.0105 (7)

C4	0.0468 (9)	0.0538 (9)	0.0677 (11)	0.0100 (7)	0.0082 (8)	0.0096 (8)
C5	0.0478 (8)	0.0433 (8)	0.0577 (10)	0.0160 (6)	0.0038 (7)	0.0060 (7)
C6	0.0529 (9)	0.0570 (9)	0.0723 (12)	0.0134 (7)	-0.0040 (8)	0.0059 (8)
C7	0.0762 (13)	0.0800 (12)	0.0596 (11)	0.0207 (10)	-0.0071 (9)	0.0047 (9)
C8	0.0792 (13)	0.0863 (13)	0.0551 (11)	0.0230 (10)	0.0082 (9)	0.0115 (9)
C9	0.0600 (10)	0.0676 (10)	0.0620 (11)	0.0160 (8)	0.0134 (8)	0.0114 (8)
C10	0.0498 (9)	0.0432 (8)	0.0561 (9)	0.0150 (6)	0.0063 (7)	0.0071 (7)
C11	0.0532 (9)	0.0548 (9)	0.0561 (10)	0.0129 (7)	0.0074 (7)	0.0042 (7)
C12	0.0585 (10)	0.0542 (9)	0.0621 (11)	0.0101 (7)	-0.0001 (8)	0.0062 (8)
C13	0.0482 (9)	0.0516 (9)	0.0629 (10)	0.0111 (7)	0.0015 (7)	0.0095 (7)
C14	0.0414 (8)	0.0522 (9)	0.0596 (10)	0.0134 (7)	0.0013 (7)	0.0090 (7)
C15	0.0518 (9)	0.0571 (9)	0.0629 (11)	0.0118 (7)	-0.0012 (8)	0.0091 (8)
C16	0.0642 (11)	0.0820 (13)	0.0599 (11)	0.0181 (9)	-0.0020 (8)	0.0110 (9)
C17	0.0624 (11)	0.0871 (13)	0.0783 (13)	0.0238 (10)	0.0067 (10)	0.0364 (11)
C18	0.0545 (10)	0.0579 (10)	0.0948 (14)	0.0149 (8)	0.0027 (9)	0.0262 (10)
C19	0.0514 (9)	0.0541 (9)	0.0727 (11)	0.0144 (7)	-0.0026 (8)	0.0101 (8)
C20	0.0449 (8)	0.0442 (8)	0.0612 (10)	0.0084 (6)	0.0009 (7)	0.0064 (7)
C21	0.0451 (8)	0.0546 (9)	0.0555 (10)	0.0136 (7)	0.0012 (7)	0.0053 (7)
C22	0.0555 (9)	0.0529 (9)	0.0624 (10)	0.0122 (7)	-0.0045 (8)	-0.0034 (8)
C23	0.0537 (9)	0.0454 (8)	0.0747 (12)	0.0110 (7)	-0.0006 (8)	0.0045 (8)
C24	0.0418 (8)	0.0465 (8)	0.0653 (10)	0.0139 (6)	0.0016 (7)	0.0095 (7)
C25	0.0592 (10)	0.0544 (9)	0.0807 (12)	0.0146 (8)	0.0026 (9)	0.0219 (9)
C26	0.0708 (12)	0.0732 (12)	0.0702 (12)	0.0192 (9)	0.0039 (9)	0.0255 (10)
C27	0.0689 (11)	0.0751 (12)	0.0568 (10)	0.0190 (9)	-0.0034 (8)	0.0085 (9)
C28	0.0585 (10)	0.0523 (9)	0.0604 (10)	0.0119 (7)	-0.0016 (8)	0.0050 (8)
C29	0.0418 (8)	0.0480 (8)	0.0563 (10)	0.0128 (6)	0.0008 (7)	0.0063 (7)
C30	0.0520 (9)	0.0614 (10)	0.0574 (10)	0.0104 (8)	0.0046 (7)	0.0045 (8)
C31	0.0550 (10)	0.0697 (11)	0.0607 (11)	0.0064 (8)	0.0034 (8)	0.0100 (8)
C32	0.0531 (9)	0.0526 (9)	0.0617 (11)	0.0094 (7)	0.0045 (8)	0.0084 (8)
C33	0.0511 (9)	0.0452 (8)	0.0593 (10)	0.0124 (7)	0.0021 (7)	0.0105 (7)
C34	0.0554 (9)	0.0561 (9)	0.0659 (11)	0.0124 (7)	0.0100 (8)	0.0159 (8)
C35	0.0761 (12)	0.0660 (11)	0.0601 (11)	0.0196 (9)	0.0090 (9)	0.0154 (8)
C36	0.0708 (12)	0.0614 (10)	0.0661 (11)	0.0182 (9)	-0.0082 (9)	0.0097 (8)
C37	0.0507 (9)	0.0572 (10)	0.0794 (12)	0.0122 (7)	-0.0023 (9)	0.0145 (9)
C38	0.0541 (9)	0.0588 (10)	0.0621 (10)	0.0131 (7)	0.0070 (8)	0.0168 (8)
O1	0.0676 (8)	0.0766 (8)	0.0653 (8)	-0.0072 (6)	0.0031 (6)	0.0079 (6)
O2	0.0733 (8)	0.0853 (9)	0.0667 (8)	-0.0125 (7)	-0.0006 (6)	0.0129 (6)

*Geometric parameters (Å, °)*

C1—C2	1.370 (2)	C20—C21	1.374 (2)
C1—C10	1.412 (2)	C20—C29	1.412 (2)
C1—H1	0.9300	C20—H20	0.9300
C2—C3	1.417 (2)	C21—C22	1.419 (2)
C2—C11	1.493 (2)	C21—C30	1.490 (2)
C3—C4	1.360 (2)	C22—C23	1.358 (2)
C3—H3	0.9300	C22—H22	0.9300
C4—C5	1.404 (2)	C23—C24	1.407 (2)
C4—H4	0.9300	C23—H23	0.9300

## supplementary materials

---

C5—C6	1.415 (2)	C24—C25	1.415 (2)
C5—C10	1.420 (2)	C24—C29	1.424 (2)
C6—C7	1.356 (2)	C25—C26	1.356 (2)
C6—H6	0.9300	C25—H25	0.9300
C7—C8	1.397 (3)	C26—C27	1.400 (2)
C7—H7	0.9300	C26—H26	0.9300
C8—C9	1.356 (2)	C27—C28	1.362 (2)
C8—H8	0.9300	C27—H27	0.9300
C9—C10	1.414 (2)	C28—C29	1.415 (2)
C9—H9	0.9300	C28—H28	0.9300
C11—O1	1.2258 (18)	C30—O2	1.2184 (18)
C11—C12	1.471 (2)	C30—C31	1.485 (2)
C12—C13	1.325 (2)	C31—C32	1.324 (2)
C12—H12	0.9300	C31—H31	0.9300
C13—C14	1.460 (2)	C32—C33	1.460 (2)
C13—H13	0.9300	C32—H32	0.9300
C14—C15	1.391 (2)	C33—C34	1.392 (2)
C14—C19	1.397 (2)	C33—C38	1.399 (2)
C15—C16	1.383 (2)	C34—C35	1.374 (2)
C15—H15	0.9300	C34—H34	0.9300
C16—C17	1.375 (3)	C35—C36	1.375 (2)
C16—H16	0.9300	C35—H35	0.9300
C17—C18	1.376 (3)	C36—C37	1.374 (2)
C17—H17	0.9300	C36—H36	0.9300
C18—C19	1.377 (2)	C37—C38	1.371 (2)
C18—H18	0.9300	C37—H37	0.9300
C19—H19	0.9300	C38—H38	0.9300
C2—C1—C10	121.64 (14)	C21—C20—C29	121.38 (14)
C2—C1—H1	119.2	C21—C20—H20	119.3
C10—C1—H1	119.2	C29—C20—H20	119.3
C1—C2—C3	118.81 (14)	C20—C21—C22	119.00 (15)
C1—C2—C11	119.27 (14)	C20—C21—C30	119.14 (14)
C3—C2—C11	121.86 (14)	C22—C21—C30	121.86 (14)
C4—C3—C2	120.73 (15)	C23—C22—C21	120.81 (15)
C4—C3—H3	119.6	C23—C22—H22	119.6
C2—C3—H3	119.6	C21—C22—H22	119.6
C3—C4—C5	121.34 (15)	C22—C23—C24	121.21 (14)
C3—C4—H4	119.3	C22—C23—H23	119.4
C5—C4—H4	119.3	C24—C23—H23	119.4
C4—C5—C6	122.83 (15)	C23—C24—C25	122.72 (14)
C4—C5—C10	118.66 (14)	C23—C24—C29	118.78 (15)
C6—C5—C10	118.50 (15)	C25—C24—C29	118.49 (15)
C7—C6—C5	120.92 (16)	C26—C25—C24	121.06 (15)
C7—C6—H6	119.5	C26—C25—H25	119.5
C5—C6—H6	119.5	C24—C25—H25	119.5
C6—C7—C8	120.41 (17)	C25—C26—C27	120.53 (17)
C6—C7—H7	119.8	C25—C26—H26	119.7
C8—C7—H7	119.8	C27—C26—H26	119.7
C9—C8—C7	120.66 (17)	C28—C27—C26	120.38 (16)



C9—C8—H8	119.7	C28—C27—H27	119.8
C7—C8—H8	119.7	C26—C27—H27	119.8
C8—C9—C10	120.75 (16)	C27—C28—C29	120.85 (15)
C8—C9—H9	119.6	C27—C28—H28	119.6
C10—C9—H9	119.6	C29—C28—H28	119.6
C1—C10—C9	122.51 (14)	C20—C29—C28	122.54 (14)
C1—C10—C5	118.75 (14)	C20—C29—C24	118.78 (14)
C9—C10—C5	118.74 (15)	C28—C29—C24	118.68 (15)
O1—C11—C12	121.57 (15)	O2—C30—C31	121.21 (16)
O1—C11—C2	119.81 (15)	O2—C30—C21	120.69 (15)
C12—C11—C2	118.62 (13)	C31—C30—C21	118.09 (14)
C13—C12—C11	121.17 (15)	C32—C31—C30	121.54 (15)
C13—C12—H12	119.4	C32—C31—H31	119.2
C11—C12—H12	119.4	C30—C31—H31	119.2
C12—C13—C14	127.04 (15)	C31—C32—C33	127.23 (15)
C12—C13—H13	116.5	C31—C32—H32	116.4
C14—C13—H13	116.5	C33—C32—H32	116.4
C15—C14—C19	118.01 (16)	C34—C33—C38	118.17 (15)
C15—C14—C13	119.50 (14)	C34—C33—C32	119.13 (14)
C19—C14—C13	122.48 (15)	C38—C33—C32	122.70 (15)
C16—C15—C14	120.84 (16)	C35—C34—C33	120.51 (16)
C16—C15—H15	119.6	C35—C34—H34	119.7
C14—C15—H15	119.6	C33—C34—H34	119.7
C17—C16—C15	120.22 (17)	C34—C35—C36	120.51 (17)
C17—C16—H16	119.9	C34—C35—H35	119.7
C15—C16—H16	119.9	C36—C35—H35	119.7
C16—C17—C18	119.73 (18)	C37—C36—C35	119.78 (17)
C16—C17—H17	120.1	C37—C36—H36	120.1
C18—C17—H17	120.1	C35—C36—H36	120.1
C17—C18—C19	120.50 (16)	C38—C37—C36	120.42 (16)
C17—C18—H18	119.8	C38—C37—H37	119.8
C19—C18—H18	119.8	C36—C37—H37	119.8
C18—C19—C14	120.70 (16)	C37—C38—C33	120.61 (16)
C18—C19—H19	119.7	C37—C38—H38	119.7
C14—C19—H19	119.7	C33—C38—H38	119.7
C10—C1—C2—C3	-0.5 (2)	C29—C20—C21—C22	-0.4 (2)
C10—C1—C2—C11	-177.88 (13)	C29—C20—C21—C30	-179.62 (13)
C1—C2—C3—C4	-1.7 (2)	C20—C21—C22—C23	-1.7 (2)
C11—C2—C3—C4	175.60 (14)	C30—C21—C22—C23	177.56 (14)
C2—C3—C4—C5	2.1 (2)	C21—C22—C23—C24	2.1 (3)
C3—C4—C5—C6	179.33 (14)	C22—C23—C24—C25	178.04 (15)
C3—C4—C5—C10	-0.3 (2)	C22—C23—C24—C29	-0.5 (2)
C4—C5—C6—C7	-179.29 (15)	C23—C24—C25—C26	-177.69 (15)
C10—C5—C6—C7	0.3 (2)	C29—C24—C25—C26	0.9 (2)
C5—C6—C7—C8	0.4 (3)	C24—C25—C26—C27	-0.2 (3)
C6—C7—C8—C9	-0.3 (3)	C25—C26—C27—C28	-0.7 (3)
C7—C8—C9—C10	-0.6 (3)	C26—C27—C28—C29	0.9 (3)
C2—C1—C10—C9	-178.13 (14)	C21—C20—C29—C28	-177.50 (14)
C2—C1—C10—C5	2.3 (2)	C21—C20—C29—C24	1.9 (2)

## supplementary materials

C8—C9—C10—C1	-178.34 (15)	C27—C28—C29—C20	179.19 (14)
C8—C9—C10—C5	1.2 (2)	C27—C28—C29—C24	-0.2 (2)
C4—C5—C10—C1	-1.9 (2)	C23—C24—C29—C20	-1.5 (2)
C6—C5—C10—C1	178.51 (13)	C25—C24—C29—C20	179.91 (13)
C4—C5—C10—C9	178.52 (14)	C23—C24—C29—C28	177.96 (14)
C6—C5—C10—C9	-1.1 (2)	C25—C24—C29—C28	-0.7 (2)
C1—C2—C11—O1	26.3 (2)	C20—C21—C30—O2	27.6 (2)
C3—C2—C11—O1	-150.94 (16)	C22—C21—C30—O2	-151.59 (17)
C1—C2—C11—C12	-153.52 (15)	C20—C21—C30—C31	-152.83 (15)
C3—C2—C11—C12	29.2 (2)	C22—C21—C30—C31	27.9 (2)
O1—C11—C12—C13	14.7 (3)	O2—C30—C31—C32	12.3 (3)
C2—C11—C12—C13	-165.43 (15)	C21—C30—C31—C32	-167.26 (15)
C11—C12—C13—C14	-178.11 (14)	C30—C31—C32—C33	-177.63 (15)
C12—C13—C14—C15	-166.36 (16)	C31—C32—C33—C34	-168.95 (16)
C12—C13—C14—C19	12.8 (3)	C31—C32—C33—C38	11.4 (3)
C19—C14—C15—C16	-0.8 (2)	C38—C33—C34—C35	-0.7 (2)
C13—C14—C15—C16	178.42 (14)	C32—C33—C34—C35	179.67 (15)
C14—C15—C16—C17	0.7 (3)	C33—C34—C35—C36	0.2 (3)
C15—C16—C17—C18	-0.1 (3)	C34—C35—C36—C37	0.3 (3)
C16—C17—C18—C19	-0.4 (3)	C35—C36—C37—C38	-0.3 (3)
C17—C18—C19—C14	0.3 (3)	C36—C37—C38—C33	-0.2 (3)
C15—C14—C19—C18	0.3 (2)	C34—C33—C38—C37	0.7 (2)
C13—C14—C19—C18	-178.88 (14)	C32—C33—C38—C37	-179.68 (14)

### Hydrogen-bond geometry ( $\text{\AA}$ , $^\circ$ )

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C13—H13 $\cdots$ O1	0.93	2.49	2.814 (2)	101
C32—H32 $\cdots$ O2	0.93	2.50	2.820 (2)	101
C18—H18 $\cdots$ Cg1 <sup>i</sup>	0.93	2.98	3.644	130
C15—H15 $\cdots$ Cg2 <sup>ii</sup>	0.93	2.94	3.642	134
C37—H37 $\cdots$ Cg3 <sup>iii</sup>	0.93	2.96	3.610	128
C1—H1 $\cdots$ Cg5 <sup>ii</sup>	0.93	2.97	3.611	127
C6—H6 $\cdots$ Cg6 <sup>iii</sup>	0.93	2.92	3.583	130

Symmetry codes: (i)  $-x+1, -y, -z+1$ ; (ii)  $-x+1, -y+1, -z+1$ ; (iii)  $-x, -y+1, -z+1$ .

Fig. 1




Fig. 2

