

## 3-Hydroxy-3-(methoxycarbonyl)-pentanedioic acid

Lawal Aliyu,<sup>a</sup> Nornisah Mohamed,<sup>a‡</sup> Ching Kheng Quah<sup>b§</sup> and Hoong-Kun Fun<sup>b\*,¶</sup>

<sup>a</sup>School of Pharmaceutical Sciences, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia, and <sup>b</sup>X-ray Crystallography Unit, School of Physics, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia  
Correspondence e-mail: hkfun@usm.my


Received 30 June 2009; accepted 1 July 2009

Key indicators: single-crystal X-ray study;  $T = 100\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.001\text{ \AA}$ ;  $R$  factor = 0.041;  $wR$  factor = 0.107; data-to-parameter ratio = 26.4.

In the title compound,  $\text{C}_7\text{H}_{10}\text{O}_7$ , the aliphatic chain is approximately planar [maximum deviation = 0.013 (1)  $\text{\AA}$ ] and makes a dihedral angle of 78.75 (7) $^\circ$  with the methoxycarbonyl group. In the crystal, molecules are linked via intermolecular  $\text{O}-\text{H}\cdots\text{O}$  and  $\text{C}-\text{H}\cdots\text{O}$  hydrogen bonds into sheets parallel to (100). In the sheet,  $\text{O}-\text{H}\cdots\text{O}$  hydrogen bonds generate  $R_2^2(9)$  and  $R_2^2(8)$  ring motifs.

### Related literature

For hydrogen-bond motifs, see: Bernstein *et al.* (1995). For bond-length data, see: Allen *et al.* (1987). For the stability of the temperature controller used for the data collection, see: Cosier & Glazer (1986).



### Experimental

#### Crystal data

$\text{C}_7\text{H}_{10}\text{O}_7$   
 $M_r = 206.15$

Orthorhombic,  $Pbca$   
 $a = 12.7110 (4)\text{ \AA}$

$b = 5.8323 (2)\text{ \AA}$   
 $c = 23.8844 (7)\text{ \AA}$   
 $V = 1770.65 (10)\text{ \AA}^3$   
 $Z = 8$

Mo  $K\alpha$  radiation  
 $\mu = 0.14\text{ mm}^{-1}$   
 $T = 100\text{ K}$   
 $0.54 \times 0.18 \times 0.11\text{ mm}$

#### Data collection

Bruker SMART APEXII CCD area-detector diffractometer  
Absorption correction: multi-scan (*SADABS*; Bruker, 2005)  
 $T_{\min} = 0.927$ ,  $T_{\max} = 0.985$

54585 measured reflections  
4404 independent reflections  
4059 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.030$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.041$   
 $wR(F^2) = 0.107$   
 $S = 1.12$   
4404 reflections

167 parameters  
All H-atom parameters refined  
 $\Delta\rho_{\text{max}} = 0.56\text{ e \AA}^{-3}$   
 $\Delta\rho_{\text{min}} = -0.25\text{ e \AA}^{-3}$

**Table 1**  
Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{O}2-\text{H1O}2\cdots\text{O}6^i$	0.84 (2)	1.85 (2)	2.6838 (8)	173 (2)
$\text{O}3-\text{H1O}3\cdots\text{O}1^{ii}$	0.87 (2)	2.01 (2)	2.8549 (8)	163 (2)
$\text{O}4-\text{H1O}4\cdots\text{O}5^{iii}$	0.91 (2)	1.73 (2)	2.6391 (9)	176 (2)
$\text{C}4-\text{H4A}\cdots\text{O}5^{iv}$	0.99 (1)	2.53 (1)	3.4035 (9)	147 (1)

Symmetry codes: (i)  $-x, y + \frac{1}{2}, -z + \frac{1}{2}$ ; (ii)  $-x, y - \frac{1}{2}, -z + \frac{1}{2}$ ; (iii)  $-x, -y, -z + 1$ ; (iv)  $-x, -y + 1, -z + 1$ .

Data collection: *APEX2* (Bruker, 2005); cell refinement: *SAINT* (Bruker, 2005); data reduction: *SAINT*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL* and *PLATON* (Spek, 2009).

LA and NM gratefully acknowledge funding from Universiti Sains Malaysia (USM) under the University Research Grant (No. 1001/PFARMASI/815025). HKF and CKQ thank USM for the Research University Golden Goose Grant (No. 1001/PFIZIK/811012). CKQ thanks USM for a Research Fellowship.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CI2844).

### References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Bruker (2005). *APEX2*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Cosier, J. & Glazer, A. M. (1986). *J. Appl. Cryst.* **19**, 105–107.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.

<sup>‡</sup> Additional correspondence author, e-mail: nornisah@usm.my.  
<sup>§</sup> Thomson Reuters ResearcherID: A-5525-2009.

<sup>¶</sup> Thomson Reuters ResearcherID: A-3561-2009.

## **supplementary materials**

*Acta Cryst.* (2009). E65, o1843 [doi:10.1107/S1600536809025598]

### 3-Hydroxy-3-(methoxycarbonyl)pentanedioic acid

L. Aliyu, N. Mohamed, C. K. Quah and H.-K. Fun

#### Comment

Research in natural products often results in the discovery of novel and interesting compounds. Chemical constituents from mango have been found to exhibit bioactivities against certain in silico disease models. Herein, we report the crystal structure of the title compound which was isolated from the mango extract.

The bond lengths (Allen *et al.*, 1987) and angles in the molecule (Fig. 1) are within normal ranges. Atoms C1, C2, C3 and C4 are approximately planar, with a maximum deviation of 0.013 (1) Å for atom C2. This plane makes a dihedral angle of 78.75 (4)° with the mean plane of methoxycarbonyl group (C6/C7/O6/O7).

In the solid state (Fig. 2), the molecules are linked *via* intermolecular O2—H1O2···O6 and O3—H1O3···O1 hydrogen bonds to generate  $R_2^2(9)$  ring motifs (Bernstein *et al.*, 1995) (Table 1) and pairs of intermolecular O4—H1O4···O5 hydrogen bonds form  $R_2^2(8)$  ring motifs. The crystal structure is further stabilized by intermolecular C4—H4A···O5 hydrogen bonds. The molecules are linked by these hydrogen bonds to form layers parallel to the (100).

#### Experimental

The title compound was extracted by Soxhlet extraction method using methanol from oven-dried mangoes. The dried extract was dissolved in water and fractionated using different solvents. The ethyl acetate fraction was evaporated using rotary evaporator and the residue was purified using column chromatography (40% ethyl acetate: 60% n-hexane) to give crystals after washing with ethyl acetate. The crystals were later found to be suitable for X-ray analysis.

#### Refinement

All H atoms were located in a difference Fourier map and refined freely.

#### Figures




Fig. 1. The molecular structure of the title compound, showing 50% probability displacement ellipsoids and the atom-numbering scheme.

## supplementary materials

---




Fig. 2. The crystal packing of the title compound, viewed along the  $a$  axis. H atoms not involved in hydrogen bonds (dashed lines) have been omitted for clarity.

### 3-Hydroxy-3-(methoxycarbonyl)pentanedioic acid

#### Crystal data

C <sub>7</sub> H <sub>10</sub> O <sub>7</sub>	$F_{000} = 864$
$M_r = 206.15$	$D_x = 1.547 \text{ Mg m}^{-3}$
Orthorhombic, <i>Pbca</i>	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2ac 2ab	Cell parameters from 9850 reflections
$a = 12.7110 (4) \text{ \AA}$	$\theta = 3.2\text{--}36.7^\circ$
$b = 5.8323 (2) \text{ \AA}$	$\mu = 0.14 \text{ mm}^{-1}$
$c = 23.8844 (7) \text{ \AA}$	$T = 100 \text{ K}$
$V = 1770.65 (10) \text{ \AA}^3$	Plate, colourless
$Z = 8$	$0.54 \times 0.18 \times 0.11 \text{ mm}$

#### Data collection

Bruker SMART APEXII CCD area-detector diffractometer	4404 independent reflections
Radiation source: fine-focus sealed tube	4059 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.030$
$T = 100 \text{ K}$	$\theta_{\text{max}} = 36.7^\circ$
$\varphi$ and $\omega$ scans	$\theta_{\text{min}} = 2.3^\circ$
Absorption correction: multi-scan ( <i>SADABS</i> ; Bruker, 2005)	$h = -21 \rightarrow 21$
$T_{\text{min}} = 0.927$ , $T_{\text{max}} = 0.985$	$k = -8 \rightarrow 9$
54585 measured reflections	$l = -39 \rightarrow 39$

#### Refinement

Refinement on $F^2$	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.041$	All H-atom parameters refined
$wR(F^2) = 0.107$	$w = 1/[\sigma^2(F_o^2) + (0.0517P)^2 + 0.4849P]$
$S = 1.12$	where $P = (F_o^2 + 2F_c^2)/3$
4404 reflections	$(\Delta/\sigma)_{\text{max}} = 0.001$
167 parameters	$\Delta\rho_{\text{max}} = 0.56 \text{ e \AA}^{-3}$
	$\Delta\rho_{\text{min}} = -0.24 \text{ e \AA}^{-3}$

Primary atom site location: structure-invariant direct methods      Extinction correction: none

### *Special details*

**Experimental.** The crystal was placed in the cold stream of an Oxford Cyrosystems Cobra open-flow nitrogen cryostat (Cosier & Glazer, 1986) operating at 100.0 (1) K.

**Geometry.** All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted R-factor wR and goodness of fit S are based on  $F^2$ , conventional R-factors R are based on F, with F set to zero for negative  $F^2$ . The threshold expression of  $F^2 > 2\text{sigma}(F^2)$  is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on  $F^2$  are statistically about twice as large as those based on F, and R- factors based on ALL data will be even larger.

### *Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.02576 (5)	0.61486 (11)	0.25826 (2)	0.01728 (11)
O2	0.17885 (5)	0.80419 (12)	0.24936 (3)	0.01938 (12)
O3	0.08405 (4)	0.23143 (9)	0.34206 (2)	0.01268 (10)
O4	0.12975 (5)	0.07178 (11)	0.47512 (3)	0.01837 (12)
O5	-0.02721 (4)	0.24507 (10)	0.46792 (2)	0.01487 (11)
O6	-0.11179 (4)	0.42068 (10)	0.35333 (2)	0.01375 (10)
O7	-0.04401 (4)	0.74583 (10)	0.38809 (3)	0.01586 (11)
C1	0.10969 (5)	0.67852 (13)	0.27732 (3)	0.01260 (11)
C2	0.14594 (5)	0.62879 (13)	0.33592 (3)	0.01235 (11)
C3	0.07801 (5)	0.45077 (12)	0.36661 (3)	0.01026 (11)
C4	0.12280 (5)	0.42501 (12)	0.42612 (3)	0.01211 (11)
C5	0.06767 (5)	0.23834 (12)	0.45814 (3)	0.01218 (11)
C6	-0.03676 (5)	0.53355 (12)	0.36864 (3)	0.01112 (11)
C7	-0.14757 (6)	0.85003 (15)	0.38333 (4)	0.02113 (15)
H2A	0.1457 (11)	0.771 (2)	0.3557 (6)	0.020 (3)*
H2B	0.2200 (10)	0.580 (2)	0.3349 (6)	0.021 (3)*
H4A	0.1116 (10)	0.567 (2)	0.4477 (6)	0.016 (3)*
H4B	0.1959 (10)	0.390 (2)	0.4241 (5)	0.015 (3)*
H7A	-0.1957 (12)	0.773 (3)	0.4079 (7)	0.033 (4)*
H7B	-0.1406 (11)	1.010 (3)	0.3934 (6)	0.023 (3)*
H7C	-0.1741 (14)	0.827 (3)	0.3455 (7)	0.039 (4)*
H1O2	0.1539 (13)	0.834 (3)	0.2177 (7)	0.038 (4)*
H1O3	0.0466 (12)	0.225 (3)	0.3117 (7)	0.032 (4)*
H1O4	0.0934 (14)	-0.032 (4)	0.4959 (8)	0.046 (5)*

### *Atomic displacement parameters ( $\text{\AA}^2$ )*

$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
----------	----------	----------	----------	----------	----------

## supplementary materials

---

O1	0.0175 (2)	0.0178 (3)	0.0166 (2)	-0.00488 (19)	-0.00472 (18)	0.00421 (19)
O2	0.0153 (2)	0.0256 (3)	0.0172 (3)	-0.0040 (2)	-0.00089 (18)	0.0107 (2)
O3	0.0149 (2)	0.0102 (2)	0.0130 (2)	0.00161 (16)	-0.00136 (16)	-0.00159 (16)
O4	0.0142 (2)	0.0180 (3)	0.0229 (3)	0.00341 (19)	0.00330 (19)	0.0089 (2)
O5	0.0123 (2)	0.0159 (2)	0.0164 (2)	0.00103 (17)	0.00198 (16)	0.00301 (18)
O6	0.0111 (2)	0.0156 (2)	0.0145 (2)	-0.00290 (17)	-0.00018 (16)	-0.00210 (17)
O7	0.0116 (2)	0.0115 (2)	0.0245 (3)	0.00166 (17)	-0.00251 (18)	-0.00385 (19)
C1	0.0128 (2)	0.0115 (3)	0.0135 (3)	0.0001 (2)	0.00038 (19)	0.0027 (2)
C2	0.0111 (2)	0.0132 (3)	0.0128 (3)	-0.0015 (2)	-0.00051 (19)	0.0029 (2)
C3	0.0101 (2)	0.0097 (2)	0.0110 (2)	-0.00015 (19)	-0.00016 (18)	0.00032 (19)
C4	0.0120 (2)	0.0133 (3)	0.0110 (3)	-0.0015 (2)	-0.00097 (19)	0.0013 (2)
C5	0.0129 (2)	0.0136 (3)	0.0100 (2)	0.0004 (2)	0.00019 (18)	0.0006 (2)
C6	0.0110 (2)	0.0106 (3)	0.0118 (3)	-0.00047 (19)	-0.00006 (18)	0.0002 (2)
C7	0.0146 (3)	0.0169 (3)	0.0319 (4)	0.0051 (3)	-0.0034 (3)	-0.0041 (3)

### Geometric parameters ( $\text{\AA}$ , $^\circ$ )

O1—C1	1.2179 (9)	C2—C3	1.5365 (9)
O2—C1	1.3251 (9)	C2—H2A	0.956 (14)
O2—H1O2	0.838 (17)	C2—H2B	0.984 (13)
O3—C3	1.4094 (9)	C3—C6	1.5373 (9)
O3—H1O3	0.869 (16)	C3—C4	1.5386 (9)
O4—C5	1.3156 (9)	C4—C5	1.5037 (10)
O4—H1O4	0.91 (2)	C4—H4A	0.984 (13)
O5—C5	1.2290 (9)	C4—H4B	0.953 (13)
O6—C6	1.2152 (8)	C7—H7A	0.960 (16)
O7—C6	1.3256 (9)	C7—H7B	0.966 (15)
O7—C7	1.4543 (10)	C7—H7C	0.973 (17)
C1—C2	1.5020 (10)		
C1—O2—H1O2	108.5 (12)	C5—C4—C3	111.60 (6)
C3—O3—H1O3	111.0 (11)	C5—C4—H4A	106.0 (8)
C5—O4—H1O4	110.8 (12)	C3—C4—H4A	110.4 (8)
C6—O7—C7	115.21 (6)	C5—C4—H4B	108.9 (8)
O1—C1—O2	124.15 (7)	C3—C4—H4B	109.6 (8)
O1—C1—C2	123.93 (6)	H4A—C4—H4B	110.4 (11)
O2—C1—C2	111.90 (6)	O5—C5—O4	123.61 (7)
C1—C2—C3	113.75 (6)	O5—C5—C4	122.06 (6)
C1—C2—H2A	107.0 (8)	O4—C5—C4	114.32 (6)
C3—C2—H2A	110.5 (8)	O6—C6—O7	123.84 (6)
C1—C2—H2B	109.1 (8)	O6—C6—C3	124.41 (6)
C3—C2—H2B	110.7 (8)	O7—C6—C3	111.74 (6)
H2A—C2—H2B	105.5 (12)	O7—C7—H7A	109.5 (10)
O3—C3—C2	112.60 (5)	O7—C7—H7B	107.5 (8)
O3—C3—C6	110.48 (5)	H7A—C7—H7B	111.0 (13)
C2—C3—C6	109.64 (5)	O7—C7—H7C	109.2 (10)
O3—C3—C4	106.00 (5)	H7A—C7—H7C	106.4 (14)
C2—C3—C4	107.38 (5)	H7B—C7—H7C	113.3 (13)
C6—C3—C4	110.65 (5)		
O1—C1—C2—C3	11.29 (11)	C3—C4—C5—O4	120.28 (7)

O2—C1—C2—C3	−170.15 (6)	C7—O7—C6—O6	−7.76 (11)
C1—C2—C3—O3	65.49 (8)	C7—O7—C6—C3	170.92 (7)
C1—C2—C3—C6	−57.93 (8)	O3—C3—C6—O6	2.90 (9)
C1—C2—C3—C4	−178.20 (6)	C2—C3—C6—O6	127.56 (7)
O3—C3—C4—C5	−54.29 (7)	C4—C3—C6—O6	−114.18 (8)
C2—C3—C4—C5	−174.86 (6)	O3—C3—C6—O7	−175.77 (6)
C6—C3—C4—C5	65.51 (7)	C2—C3—C6—O7	−51.11 (8)
C3—C4—C5—O5	−60.67 (9)	C4—C3—C6—O7	67.15 (7)

*Hydrogen-bond geometry (Å, °)*

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
O2—H1O2···O6 <sup>i</sup>	0.84 (2)	1.85 (2)	2.6838 (8)	173 (2)
O3—H1O3···O1 <sup>ii</sup>	0.87 (2)	2.01 (2)	2.8549 (8)	163 (2)
O4—H1O4···O5 <sup>iii</sup>	0.91 (2)	1.73 (2)	2.6391 (9)	176 (2)
C4—H4A···O5 <sup>iv</sup>	0.99 (1)	2.53 (1)	3.4035 (9)	147 (1)

Symmetry codes: (i)  $-x, y+1/2, -z+1/2$ ; (ii)  $-x, y-1/2, -z+1/2$ ; (iii)  $-x, -y, -z+1$ ; (iv)  $-x, -y+1, -z+1$ .

## **supplementary materials**

---

**Fig. 1**




Fig. 2

